



A summary list of fossil spiders and their relatives

compiled by

**Jason A. Dunlop (Berlin), David Penney (Manchester)
& Denise Jekel (Berlin)**

with additional contributions from Lyall I. Anderson, Simon J. Braddy,
James C. Lamsdell, Paul A. Selden & O. Erik Tetlie



A summary list of fossil spiders and their relatives

compiled by

Jason A. Dunlop (Berlin), David Penney (Manchester) & Denise Jekel (Berlin)

with additional contributions from Lyall I. Anderson, Simon J. Braddy,
James C. Lamsdell, Paul A. Selden & O. Erik Tetlie

Suggested citation:

Dunlop, J. A., Penney, D. & Jekel, D. 2015. A summary list of fossil spiders and their relatives. In *World Spider Catalog*. Natural History Museum Bern, online at <http://wsc.nmbe.ch>, version 15.5, accessed on {date of access}.

Last updated: 09.01.2015

INTRODUCTION

Fossil spiders have not been fully catalogued since Bonnet's *Bibliographia Araneorum* and are not included in the current *World Spider Catalog*. Since Bonnet's time there has been considerable progress in our understanding of the fossil record of spiders – and other arachnids – and numerous new taxa have been described. For an overview see Dunlop & Penney (2012). Spiders remain the single largest fossil group, but our aim here is to offer a summary list of all fossil Chelicerata in their current systematic position; as a first step towards the eventual goal of combining fossil and Recent data within a single arachnological resource.

To integrate our data as smoothly as possible with standards used for living spiders, our list for Araneae follows the names and sequence of families adopted in the previous Platnick Catalog. For this reason some of the family groups proposed in Wunderlich's (2004, 2008, 2012) monographs of amber and copal spiders are not reflected here, and we encourage the reader to consult these studies for details and alternative opinions. Extinct families have been inserted in the position which we hope best reflects their probable affinities. For other arachnid groups we have largely followed the nomenclature and family sequences adopted in other online or printed summaries; for example Victor Fet *et al.*'s work on scorpions, Mark Harvey's catalogues of pseudoscorpions and the 'minor' orders – all of which also list the fossils – Adriano Kury's harvestman overviews and the third edition of the Manual of Acarology for mites. For all groups, genus and species names were compiled from established lists and cross-referenced against the primary literature.

We aim to reflect the latest published opinions on the taxonomy of fossil species. A caveat here is that some synonomies and transfers proposed in the literature were only provisional or tentative in nature. At times we were forced to interpret whether a formal nomenclatural change had actually been made, and we have tried to accommodate these difficulties as best as possible. We should also stress that many historical fossil types require revision. Older species names assigned to common, modern genera such as *Araneus*, *Clubiona* or *Linyphia* among the spiders, should be treated with caution. The list has been extended to include Recent species – particularly some spiders and numerous oribatid mites – found as (sub)fossils. These are generally specimens of Quaternary age found in copal, or recovered from peats or archeological sites.

We have provided references for the first descriptions of all the fossil species, and where possible we have added the relevant taxonomic literature for all the taxon names which we mention here. We should, however, note that for some groups (especially mites) recovering the correct author and date for higher taxa proved challenging, and we hope in future releases to be able to clarify these names and augment the reference list accordingly. Formal synonymy lists for the fossil species are being compiled and that which we have for individual taxa can be made available upon request upon a ‘fair use’ basis. As with any project of this size, we cannot guarantee the accuracy of all these entries and we encourage readers to forward omissions or corrections to jason.dunlop@mfn-berlin.de or David.Penney@manchester.ac.uk.

PRINCIPAL CHANGES SINCE THE LAST UPDATE

Recent work included descriptions of new Mesozoic spiders: Palpimanoidea from the Jurassic of Germany and Theridosomatidae from the Cretaceous of France. Some fossil spiders needed to be transferred to the recently accepted families Eutichuridae, Trachelidae and Phrurolithidae. Additionally, the oldest opilioacarid mite was described from Cretaceous Myanmar amber as well as two amber whip spiders (from Myanmar and India respectively). A new horseshoe crab assigned to *Limulus* was described from the Jurassic of Poland. The controversial Burgess Shale fossil *Sanctacaris* has been tentatively included under Chelicerata based on a recent phylogeny.

ACKNOWLEDGMENTS

We are very grateful to Wolfgang Nentwig and the Bern team for agreeing to host this list as an appendix to the Catalog, to Paul Selden for encouragement and support and to those colleagues who have advised us on oversights and/or provided further literature.

EXPLANATIONS

- † indicates an entirely extinct genus, family or other higher taxon
- all species listed assumed to be extinct unless marked [Recent]
- * indicates the type species of (fossil) genera

Stratigraphical abbreviations:

pC = Precambrian, C = Cambrian, O = Ordovician, S = Silurian,

D = Devonian, C = Carboniferous, P = Permian

Tr = Triassic, J = Jurassic, K = Cretaceous

Pa = Palaeogene, Ne = Neogene, Qt = Quaternary

PYCGONOIDA

11 currently valid species of fossil sea spider

- note that in some modern phylogenies the Palaeozoic genera resolve *within* the crown group

PYCGONOIDA Latreille, 1810 Cambrian – Recent

= ARACHNOPODA Dana, 1853

† **Cambropycnogon Waloszek & Dunlop, 2002** Cambrian

1. *Cambropycnogon klausmuelleri* Waloszek & Dunlop, 2002* € ‘Orsten’, Sweden

Pycnogonid affinities of this taxon were questioned by Bamber (2007)

† **Haliestes Siveter, Sutton, Briggs & Siveter, 2004** Silurian

2. *Haliestes dasos* Siveter, Sutton, Briggs & Siveter, 2004* S Herefordshire Lgst.

† **Flagellopantopus Poschmann & Dunlop, 2006** Devonian

3. *Flagellopantopus blocki* Poschmann & Dunlop, 2006* D Hunsruckschiefer

† **Palaeomarachne Rudkin, Cuggy, Young & Thompson, 2013** Ordovician

4. *Palaeomarachne granulata* Rudkin, Cuggy, Young & Thompson, 2013* O Manitoba, Canada

† **Pentapantopus Kühl, Poschmann & Rust, 2013** Devonian

5. *Pentapantopus vogteli* Kühl, Poschmann & Rust, 2013* D Hunsruckschiefer

† **PALAEOSOPHIDAE Dubinin, 1957** Devonian

† **Palaeoisopus Broili, 1928** Devonian

6. *Palaeoisopus problematicus* Broili, 1928* D Hunsruckschiefer

† **PALAEOPANTOPODIDAE Broili, 1930** Devonian

† **Palaeopantopus Broili, 1928** Devonian

7. *Palaeopantopus maucherii* Broili, 1928* D Hunsruckschiefer

PANTOPODA Gerstaecker, 1863 Devonian – Recent

= PEGMATA Fry, 1978

family uncertain

† **Palaeothea Bergström, Stürmer & Winter, 1980** Devonian

8. *Palaeothea devonica* Bergström, Stürmer & Winter, 1980* D Hunsruckschiefer

AUSTRODECIDAE Stock, 1954 Recent

no fossil record

PYCGONIDAE Wilson, 1878 Recent

no fossil record

- COLOSSENDEIDAE Hoek, 1881** ?Jurassic – Recent
- = PASITHOIDAE Sars, 1891
 - = RHOPALORHYNCHIDAE Fry, 1978
- † **Colossopantopodus Charbonnier, Vannier & Riou, 2007** Jurassic
9. *Colossopantopodus boissinensis* Charbonnier, Vannier & Riou, 2007* . J La Voulte-sur-Rhône
tentative referal
- AMMOTHEIDAE Dohrn, 1881** ?Jurassic – Recent
- = EURYCIDIIDAE Sars, 1891
 - = OORHYNCHIDAE Schimkewitsch, 1913
 - = TANYSTYLIDAE Schimkewitsch, 1913
 - = AMMOTHELLIDAE Fry, 1978
 - = EPHYROGYMNIDAE Fry, 1978
 - = PARANYMPHONIDAE Fry, 1978
 - = SERICOSURIDAE Fry, 1978
 - = TRYGAEIDAE Fry, 1978
- † **Palaeopycnogonides Charbonnier, Vannier & Riou, 2007** Jurassic
10. *Palaeopycnogonides gracilis* Charbonnier, Vannier & Riou, 2007* J La Voulte-sur-Rhône
tentative referal
- CALLIPALLENIIDAE Hilton, 1942** Recent
- = PALLENIIDAE Wilson, 1878 [Pallene is a preoccupied genus]
 - = CHEILAPALLENIIDAE Fry, 1978
 - = CLAVIGEROPALLENIIDAE Fry, 1978
 - = HANNONIIDAE Fry, 1978
 - = METAPALLENIIDAE Fry, 1978
 - = QUEUBIDAE Fry, 1978
 - = STYLOPALLENIIDAE Fry, 1978
- no fossil record
- NYMPHONIDAE Wilson, 1878** Recent
- no fossil record
- PALLENOPSIDAE Fry, 1978** Recent
- no fossil record
- ENDEIDAE Norman, 1904** ?Jurassic – Recent
- † **Palaeoendeis Charbonnier, Vannier & Riou, 2007** Jurassic
11. *Palaeoendeis elmii* Charbonnier, Vannier & Riou, 2007* J La Voulte-sur-Rhône
tentative referal
- PHOXICHILIIDIIDAE Sars, 1891** Recent
- = ANOPLODACTYLIDAE Fry, 1978
 - = PHOXIPHILYRIDAE Fry, 1978

no fossil record

RHYNCHOTHORACIDAE Thompson, 1909 Recent

no fossil record

MISIDENTIFICATIONS

1. *Pentapalaæopycnon inconspicua* Hedgpeth, 1978 [crustacean] J Solnhofen
2. *Pycnogonites uncinatus* Quenstedt, 1852 [crustacean] J Solnhofen

c. 1,300 Recent species

(EU)CHELICERATA

5 currently valid, but unplaced (eu)chelicerate fossil species

- *Sanctacaris* has been recovered as an early chelicerate in some phylogenetic studies – most recently by Legg (2014) – although this interpretation is not universal
- *Offacolus* has been described in detail from reconstructions based on serial sections, and was resolved in some phylogenies to a basal position within Euchelicera
- *Dibasterium* was described as a horseshoe crab, albeit one with multiple biramous appendages
- the other listed taxa are mostly poor or incomplete specimens which have been treated as either xiphosurans, chasmataspids or eurypterids
- resting impressions imply that Chasmataspida were probably present in the late Cambrian

CHELICERATA Heymons, 1901 ?Cambrian – Recent

† *Sanctacaris* Briggs & Collins, 1988 Cambrian
 1. *Sanctacaris uncata* Briggs & Collins, 1988* C Burgess Shale

EUCHELICERATA Weygoldt & Paulus, 1979 ?Cambrian – Recent

STEM-EUCHELICERATA?

† *Offacolus* Orr, Siveter, Briggs, Siveter & Sutton, 2000 Silurian
 2. *Offacolus kingi* Orr, Siveter, Briggs, Siveter & Sutton, 2000* S Herefordshire Lgst.
 † *Dibasterium* Briggs, Siveter, Siveter, Sutton, Garwood & Legg, 2012 Silurian
 3. *Dibasterium durgae* Briggs, Siveter, Siveter, Sutton, Garwood & Legg,
 2012* S Herefordshire Lgst.

EUCHELICERATA INCERTAE SEDIS

† *Polystomurum* Novojilov, 1958 Devonian
 4. *Polystomurum stormeri* Novojilov, 1958* D Voroneje, Siberia
 † *Thurandina* Størmer, 1974 Devonian
 5. *Thurandina waterstoni* Størmer, 1974* D Alken an der Mosel

XIPHOSURA s. lat.

104 currently valid species traditionally assigned to horseshoe crabs, of which 82 are unequivocal Xiphosura

- Lamsdell (2013) argued that Xiphosura may not be monophyletic and that a number of fossils traditionally placed as stem-group (synziphosurine) horseshoe crabs are actually stem-group euhelicerates. The list below attempts to reflect this position, whereby it should be noted that in this scheme the Planaterga clade would also include Chasmataspidida, Eurypterida and Arachnida and Planaterga is nested within Prosomapoda.

PROSOMAPODA Lamsdell, 2013a Siliurian – Recent

FAMILY UNSPECIFIED

† Anderella Moore, McKenzie & Lieberman, 2007	Carboniferous
1. <i>Anderella parva</i> Moore, McKenzie & Lieberman, 2007*	C Bear Gulch
† Borchgrevinkium Novojilov, 1959	Devonian
2. <i>Borchgrevinkium taimyrensis</i> Novojilov, 1959*	D Taimyr, Siberia
† Camanchia Moore, Briggs, Braddy & Shultz, 2011	Silurian
3. <i>Camanchia grovensis</i> Moore, Briggs, Braddy & Shultz, 2011*	S Scotch Grove, Iowa
† Legrandella Eldredge, 1974	Devonian
4. <i>Legrandella lombardii</i> Eldredge, 1974*	D Cochabamba, Bolivia
† Venustulus Moore, 2005 <i>in Moore et al.</i>	Silurian
5. <i>Venustulus waukeshaensis</i> Moore, 2005 <i>in Moore et al.*</i>	S Waukesha Lgst.
† WEINBERGINIDAE Richter & Richter, 1929	Devonian
† Weinbergina Richter & Richter, 1929	Devonian
6. <i>Weinbergina opitzi</i> Richter & Richter, 1929*	D Hunsruckschiefer

PLANATERGA Lamsdell, 2013a Siliurian – Recent

FAMILY UNSPECIFIED

† Bembicosoma Laurie, 1899	Silurian
7. <i>Bembicosoma pomphicus</i> Laurie, 1899*	S Pentland hills
† Cyamocephalus Currie, 1927	Silurian
8. <i>Cyamocephalus loganensis</i> Currie, 1927*	S Lesmahagow
† Pseudoniscus Nieszkowski, 1859	Silurian
= † <i>Neolimulus</i> Woodward, 1868a		
9. <i>Pseudoniscus aculeatus</i> Nieszkowski, 1859*	S Saaremaa
10. <i>Pseudoniscus clarkei</i> Ruedemann, 1916	S Pittsford, New York
11. <i>Pseudoniscus falcatus</i> (Woodward, 1868a)	S Lesmahagow
12. <i>Pseudoniscus roosevelti</i> Clarke, 1902	S ‘Bertie Waterlime’
† Bunaia Clarke, 1919	Silurian

13. ' <i>Bunaia</i> ' <i>heintzi</i> Størmer, 1934a	S Spitsbergen
14. <i>Bunaia woodwardi</i> Clarke, 1919*	S 'Bertie Waterlime'
† BUNODIDAE Packard, 1896	Silurian
† <i>Bunodes</i> Eichwald, 1854	Silurian
= † <i>Exapinurus</i> Nieszkowski, 1859	
15. <i>Bunodes lunula</i> Eichwald, 1854*	S Saaremaa
i. = <i>Bunodes rugosus</i> Eichwald, 1854	S Saaremaa
ii. = <i>Exapinurus schrenki</i> Nieszkowski, 1859	S Saaremaa
† <i>Limuloides</i> Woodward, 1865	Silurian
= † <i>Hemiaspis</i> Woodward, 1864 [preoccupied]	
16. <i>Limuloides limuloides</i> (Woodward, 1865)	S Ludlow
17. <i>Limuloides horridus</i> (Woodward, 1872a)	S Ludlow
18. <i>Limuloides salweyi</i> (Woodward, 1872a)	S Ludlow
i. = <i>Hemiaspis tuberculatus</i> (Salter in Woodward, 1872a) S Ludlow	
19. <i>Limuloides speratus</i> Woodward, 1872a	S Ludlow
i. = <i>Hemiaspis optatus</i> (Salter in Woodward, 1872a) S Ludlow	
† <i>Pasternakevia</i> Selden & Drygant, 1987	Silurian
20. <i>Pasternakevia podolica</i> Selden & Drygant, 1987*	S Podolia

Planaterga sensu Lamsdell (2013a) also includes chasmataspids, eurypterids and arachnids

XIPHOSURA Latreille, 1802	Ordovician – Recent
= MEROSTOMATA Dana, 1852	

FAMILY UNSPECIFIED

† <i>Kiaeria</i> Størmer, 1934b	Silurian
21. <i>Kiaeria limuloides</i> Størmer, 1934b*	S Ringerike
† <i>Maldybulakia</i> Tesakov & Alekseev, 1998	Devonian
= † <i>Lophodesmus</i> Tesakov & Alekseev, 1992 [preoccupied]	
NB: Originally described as possible myriapods	
22. <i>Maldybulakia angusi</i> Edgecombe, 1998	D New South Wales
23. <i>Maldybulakia malcomi</i> Edgecombe, 1998	D New South Wales
24. <i>Maldybulakia mirabilis</i> (Tesakov & Alekseev, 1992)*	D Kazakhstan
† <i>Willwerathia</i> Størmer, 1969	Devonian
25. <i>Willwerathia laticeps</i> (Størmer, 1936a)*	D Willwerath

† KASIBELINURIDAE Pickett, 1993	Devonian
† <i>Kasibelinurus</i> Pickett, 1993	Devonian
26. <i>Kasibelinurus amicorum</i> Pickett, 1993*	D New South Wales
27. <i>Kasibelinurus yueya</i> Lamsdell, Xue & Selden, 2013	D Yunann, China

possible kasibelinurids?

28. '*Belinurus*' *alleghenyensis* Eller, 1938a D New York State
 29. '*Belinurus*' *carterae* Eller, 1940 D Pennsylvania
 30. '*Prestwichia*' *randalli* Beecher, 1902 D Pennsylvania

- † **ELLERIDAE** Raymond, 1944 Devonian
 † ***Elleria*** Raymond, 1944 Devonian
 31. *Elleria morani* (Eller, 1938b)* D Pennsylvania

XIPHOSURIDA Latreille, 1802 Ordovician – Recent

family uncertain

- † ***Lunataspis*** Rudkin, Young & Nowlan, 2008 Ordovician
 32. *Lunataspis aurora* Rudkin, Young & Nowlan, 2008 O Manitoba

- † **BELINURINA** Zittel & Eastman, 1913 Carboniferous

- † **BELINURIDAE** Zittel & Eastman, 1913 Carboniferous

- † ***Bellinurus*** Pictet, 1846 Carboniferous

= † *Belinurus* König, 1851

= † *Steropsis* Baily, 1869

= † *Koenigiella* Raymond, 1944

NB: Pictet's 1846 name *Bellinurus* [sic] was based on a misspelling of *Belinurus* from König's unpublished plates, which themselves only became available posthumously as of 1851

33. *Bellinurus arcuatus* Baily, 1863 C Coal Measures
 34. *Bellinurus baldwini* Woodward, 1907b C Coal Measures
 35. *Bellinurus bellulus* Pictet, 1846 C Coalbrookdale, UK
 36. *Bellinurus carwayensis* Dix & Pringle, 1929 C South Wales, UK
 37. *Bellinurus concinnus* Dix & Pringle, 1929 C South Wales, UK
 38. *Bellinurus grandaevis* Jones & Woodward, 1899 C Nova Scotia
 39. *Bellinurus iswariensis* (Chernyshev, 1928) C Donetsk Basin
 40. *Bellinurus kiltorkensis* Baily, 1869 C Coal Measures
 41. *Bellinurus koenigianus* Woodward, 1872a C Coal Measures
 42. *Bellinurus lacoei* Packard, 1885 C Mazon Creek
 43. *Bellinurus longicaudatus* Woodward, 1907b C Coal Measures
 44. *Bellinurus lunatus* (Martin, 1809) C Mansfield, UK
 45. *Bellinurus metschetensis* (Chernyshev, 1928) C Donetsk Basin
 46. *Bellinurus morgani* Dix & Pringle, 1930 C South Wales, UK
 47. *Bellinurus pustulosus* Dix & Pringle, 1929 C South Wales, UK
 48. *Bellinurus reginae* Baily, 1863 C Coal Measures
 49. *Bellinurus stepanovi* (Chernyshev, 1928) C Donetsk Basin
 50. *Bellinurus trechmanni* Woodward, 1918 C Coal Measures
 51. *Bellinurus trilobitoides* (Buckland, 1837)* C Coalbrookdale, UK
 52. *Bellinurus truemani* Dix & Pringle, 1929 C South Wales, UK

† EUPROOPIDAE Eller, 1938b

= † LIOMESASPIDIDAE Raymond, 1944

† <i>Anacontium</i> Raymond, 1944	Permian
53. <i>Anacontium brevis</i> Raymond, 1944	P Oklahoma
54. <i>Anacontium carpenteri</i> Raymond, 1944	P Oklahoma
† <i>Euproops</i> Meek, 1867	Carbon. – ?Permian
= † <i>Prestwichia</i> Woodward, 1867 [preoccupied]	
= † <i>Prestwichianella</i> Cockerell, 1905 [replacement name for <i>Prestwichia</i>]	
55. <i>Euproops anthrax</i> (Prestwich, 1840)	C Coal Measures
56. <i>Euproops bifidus</i> Siegfried, 1972	C Coal Measures
57. <i>Euproops cambrensis</i> Dix & Pringle, 1929	C Coal Measures
58. <i>Euproops danae</i> (Meek & Worthen, 1865)*	C Coal Measures
i. = <i>Euproops amiae</i> Woodward, 1918	C Coal Measures
ii. = <i>Euproops darrahi</i> Raymond, 1944	C Coal Measures
iii. = <i>Euproops graigolae</i> Dix & Pringle, 1929	C South Wales
iv. = <i>Euproops gwenti</i> Dix & Pringle, 1929	C South Wales
v. = <i>Euproops islwyni</i> Dix & Pringle, 1929	C South Wales
vi. = <i>Euproops kilmersdonensis</i> Ambrose & Romano, 1972	C Kilmersdon, UK
vii. = <i>Euproops laevicula</i> Raymond, 1944	C Coal Measures
viii. = <i>Euproops laticephalus</i> Raymond, 1944	C Coal Measures
ix. = <i>Euproops packardi</i> Willard & Jones, 1935	C Coal Measures
x. = <i>Prestwichia (Euproops) scheeleana</i> Ebert, 1892	C Coal Measures
xi. = <i>Euproops thompsoni</i> Raymond, 1944	C Coal Measures
59. <i>Euproops longispina</i> Packard, 1885	C Mazon Creek
60. <i>Euproops mariae</i> Crônier & Courville, 2005	C Massif Central
61. <i>Euproops meeki</i> Dix & Pringle, 1929	C South Wales
62. <i>Euproops nitida</i> Dix & Pringle, 1929	C South Wales
63. <i>Euproops orientalis</i> Kobayashi, 1933	?P Korea
64. <i>Euproops rotundatus</i> Prestwich, 1840	C Coal Measures
<i>Euproops</i> sp. in Brauckmann (1982)	C Piesberg, Germany
† <i>Liomesaspis</i> Raymond, 1944	Carbon. – Permian
= † <i>Pringlia</i> Raymond, 1944	
= † <i>Palatinaspis</i> Malz & Poschmann, 1993	
65. ? <i>Liomesaspis birtwelli</i> (Woodward, 1872a)	C Coal Measures
66. <i>Liomesaspis laevis</i> Raymond, 1944*	C Coal Measures
i. = <i>Palatinaspis beimbaueri</i> Malz & Poschmann, 1993	C Saar-Nahe Basin
ii. = <i>Pringlia bispinosa</i> Raymond, 1944	C Coal Measures
iii. = <i>Pringlia demaisterei</i> Vandenbergh, 1961	C Coal Measures
iv. = <i>Pringlia fritschii</i> Remy & Remy, 1959	C Coal Measures
67. <i>Liomesaspis leonardensis</i> (Tasch, 1961)	P Annelly, Kansas
† <i>Prolimulus</i> Frič, 1899	Carboniferous
68. <i>Prolimulus woodwardi</i> Frič, 1899*	C Nýřany

UNNAMED TAXON

- † *Bellinuroopsis* Chernyshev, 1933 Carboniferous
 = † *Neobelinuroopsis* Eller, 1938a
69. *Bellinuroopsis rossicus* Chernyshev, 1933* C Coal Measures
- † ROLFEIIDAE Selden & Siveter, 1987 Carboniferous
- † *Rolfeia* Waterston, 1985 Carboniferous
 70. *Rolfeia fouldenensis* Waterston, 1985* C Fouldon, Scotland
- LIMULINA Richter & Richter, 1929 Carbon. – Recent
 Unanmed specimen in Krause et al. (2009) Tr Ohrdruf, Germany
- † PALEOLIMULOIDEA Raymond, 1944 Carbon. – Jurassic
- † PALEOLIMULIDAE Raymond, 1944 Carbon. – Jurassic
 = † MESOLIMULIDAE (Størmer, 1952) [in part; see Reik & Gill 1971]
 = † DUBBOLIMULIDAE Pickett, 1984
- † *Limulitella* Størmer, 1952 Triassic – Jurassic
 = † *Limulites* Schimper, 1853 [preoccupied]
 Limulitella sp. in Hauschke et al. (2004) Tr Madagascar
 ?*Limulitella* sp. in Hauschke & Wilde (2008) Tr Dallau, Germany
 ?*Limulitella* sp. in Hauschke et al. (2009) Tr Winterswijk
 71. *Limulitella bronni* (Schimper, 1853)* Tr Grés à Voltzia
 i. = *Limulus sandbergeri* Kirchner, 1923 Tr Germany
 72. *Limulitella henkeli* Fritsch, 1906 Tr Halle, Germany
 73. ?*Limulitella liasokeuperensis* (Braun, 1860) J Germany
 74. *Limulitella vicensis* (Bleicher, 1897) Tr Lorraine
 75. *Limulitella volgensis* Ponomarenko, 1985 Tr Moscow
- † *Paleolimulus* Dunbar, 1923 Carbon. – Triassic
 = † *Dubbolimulus* Pickett, 1984
 ?*Palaeolimulus* sp. in Hauschke & Wilde (2000) Tr Harz, Germany
 76. *Paleolimulus fuchsbergensis* Hauschke & Wilde, 1987 Tr northwest Germany
 77. *Paleolimulus jakovlevi* Glushenko in Glushenko & Ivanov, 1961 P Novoselovka, Ukraine
 78. ?*Paleolimulus juresanensis* Chernyshev, 1933 C Ural region
 79. *Paleolimulus longispinus* Schram, 1979 C Bear Gulch, Montana
 80. *Paleolimulus peetae* (Pickett, 1984) Tr New South Wales
 81. *Paleolimulus signatus* (Beecher, 1904) C-P Kansas, Illinois
 i. = *Paleolimulus avitus* Dunbar, 1923* P Kansas
- MORAVURIDAE Příbyl, 1967 Carboniferous
- † *Moravurus* Příbyl, 1967 Carboniferous
 82. *Moravurus rehori* Příbyl, 1967 C Ostrava-Karviná

- † *Xaniopyramis* Siveter & Selden, 1987 Carboniferous
 83. *Xaniopyramis linseyi* Siveter & Selden, 1987* C Weardale, UK
- LIMULOIDEA Zittel, 1885** Carbon. – Recent
 unnamed specimen *in* Hauschke & Wilde (1989) P Korbacher Bucht
- † *Alanops* Racheboeuf et al., 2002 Carboniferous
 84. *Alanops magnifica* Racheboeuf et al., 2002 C Montceau-les-Mines
- † *Casterolimulus* Holland, Erickson & O'Brien, 1975 Cretaceous
 85. *Casterolimulus kletti* Holland, Erickson & O'Brien, 1975* K North Dakota
- † *Panduralimulus* Allen & Feldman, 2005 Permian
 86. *Panduralimulus babcocki* Allen & Feldman, 2005 P Texas
- † *Valloisella* Racheboeuf, 1992 Carboniferous
 87. *Valloisella lievinensis* Racheboeuf, 1992* C northern France
- † **AUSTROLIMULIDAE Riek, 1955** Triassic
 † *Austrolimulus* Riek, 1955 Triassic
 88. *Austrolimulus fletcheri* Riek, 1955* Tr New South Wales
- LIMULIDAE Zittel, 1885** Triassic – Recent
 = † **MESOLIMULIDAE** (Størmer, 1952) [in part; see Reik & Gill 1971]
 ?Limulidae gen. et sp. indet *in* Hauschke et al. (1992) Tr Rüdersdorf, Germany
 † *Crenatolimulus* Feldmann, Schweitzer, Dattilo & Farlow, 2011 Cretaceous
 89. *Crenatolimulus paluxyenis* Feldmann, Schweitzer, Dattilo & Farlow,
 2011* K Texas
- Limulus* Müller, 1785 Triassic – Recent
 90. *Limulus coffini* Reeside & Harris, 1952 K Colorado
 91. *Limulus darwini* Kin & Błażejowski, 2014 J Kcynia, Poland
 92. "Limulus" decheni Zinken, 1862 Pa Teuchern, Germany
 [NB: Hauschke & Wilde (2004) considered this intermediate between *Limulus* and *Tachypleus*]
 93. *Limulus priscus* Münster, 1839 Tr Rottweil, Germany
 94. *Limulus woodwardi* Watson, 1909 J Northamptonshire
- † **Mesolimulus** Størmer, 1952 Triassic – Cretaceous
 Mesolimulus sp. *in* Ross & Vannier (2002) J southern England
 95. *Mesolimulus cespelli* Via Boada, 1987 Tr Tarragona, Spain
 96. *Mesolimulus sibiricus* Ponomarenko, 1985 J Siberia
 97. ?*Mesolimulus syriacus* (Woodward, 1879) K Lebanon
 98. *Mesolimulus walchi* (Desmarest, 1822)* J Solnhofen, etc.
 i. = *Limulus brevicauda* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 ii. = *Limulus brevispina* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 iii. = *Limulus intermedius* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 iv. = *Limulus ornatus* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 v. = *Limulus sulcatus* Münster *in* v. d. Hoeven, 1838 J Solnhofen

- vi. = *Limulus giganteus* Münster, 1840 J Solnhofen
 NB: not entirely clearly that all these names have been formally synonymised
- † ***Psammolimulus* Lange, 1923** Triassic
99. *Psammolimulus gottingensis* Lange, 1923* Tr Göttingen, Germany
- Tachypleus* Leach, 1819** Triassic – Recent
- = † *Heterolimulus* Via Boada & Villalta, 1966
100. *Tachypleus gadeai* (Via Boada & Villalta, 1966) Tr Tarragona, Spain
- † ***Tarracolimulus* Romero & Via Boada, 1977** Triassic
101. *Tarracolimulus rieki* Romero & Via Boada, 1977* Tr Tarragona, Spain
- † ***Victalimulus* Riek & Gill, 1971** Cretaceous
102. *Victalimulus mcqueeni* Riek & Gill, 1971* K Koonwarra
- † ***Yunnanolimulus* Zhang, Hu, Zhou, Lv & Bai, 2009** Triassic
103. *Yunnanolimulus luopingensis* Zhang, Hu, Zhou, Lv & Bai, 2009* Tr Luoping, China

INCERTAE SEDIS

† ***Belinuopsis* Matthew 1910**

104. *Belinuopsis wigudensis* Matthew, 1910 C Coal Measures

NOMEN DUBIUM

1. *Limulus nathorsti* Jackson, 1906 J southern Sweden

NOMINA NUDA

1. *Euproops rotunda major* (Woodward, 1907) C Sparth Bottoms
2. *Veltheimia bicornis* Beyschlag & von Fritsch, 1899 C? Rotliegend

MISIDENTIFICATIONS

1. *Belinurus carterae* Eller, 1940 [synonym of *P. eriensis*; see below]
2. *Bifarius comptae* Tasch, 1961 [insect] P Kansas
3. *Eolimulus alatus* Moberg, 1892 [doubtful xiphosuran] C Öland, Sweden
4. *Elmocephalus carltonensis* (Tasch, 1963) [?crustacean] P Kansas
5. *Hemiaspis tunnecliffei* Chapman, 1932 [trilobite] S Victoria
6. *Hypatocephala rugosa* Tasch, 1961 [insect] P Kansas
7. *Lemoneites ambiguus* Flower, 1969 [Echinodermata] O Texas
8. *Lemoneites gomphocaudatus* Flower, 1969 [Echinodermata] O Texas
9. *Lemoneites mirabilis* Flower, 1969 [Echinodermata] O Texas
10. *Lemoneites simplex* Flower, 1969 [Echinodermata] O Texas
11. *Pincombella belmontensis* Chapman, 1932 [insect – Hemiptera] P New South Wales
12. *Permolimulinella rarissima* Tasch, 1963 [insect] P Kansas
13. *Strongylocephalus charactis* Tasch, 1961 [insect] P Kansas
14. *Protolimulus eriensis* [Xiphosuran trace fossil: see *Selenichnites*]

CHASMATASPIDIDA

11 currently valid species of fossil chasmataspidid

- there are some doubts about the monophyly of Chasmataspidida

† CHASMATASPIDIDA Caster & Brooks, 1956 ?Camb. – Devonian

= † DIPLOASPIDIDA Simonetta & Delle Cave, 1978

† CHASMATASPIDIDAE Caster & Brooks, 1956 ?Camb. – Ordovician

† Chasmataspis Caster & Brooks, 1956 ?Camb. – Ordovician

?*Chasmataspis* sp. resting traces in Dunlop et al. (2004) E Texas

1. *Chasmataspis laurencii* Caster & Brooks, 1956* O Tennessee

† DIPLOASPIDIDAE Størmer, 1972 Silurian – Devonian

= † HETEROASPIDIDAE Størmer, 1972

† Achanarraspis Anderson, Dunlop & Trewin, 2000 Devonian

2. *Achanarraspis reedi* Anderson, Dunlop & Trewin, 2000* D Achanarras, Scotland

† Diploaspis Størmer, 1972 Devonian

3. *Diploaspis casteri* Størmer, 1972* D Alken an der Mosel

4. *Diploaspis muelleri* Poschmann, Anderson & Dunlop, 2005 D Hombach, Germany

† Dvulikiaspis Marshall, Lamsdell, Shpinev & Braddy, 2014 Devonian

5. *Dvulikiaspis menneri* (Novojilov, 1959)* D Siberia

† Forfarella Dunlop, Anderson & Braddy, 1999 Devonian

6. *Forfarella mitchelli* Dunlop, Anderson & Braddy, 1999* D Arbroath, Scotland

† Heteroaspis Størmer, 1972

7. *Heteroaspis stoermeri* (Novojilov, 1959)* D Siberia; Alken

i. = *Heteroaspis novojilovi* Størmer, 1972 D Alken an der Mosel

† Loganamaraspis Tetlie & Braddy, 2004a Silurian

8. *Loganamaraspis dunlopi* Tetlie & Braddy, 2004a* S Lesmahagow

† Nahlyostaspis Marshall, Lamsdell, Shpinev & Braddy, 2014 Devonian

9. *Nahlyostaspis bergstroemi* Marshall, Lamsdell, Shpinev & Braddy,
2014* D Siberia

† Octoberaspis Dunlop, 2002 Devonian

10. *Octoberaspis ushakovi* Dunlop, 2002* D October Rev. Is

† Skrytyaspis Marshall, Lamsdell, Shpinev & Braddy, 2014 Devonian

11. *Skrytyaspis andersoni* Marshall, Lamsdell, Shpinev & Braddy, 2014* D Siberia

no Recent species

EURYPTERIDA

250 currently valid species of fossil sea scorpion

- Tollerton (1989) suggested removing Hibbertopteroidea from Eurypterida s.s., but this has not been adopted by subsequent workers and they are treated here as derived stylonurid eurypterids

† EURYPTERIDA Burmeister, 1843	Ordovician – Permian
	= † GIGANTOSTRACA Haeckel, 1866	
	= † CYRTOCTENIDA Størmer & Waterston, 1968	
† STYLONURINA Diener, 1924	Ordovician – Permian
	= † WOODWARDOPTERINA Kjellesvig-Waering, 1959	
	= † HIBBERTOPTERINA Størmer, 1974	
† RHENOPTEROIDEA Størmer, 1951	Ordovician – Devonian
	= † BRACHYOPTERELLOIDEA Tollerton, 1989	
† RHENOPTERIDAE Størmer, 1951	Ordovician – Devonian
	= † BRACHYOPTERELLIDAE Tollerton, 1989	
	= † ALKENOPTERIDAE Poschmann & Tetlie, 2004	
† Brachyopterella Kjellesvig-Waering, 1966a	Silurian
1. <i>Brachyopterella pentagonalis</i> (Størmer, 1934b)*	S Ringerike, Norway
2. <i>Brachyopterella ritchiei</i> Waterston, 1979	S Slot Burn, Scotland
† Brachypterus Størmer, 1951	Ordovician
3. <i>Brachypterus stubblefieldi</i> Størmer, 1951*	O Montgomeryshire
† Kiaeropterus Waterston, 1979	Silurian
4. <i>Kiaeropterus cyclophthalmus</i> (Laurie, 1892)	S Pentland Hills, Scotl.
5. <i>Kiaeropterus ruedemanni</i> (Størmer, 1934b)*	S Ringerike, Norway
† Leiopterala Lamsdell, Braddy, Loeffler & Dineley, 2010	Devonian
6. <i>Leiopterala tetliei</i> Lamsdell, Braddy, Loeffler & Dineley, 2010	D Nunavut, Canada
† Rhenopterus Størmer, 1936a	Devonian
7. <i>Rhenopterus diensti</i> Størmer, 1936a*	D Willwerath, Germ.
i. = <i>Rhenopterus latus</i> Størmer, 1936a	D Willwerath, Germ.
8. <i>Rhenopterus macrotuberculatus</i> Størmer, 1974	D Alken an der Mosel
9. <i>Rhenopterus tuberculatus</i> Størmer, 1936a	D Overath, Germ.
† STYLONUROIDEA Kjellesvig-Waering, 1959	Silurian – Devonian
† PARASTYLONURIDAE Waterston, 1979	Silurian – Devonian
† Parastylonurus Kjellesvig-Waering, 1966a	Silurian
10. <i>Parastylonurus hendersoni</i> Waterston, 1979	S Pentland Hills, Scotl.
11. <i>Parastylonurus ornatus</i> (Laurie, 1892)*	S Scotland

12. ?*Parastylonurus sigmoidalis* Kjellesvig-Waering, 1971 S Shropshire, UK
- † *Stylorella* Kjellesvig-Waering, 1966a Silurian – Devonian
13. *Stylorella ?arnoldi* (Ehlers, 1935) D Pennsylvania, USA
14. *Stylorella ?beecheri* (Hall, 1884c) D Pennsylvania, USA
15. *Stylorella spinipes* (Page, 1859)* S Kip Burn, Scotland
 i. = *Stylourus logani* Woodward, 1872 S Kip Burn, Scotland
- † STYLOURIDAE Diener, 1924 Silurian–Devonian
- = † LAURIEIPTERIDAE Kjellesvig-Waering, 1966a
- = † PAGEIDAE Kjellesvig-Waering, 1966a
- † *Ctenopterus* Clarke & Ruedemann, 1912 Silurian
16. *Ctenopterus cestrotus* (Clarke, 1907)* S Otisville, New York
- † *Laurieipterus* Kjellesvig-Waering, 1966a Silurian
17. *Laurieipterus elegans* (Laurie, 1899)* S Pentland Hills, Scotl.
- † *Pagea* Waterston, 1962 Devonian
18. *Pagea plotnicki* Lamsdell, Braddy, Loeffler & Dineley, 2010 D Nunavut, Canada
19. *Pagea sturrocki* Waterston, 1962* D Old Red Sandstone
20. *Pagea symondsii* (Salter, 1859) D Old Red Sandstone
- † *Stylourus* Page, 1856 Devonian
21. *Stylourus powriensis* Page, 1856* D Mid. Valley Scotland
 i. = *Stylourus ensiformis* Woodward, 1864 D Mid. Valley Scotland
22. ?*Stylourus shaffneri* Willard, 1933 D Pennsylvania
- † KOKOMOPTEROIDEA Kjellesvig-Waering, 1966a Silurian
- † KOKOMOPTERIDAE Kjellesvig-Waering, 1966a Silurian
- † *Kokomopterus* Kjellesvig-Waering, 1966a Silurian
23. *Kokomopterus longicaudatus* (Clarke & Ruedemann, 1912)* S Kokomo, Indiana
- † *Lamontopterus* Waterston, 1979 Silurian
24. *Lamontopterus knoxae* (Lamont, 1955)* S Pentland Hills, Scotl.
- † HARDIEOPTERIDAE Tollerton, 1989 Silurian – Devonian
- † *Hallipterus* Kjellesvig-Waering, 1963a Devonian
25. *Hallipterus excelsior* (Hall, 1884a)* D New York
 i. = *Dolichocephala lacoana* Claypole, 1883 D Pennsylvania
- † *Hardieopterus* Waterston, 1979 Silurian
26. ?*Hardieopterus lanarkensis* Waterston, 1979 S Patrick Burn, Scotl.
27. *Hardieopterus macrorthalmus* (Laurie, 1892)* S Pentland Hills, Scotl.
28. *Hardieopterus megalops* (Salter, 1859) S Herefordshire, Engl.
29. *Hardieopterus myops* (Clarke, 1907) S eastern USA
- † *Tarsopterella* Størmer, 1951 Devonian
30. *Tarsopterella scotica* (Woodward, 1872)* D Mid. Valley Scotland
 i. = ?*Erieopterus brewsteri* Woodward, 1864 D Mid. Valley Scotland

- ii. = *Stylonurus armatus* Page, 1867 D Mid. Valley Scotland
- † MYCTEROPOIDEA Cope, 1886** Silurian – Permian
- = † HIBBERTOPTEROIDEA Kjellesvig-Waering, 1959
- † DREPANOPTERIDAE Kjellesvig-Waering, 1966a** Silurian – Devonian
- † Drepanopterus Laurie, 1892** Silurian – Devonian
31. *Drepanopterus abonensis* Simpson, 1951 D Portishead, England
32. *Drepanopterus odontospathus* Lamsdell, 2012 D Arctic Canada
33. *Drepanopterus pentlandicus* Laurie, 1892* S Pentland Hills, Scotl.
- † HIBBERTOPTERIDAE Kjellesvig-Waering, 1959** Devonain – Permian
- = † CYRTOCTENIDAE Waterston, Oelofsen & Oosthuizen, 1985
- † Campylocephalus Eichwald, 1860** Carboniferous – Perm.
34. *Campylocephalus oculatus* (Kutorga, 1838)* P Dourasovo, Russia
35. *Campylocephalus permianus* (Ponomarenko, 1985) P Komi, Russia
36. ?*Campylocephalus salmi* Stur, 1877 C Ostrava, Czech Rep.
- † Cyrtocetus Størmer & Waterston, 1968** Devonian – Carbon.
37. *Cyrtocetus caledonicus* (Salter, 1863) C East Lothian, Scotl.
38. *Cyrtocetus dewalquei* (Fraipont, 1889) D Pont-de-Bonne, Belg.
- i. = *Eurypterus dewalquei* var. *longimanus* Fraipont,
1889 D Pont-de-Bonne, Belg.
39. *Cyrtocetus dicki* (Peach, 1883) C Thurso, Scotland
40. *Cyrtocetus ostraviensis* (Augusta & Přibyl, 1951) C Ostrava, Czech Rep.
41. *Cyrtocetus peachi* Størmer & Waterston, 1968* C Berwickshire, Scotl.
42. *Cyrtocetus wittebergensis* Waterston, Oelofsen & Oosthuizen, 1985 ... C Cape Province
- † Dunsopterus Waterston, 1968** Carboniferous
43. *Dunsopterus stevensoni* (Etheridge Jr, 1877)* C Berwickshire, Scotl.
- † Hastimima White, 1908** Permian
44. *Hastimima whitei* White, 1908* P Brazil
- † Hibbertopterus Kjellesvig-Waering, 1959** Carboniferous – Perm.
45. ?*Hibbertopterus hibernicus* (Baily, 1872) C Kiltoran, Ireland
46. *Hibbertopterus scouleri* (Hibbert, 1836)* C West Lothian, Scotl.
- † Vernonopterus Waterston, 1957** Carboniferous
47. *Vernonopterus minutisculptus* (Peach, 1907)* C Lanarkshire, Scotland
- † MYCTEROPIDAE Cope, 1886** Carboniferous – Perm.
- = † WOODWARDOPTERIDAE Kjellesvig-Waering, 1959
- † Megarachne Hünicken, 1980** Carboniferous – Perm.
48. *Megarachne servinei* Hünicken, 1980* C-P Santa Rosa, Argen.
- † Mycterops Cope, 1886** Carboniferous
49. ?*Mycterops blairi* Waterston, 1968 C Loanhead, Scotland
50. *Mycterops matthieu* Pruvost, 1924 C Charleroi, Belgium

51. *Mycterops ordinatus* Cope, 1886* C Channelton, PA
52. ?*Mycterops whitei* Schram, 1984 C Crescent, Iowa
- † **Woodwardopterus** Kjellesvig-Waering, 1959 Carboniferous
53. *Woodwardopterus scabrosus* (Woodward, 1887)* C Glencarholm, Scotl.
- STYLONURINA incertae sedis**
- † **Stylonuroides** Kjellesvig-Waering, 1966a Silurian – Devonian
54. *Stylonuroides dolichopteroides* (Størmer, 1934b)* S Ringerike, Norway
55. *Stylonuroides orientalis* Shpinev, 2012 D Lake Shunet, Siberia
- † **EURYPTERINA** Burmeister, 1843 Ordovician – Permian
- † **ONYCHOPTERELLOIDEA** Lamsdell, 2011 Ordovician–Silurian
- † **ONYCHOPTERELLIDAE** Lamsdell, 2011 Ordovician–Silurian
- † **Alkenopterus** Størmer, 1974 Devonian
56. *Alkenopterus brevitelson* Størmer, 1974* D Alken an der Mosel
57. *Alkenopterus burglahrensis* Poschmann & Tetlie, 2004 D Westerwald, Germ.
- † **Onychopterella** Størmer, 1951 Ordovician–Silurian
58. *Onychopterella augusti* Braddy, Aldridge & Theron, 1995 O Soom Shale, S. Afr.
59. *Onychopterella kokomoensis* (Miller & Gurley, 1896)* S Kokomo, Indiana
- i. = *Eurypterus ranilarva* Clarke & Ruedemann, 1912 S Kokomo, Indiana
60. ?*Onychopterella pumilus* (Savage, 1916) S Essex, Illinois
- † **Tyloptera** Størmer, 1951 Silurian
61. *Tyloptera boylei* (Whiteaves, 1884) S Ontario, Canada
- † **MOSELOPTEROIDEA** Lamsdell, Braddy & Tetlie, 2010 Silurian – Devonian
- † **MOSELOPTERIDAE** Lamsdell, Braddy & Tetlie, 2010 Devonian
- † **Moselopterus** Størmer, 1974 Devonian
62. *Moselopterus aencylotelson* Størmer, 1974* D Alken an der Mosel
63. *Moselopterus elongatus* Størmer, 1974 D Alken an der Mosel
64. *Moselopterus lancmani* (Delle, 1937) D Plavinas, Latvia
- † **Stoermeropterus** Lamsdell, 2011 Silurian
65. *Stoermeropterus conicus* (Laurie, 1892)* S Pentland Hills
- i. = *Drepanopterus bembycoides* Laurie, 1899 S Pentland Hills
- ii. = *Drepanopterus lobatus* Laurie, 1899 S Pentland Hills
66. *Stoermeropterus latus* (Størmer, 1934b) S Ringerike, Norway
67. *Stoermeropterus nodosus* (Kjellesvig-Waering & Leutze, 1966) S Bass, West Virginia
- † **Vinetopterus** Poschmann & Tetlie, 2004 Devonian
68. *Vinetopterus martini* Poschmann & Tetlie, 2004 D Westerwald, Germ.
69. *Vinetopterus struvei* (Størmer, 1974)* D Alken an der Mosel
- † **MEGALOGRAPTOIDEA** Caster & Kjellesvig-Waering, 1955 Ordovician
- † **MEGALOGRAPTIDAE** Caster & Kjellesvig-Waering, 1955 Ordovician

† <i>Echinognathus</i> Walcott, 1882	Ordovician
70. <i>Echinognathus clevelandi</i> Walcott, 1882*	O New York
† <i>Megalograptus</i> Miller, 1874	Ordovician
71. <i>Megalograptus alveolatus</i> (Shuler, 1915)	O Virginia
72. <i>Megalograptus ohioensis</i> Caster & Kjellesvig-Waering, 1955	O Ohio
73. <i>Megalograptus shideleri</i> Caster & Kjellesvig-Waering, 1964	O Ohio
74. <i>Megalograptus welchi</i> Miller, 1874*	O Ohio
75. <i>Megalograptus williamsae</i> Caster & Kjellesvig-Waering, 1964	O Ohio
† 'EURYPTEROIDEA' Burmeister, 1843	Ordovician – Devonian
NB: Lamsdell <i>et al.</i> (2013) questioned the monophyly of this superfamily	
Family uncertain	
† <i>Pentlandopterus</i> Lamsdell, Hoşgör & Selden, 2013	Ordovician
76. <i>Pentlandopterus minor</i> (Laurie, 1899)*	S Pentland Hills, Scotl.
† <i>Paraeurypterus</i> Lamsdell, Hoşgör & Selden, 2013	Ordovician
77. <i>Paraeurypterus anatoliensis</i> Lamsdell, Hoşgör & Selden, 2013*	O Şort Tepe, Turkey
† DOLICOPTERIDAE Kjellesvig-Waering & Størmer, 1952	Silurian – Devonian
† <i>Clarkeipterus</i> Kjellesvig-Waering, 1966 [a/b?]	Silurian
78. <i>Clarkeipterus ?otisius</i> (Clarke, 1907)	S eastern USA
79. <i>Clarkeipterus testudineus</i> (Clarke & Ruedeman, 1912)*	S New York
† <i>Dolichopterus</i> Hall, 1859	Silurian
80. <i>Dolichopterus gotlandicus</i> Kjellesvig-Waering, 1979	S Gotland, Sweden
81. <i>Dolichopterus jewetti</i> Caster & Kjellesvig-Waering, 1956	S New York
82. <i>Dolichopterus macrocheirus</i> Hall, 1859*	S New York / Canada
83. <i>Dolichopterus siluriceps</i> Clarke & Ruedemann, 1912	S New York / Canada
† <i>Ruedemannipterus</i> Kjellesvig-Waering, 1966	Silurian
84. <i>Ruedemannipterus stylonuroides</i> (Clarke & Ruedemann, 1912)*	S Otisville, New York
† EURYPTERIDAE Burmeister, 1843	Silurian
† <i>Eurypterus</i> de Kay, 1825	Silurian
= † <i>Baltoeurypterus</i> Størmer, 1973	
85. ? <i>Eurypterus cephalaspis</i> Salter, 1856	S Herefordshire, Engl.
86. <i>Eurypterus dekayi</i> Hall, 1859	S New York / Ontario
87. <i>Eurypterus flintstonensis</i> Swartz, 1923	S eastern USA
88. <i>Eurypterus hankeni</i> Tetlie, 2006a	S Ringerike, Norway
89. <i>Eurypterus henningsmoeni</i> (Tetlie, 2002)	S Bærum, Norway
90. <i>Eurypterus laculatus</i> Kjellesvig-Waering, 1958	S New York / Ontario
91. <i>Eurypterus lacustris</i> Harlan, 1834	S New York / Ontario
i. = <i>Eurypterus pachycheirus</i> Hall, 1859	S New York / Ontario
ii. = <i>Eurypterus robustus</i> Hall, 1859	S New York / Ontario
92. <i>Eurypterus leopoldi</i> Tetlie, 2006a	S Somerset Is., Canada

93. *Eurypterus megalops* Clarke & Ruedemann, 1912 S New York
94. *Eurypterus ornatus* Leutze, 1958 S Fayette, Ohio
95. *Eurypterus pittsfordensis* Sarle, 1903 S Pittsford, New York
96. *Eurypterus quebecensis* Kjellesvig-Waering, 1958 S Québec, Canada
97. *Eurypterus remipes* DeKay, 1825* S New York / Ontario
- i. = *Carcinosoma trigona* (Ruedemann, 1916) S New York
98. *Eurypterus serratus* (Jones & Woodward, 1888) S Gotland, Sweden
99. *Eurypterus tetragonophthalmus* Fischer, 1839 S Saaremaa, Estonia
- i. = *Eurypterus fischeri* Eichwald, 1854 S Estonia / Ukraine
- ii. = *Eurypterus fischeri* var. *rectangularis* Schmidt, 1883...S Saaremaa, Estonia
- † ERIEOPTERIDAE Tollerton, 1989** Silurian – Devonian
- † Erieopterus Kjellesvig-Waering, 1958** Silurian – Devonian
100. *Erieopterus eriensis* (Whitfield, 1882) S Ohio
101. *Erieopterus hypsophthalmus* Kjellesvig-Waering, 1958 S Ohio
102. ?*Erieopterus laticeps* (Schmidt, 1883) S Saaremaa, Ringerike
103. ?*Erieopterus limuloides* (Kjellesvig-Waering, 1948a) S Kokomo, Indiana
104. *Erieopterus microphthalmus* (Hall, 1859)* D New York / Canada
105. ?*Erieopterus phillipsensis* Copeland, 1971 S Cornwallis Is. Canada
106. ?*Erieopterus statzi* Størmer, 1936a D Siegburg, Germany
107. ?*Erieopterus turgidus* Stumm & Kjellesvig-Waering, 1962 S Michigan
- † STROBILOPTERIDAE Lamsdell & Selden, 2013** Silurian – Devonian
- † Buffalopterus Kjellesvig-Waering & Heubusch, 1962** Silurian
108. *Buffalopterus pustulosus* (Hall, 1859)* S New York / Ontario
- i. = *Eurypterus giganteus* Pohlman, 1882 S New York / Ontario
- ii. = *Pterygotus globicaudatus* Pohlman, 1882 S New York / Ontario
- † Strobilopterus Ruedemann, 1935** Silurian – Devonian
- = † *Syntomopterus* Kjellesvig-Waering, 1961 [preoccupied]
- = † *Syntomopterella* Tetlie, 2007 [replacement name]
109. *Strobilopterus laticeps* (Schmidt, 1883) S Saaremaa, Estonia
- i. = *Dolichopterus stoermeri* Caster & Kjellesvig-Waering,
1956 S Saaremaa, Estonia
110. *Strobilopterus princetonii* (Ruedemann, 1934)* D Wyoming, USA
- i. = *Erieopterus latus* Ruedemann, 1935 D Wyoming, USA
111. *Strobilopterus proteus* Lamsdell & Selden, 2013 D Wyoming, USA
112. *Strobilopterus richardsoni* (Kjellesvig-Waering, 1961a*) D Ohio
- † DIPLOPERCULATA Lamsdell, Hoşgör & Selden, 2013** Ordovician – Devonian
- † CARCINOSOMATOIDEA Størmer, 1934b** Ordovician – Devonian
- = † MIXOPTEROIDEA Caster & Kjellesvig-Waering, 1955
- † CARCINOSOMATIDAE Størmer, 1934b** Ordovician – Devonian

- † **Carcinosoma Claypole, 1890b** **Silurian**
- = † *Euryxoma* Claypole, 1890a [preoccupied]
113. ?*Carcinosoma harleyi* Kjellesvig-Waering, 1961b S England
114. *Carcinosoma libertyi* Copeland & Bolton, 1960 S Manitoulin I., Canada
115. *Carcinosoma newlini* (Claypole, 1890a)* S Kokomo, Indiana
- i. = *Carcinosoma ingens* Claypole, 1894 S Kokomo, Indiana
116. ?*Carcinosoma punctatum* (Salter in Huxley & Salter, 1859) S England
117. *Carcinosoma scorpioides* (Woodward, 1868) S Lesmahagow
- i. = *Pterygotus raniceps* Woodward, 1868 S Lesmahagow
118. *Carcinosoma scoticus* (Laurie, 1899) S Pentland Hills, Scotl.
119. ?*Carcinosoma spiniferum* Kjellesvig-Waering & Heubusch, 1962 S Pittsford, New York
- † **Eocarcinosoma Caster & Kjellesvig-Waering, 1964** **Ordovician**
120. *Eocarcinosoma batrachophthalmus* Caster & Kjellesvig-Waering,
1964* O Ohio
- † **Eusarcana Strand, 1942** **Silurian – Devonian**
- = † *Eusarcus* Grote & Pitt, 1875 [preoccupied]
- = † *Paracarcinosoma* Caster & Kjellesvig-Waering, 1964
121. *Eusarcana acrocephalus* (Semper, 1898) S–D Barrandian area
122. *Eusarcana obesus* (Woodward, 1868) S Lesmahagow
123. *Eusarcana scorpionis* (Grote & Pitt, 1875)* S New York / Ontario
- † **Rhinocarcinosoma Novojilov, 1962** **Silurian**
124. *Rhinocarcinosoma cicerops* (Clarke, 1907) S Otisville, New York
125. *Rhinocarcinosoma dosonensis* Braddy, Selden & Doan Nhat, 2002 S Dô Son, Vietnam
126. *Rhinocarcinosoma vaningeni* (Clarke & Ruedemann, 1912)* S Clinton, New York
- † **MIXOPTERIDAE Caster & Kjellesvig-Waering, 1955** **Silurian**
- = † LANARKOPTERIDAE Tollerton, 1989
- † **Lanarkopterus Ritchie, 1968** **Silurian**
127. *Lanarkopterus dolichoschelus* (Størmer, 1936b)* S Scotland
- † **Mixopterus Ruedemann, 1921** **Silurian**
128. *Mixopterus kiaeri* Størmer, 1934b S Ringerike, Norway
129. *Mixopterus multispinosus* (Clarke & Ruedemann, 1912)* S New York
130. *Mixopterus simonsoni* Schmidt, 1883 S Saaremaa, Estonia
- † ‘WAERINGOPTEROIDEA’ **Silurian – Devonian**
- NB: Superfamily name appears to be derived from a thesis; a family Waeringopteridae has not been formally published
- † **Grossopterus Størmer, 1934c** **Devonian**
131. *Grossopterus overathi* (Gross, 1933)* D Overath
132. *Grossopterus inexpectans* (Ruedemann, 1921) D Gilboa
- † **Orcanopterus Stott, Tetlie, Braddy, Nowlan, Glasser & Devereux, 2005** **Ordovician**
133. *Orcanopterus manitoulinensis* Stott, Tetlie, Braddy, Nowlan, Glasser

& Devereux, 2005*	O Manitoulin I., Canada
† <i>Waeringopterus</i> Leutze, 1961	Silurian
134. <i>Waeringopterus apfeli</i> Leutze, 1961	S New York / Ontario
135. <i>Waeringopterus cumberlandicus</i> (Swartz, 1923)*	S West Virginia
i. = <i>Eurypterus swartzi</i> Kjellesvig-Waering, 1958	S West Virginia
† ADELOPHTHALMOIDEA Tollerton, 1989	Devonian – Permian
† ADELOPHTHALMIDAE Tollerton, 1989	Devonian – Permian
† <i>Adelophthalmus</i> Jordan in Jordan & von Mayer, 1854	Devonian – Permian
= † <i>Lepidoderma</i> Reuss, 1855	
= † <i>Anthraconectes</i> Meek & Worthen, 1868 [a/b?]	
= † <i>Polyzosternites</i> Goldenberg, 1873	
= † <i>Glyptoscorpius</i> Peach, 1882	
136. <i>Adelophthalmus approximatus</i> (Hall & Clarke, 1888)	C Pennsylvania, USA
137. <i>Adelophthalmus asturica</i> (Melendez, 1971)	C d'Ablana, Spain
138. <i>Adelophthalmus bradorensis</i> (Bell, 1922)	C N. Campbelltown
139. <i>Adelophthalmus cambieri</i> (Pruvost, 1930)	C Charleroi, Belgium
140. ? <i>Adelophthalmus carbonarius</i> (Chernyshev, 1933)	C Donets, Ukraine
141. <i>Adelophthalmus chinensis</i> (Grabau, 1920)	C–P Zhaozhuang
142. <i>Adelophthalmus corneti</i> (Pruvost, 1939)	C Quaregnon, Belgium
143. <i>Adelophthalmus douvillei</i> (de Lima, 1890)	P Bussaco, Portugal
144. <i>Adelophthalmus dumonti</i> (Stainier, 1917)	C Mechelen-sur-Meuse
145. <i>Adelophthalmus granosus</i> Jordan in Jordan & von Meyer, 1854*	C Saarbrücken, Germ.
146. <i>Adelophthalmus imhofi</i> (Reuss, 1855)	C Vlkys, Czech Rep.
147. <i>Adelophthalmus irinae</i> Shpinev, 2006	C Krasnoyarsk, Russia
148. <i>Adelophthalmus kidstoni</i> (Peach, 1888)	C Radstock, England
149. ? <i>Adelophthalmus lohesti</i> (Dewalque in Fraipont 1889)	D Pont de Bonne, Belg.
150. <i>Adelophthalmus luceroensis</i> Kues & Kietzke, 1981	P New Mexico
151. <i>Adelophthalmus mansfieldi</i> (Hall, 1877)	C Pennsylvania
i. = <i>Eurypterus stylus</i> Hall, 1884	C Pennsylvania
152. <i>Adelophthalmus mazonensis</i> (Meek & Worthen, 1868)	C Illinois
153. <i>Adelophthalmus moyseyi</i> (Woodward, 1907a)	C Ilkeston, Blaengarw
i. = <i>Eurypterus derbiensis</i> Woodward, 1907a	C Ilkeston, England
154. <i>Adelophthalmus nebraskensis</i> (Barbour, 1914)	P Nebraska
155. <i>Adelophthalmus pennsylvanicus</i> (Hall, 1877)	C Pennsylvania
156. ? <i>Adelophthalmus perornatus</i> (Peach, 1882)	C Glencarholm, Scotl.
157. <i>Adelophthalmus pruvosti</i> Kjellesvig-Waering, 1948b	C Lens, France
158. <i>Adelophthalmus piussii</i> Lamsdell, Simonetto & Selden 2013	C Carnic Alps, Italy
159. ? <i>Adelophthalmus raniceps</i> Goldenberg, 1873	C Saarbrücken, Germ.
160. <i>Adelophthalmus sellardsi</i> (Dunbar, 1924)	P Elmo, Kansas
161. <i>Adelophthalmus sievertsi</i> (Størmer, 1969)	D Willwerath, Germ.
i. = ? <i>Eurypterus trapezoides</i> Størmer, 1974	D Nellenköpfchen, Ger.

162. *Adelophthalmus waterstoni* (Tetlie et al., 2004) D Kimberley, Australia
163. *Adelophthalmus wilsoni* (Woodward, 1888) C Radstock, England
164. *Adelophthalmus zadrai* Přibyl, 1952 C Moravo-Silesia
- † ***Bassipterus* Kjellesvig-Waering & Leutze, 1966** Silurian
165. *Bassipterus virginicus* Kjellesvig-Waering & Leutze, 1966* S Bass, West Virginia
- † ***Eysyslopterus* Tetlie & Poschmann, 2008** Silurian
166. *Eysyslopterus patteni* (Størmer, 1934d) S Saaremaa, Estonia
- † ***Nanahughmilleria* Kjellesvig-Waering, 1961b** Silurian – Devonian
167. *Nanahughmilleria clarkei* Kjellesvig-Waering, 1964b S Otisville, New York
168. *Nanahughmilleria norvegica* (Kiær, 1911)* S Ringerike, Norway
i. = *Eurypterus minutus* Kiær, 1911 S Ringerike, Norway
169. *Nanahughmilleria notosiberica* Shpinev, 2012 D Krasnoyarsk, Siberia
170. ?*Nanahughmilleria prominens* (Hall, 1884b) S Cayuga, New York
171. *Nanahughmilleria pygmaea* (Salter, 1859) S Herefordshire, Engl.
- 172.?*Nanahughmilleria schiraensis* (Pirozhnikov, 1957) D Khakassia, Russia
- † ***Parahughmilleria* Kjellesvig-Waering, 1961b** Silurian – Devonian
173. *Parahughmilleria bellistriata* (Kjellesvig-Waering, 1950a) S West Virginia
174. *Parahughmilleria hefteri* Størmer, 1973 D Rhenish Massif, Ge.
175. *Parahughmilleria longa* Shpiney, 2012 D Lake Shunet, Siberia
176. *Parahughmilleria maria* (Clarke, 1907) S New York
177. *Parahughmilleria matarakensis* (Pirozhnikov, 1957) D Khakassia, Russia
178. *Parahughmilleria salteri* Kjellesvig-Waering, 1961b* S Herefordshire, Engl.
- † ***Pittsfordipterus* Kjellesvig-Waering & Leutze, 1966** Silurian
179. *Pittsfordipterus phelpae* (Ruedemann, 1921)* S Pittsford, New York
- † **PTERYGOTIOIDEA Clarke & Ruedemann, 1912** Silurian – Devonian
- † **HUGHMILLERIIDAE Kjellesvig-Waering, 1951** Silurian
- † ***Herefordopterus* Tetlie, 2006b** Silurian
180. *Herefordopterus banksii* (Salter, 1856)* S Herefordshire, Engl.
i. = *Eurypterus acuminatus* Salter, 1859a S Herefordshire, Engl.
- † ***Hughmilleria* Sarle, 1903** Silurian
181. *Hughmilleria shawangunk* Clarke, 1907 S eastern USA
182. *Hughmilleria socialis* Sarle, 1903* S Pittsford, New York
i. = *Hughmilleria robusta* Sarle, 1903 S Pittsford, New York
183. *Hughmilleria wangi* Tetlie, Selden & Ren, 2007 S Hunan, China
- † **SLIMONIDAE Novojilov, 1968** Silurian
- † ***Salteropterus* Kjellesvig-Waering, 1951** Silurian
184. *Salteropterus abbreviatus* (Salter, 1859)* S Herefordshire, Engl.
- † ***Slimonia* Page, 1856** Silurian
185. *Slimonia acuminata* Salter, 1856* S Lesmahagow
i. = *Himantopterus maximus* Salter, 1856 S Lesmahagow

186. *Slimonia boliviensis* Kjellesvig-Waering, 1973 S Cochambamba, Bol.
187. *Slimonia dubia* Laurie, 1899 S Pentland Hills, Scotl.
- † **PTERYGOTIDAE Clarke & Ruedemann, 1912** **Silurian – Devonian**
- = † JAEKELOPTERIDAE Størmer, 1974
- † **Acutiramus Ruedemann, 1935** **Silurian – Devonian**
188. *Acutiramus bohemicus* (Barrande, 1872) S Barrandian area
- i. = *Pterygotus comes* Barrande, 1872 S Barrandian area
- ii. = *Pterygotus mediocris* Barrande, 1872 S Barrandian area
- iii. = *Pterygotus blahai* Semper, 1898 S Barrandian area
- iv. = *Pterygotus fissus* Seemann, 1906 S Barrandian area
189. *Acutiramus cummingsi* (Grote & Pitt, 1875) S USA / Canada
- i. = *Pterygotus acuticaudatus* Pohlman, 1882 S New York
- ii. = *Pterygotus buffaloensis* Pohlman, 1881 S New York
- iii. = *Pterygotus quadraticaudatus* Pohlman, 1882 S New York
190. *Acutiramus floweri* Kjellesvig-Waering & Caster, 1955 S Kenwood, New York
191. *Acutiramus macrophthalmus* (Hall, 1859)* S USA / Canada
- i. = *Pterygotus osborni* Hall, 1859 S New York
- ii. = *Pterygotus cobbi* var. *juvenis* Clarke & Ruedemann,
1912 S New York
192. *Acutiramus perneri* Chlupáč, 1994 D Barrandian area
193. *Acutiramus perryensis* Leutze, 1958 S Ohio
194. *Acutiramus suwanneensis* Kjellesvig-Waering, 1955 S? Florida
- † **Ciurcopterus Tetlie & Briggs, 2009** **Silurian**
195. *Ciurcopterus sarlei* (Ciurca & Tetlie, 2007) S Pittsford, New York
196. *Ciurcopterus ventricosus* (Kjellesvig-Waering, 1948a)* S Kokomo, Indiana
- † **Erettopterus Salter in Huxley & Salter, 1859** **Silurian – Devonian**
- = † *Truncatiramus* Kjellesvig-Waering, 1961b
197. *Erettopterus bilobus* (Salter, 1856)* S Lesmahagow
- i. = *Eurypterus perornatus* Salter, 1856 S Lesmahagow
- ii. = *Pterygotus bilobus* var. *acidens* Woodward, 1878 S Lesmahagow
- iii. = *Pterygotus bilobus* var. *crassus* Woodward, 1878 S Lesmahagow
- iv. = *Pterygotus bilobus* var. *inornatus* Woodward, 1878 S Lesmahagow
- v. = *Pterygotus bilobus* var. *perornatus* Woodward, 1878 S Lesmahagow
- vi. = *Pterygotus perornatus* var. *plicatissimus* Salter in
Huxley & Salter, 1859 S Lesmahagow
198. *Erettopterus brodiei* Kjellesvig-Waering, 1961b S Herefordshire, Engl.
199. *Erettopterus canadensis* (Dawson, 1879) S Ontario, Canada
200. *Erettopterus exophthalmus* Kjellesvig-Waering & Leutze, 1966 S Bass, West Virginia
201. *Erettopterus gigas* Salter in Huxley & Salter, 1859 S Herefordshire, Engl.
202. *Erettopterus globiceps* Clarke & Ruedemann, 1912 S eastern USA
203. *Erettopterus grandis* Pohlman, 1881 S New York

204. *Erettopterus holmi* (Størmer, 1934b) S Ringerike, Norway
205. *Erettopterus laticauda* Schmidt, 1883 S Saaremaa, Estonia
206. *Erettopterus marstoni* Kjellesvig-Waering, 1961b S England
207. *Erettopterus megalodon* Kjellesvig-Waering, 1961b S England
208. *Erettopterus osiliensis* Schmidt, 1883 S Saaremaa, Estonia
209. *Erettopterus saetiger* Kjellesvig-Waering, 1964a S Pennsylvania
210. *Erettopterus serratus* Kjellesvig-Waering, 1961b D Ohio
211. *Erettopterus spatulatus* Kjellesvig-Waering, 1961b S Herefordshire, Engl.
212. ?*Erettopterus vogti* Størmer, 1934a D Spitsbergen
213. *Erettopterus waylandsmithi* Kjellesvig-Waering & Caster, 1955 S Kenwood, New York
- † ***Jaekelopterus* Waterston, 1964** **Devonian**
214. *Jaekelopterus howelli* Kjellesvig-Waering & Størmer, 1952 D Wyoming
- i. = *Pterygotus mcgrewi* Kjellesvig-Waering & Richardson
In Kjellesvig-Waering (1986) [nomen nudum] D Wyoming
215. *Jaekelopterus rhenanae* (Jaekel, 1914)* D Rhenish Massif, Ger.
- † ***Necrogammarus* Woodward, 1870** **Silurian**
216. *Necrogammarus salweyi* Woodward, 1870 S Herefordshire, Engl.
- † ***Pterygotus* Agassiz, 1839** **Silurian – Devonian**
- = † *Curviramus* Reudemann, 1935
217. *Pterygotus anglicus* Agassiz, 1844* D Scotland, Canada
- i. = *Pterygotus atlanticus* Clarke & Ruedemann, 1912 D New Brunswick, Can.
- ii. = *Pterygotus minor* Woodward, 1864 D Scotland
218. *Pterygotus arcuatus* Salter *in* Huxley & Salter, 1859 S Herefordshire, Engl.
219. ?*Pterygotus australis* McCoy, 1899 S Melbourne, Australia
220. *Pterygotus barrandei* Semper, 1898 S Barrandian area
- i. = *Pterygotus beraunensis* Semper, 1898 S Barrandian area
221. *Pterygotus bolivianus* Kjellesvig-Waering, 1964a D Belen, Bolivia
222. *Pterygotus carmani* Kjellesvig-Waering, 1961 D Ohio
223. *Pterygotus cobbi* Hall, 1859 S New York / Canada
224. *Pterygotus denticulatus* Kjellesvig-Waering, 1961b S Herefordshire, Engl.
225. *Pterygotus floridanus* Kjellesvig-Waering, 1950b D Florida
226. *Pterygotus gaspesiensis* Russell, 1953 D Québec, Canada
227. ?*Pterygotus grandidentatus* Kjellesvig-Waering, 1961b S England
228. ?*Pterygotus impacatus* Kjellesvig-Waering, 1964a S Saaremaa, Estonia
229. *Pterygotus kopaninensis* Barrande, 1872 S Barrandian area, Cz.
230. *Pterygotus lanarkensis* Kjellesvig-Waering, 1964a S Lesmahagow, Scotl.
231. *Pterygotus lightbodyi* Kjellesvig-Waering, 1961b S England
232. *Pterygotus ludensis* Salter *in* Huxley & Salter, 1859 S Herefordshire, Engl.
233. *Pterygotus marylandicus* Kjellesvig-Waering, 1964a S Maryland
234. *Pterygotus monroensis* Sarle 1902 S New York

EURYPTERIDA incertae sedis

- † *Dorfopterus* Kjellesvig-Waering, 1955 Devonian
235. *Dorfopterus angusticollis* Kjellesvig-Waering, 1955* D Wyoming
- † ?*Dolichopterus*
236. ?*Dolichopterus asperatus* Kjellesvig-Waering, 1961 [a/b?] D Ohio
237. ?*Dolichopterus bulbosus* Kjellesvig-Waering, 1961b S Herefordshire, Engl.
238. ?*Dolichopterus herkimerensis* Caster & Kjellesvig-Waering, 1956 S New York / Canada
- † ?*Eurypterus*
239. ?*Eurypterus loi* Chang, 1957 [non eurypterid?] S Hubei, China
240. ?*Eurypterus podolicus* Chernyshev, 1947 S Ukraine
241. ?*Eurypterus satpaevi* Simorin, 1956 C Karaganda, Kazakh.
242. ?*Eurypterus styliformis* Chang, 1957 [non eurypterid?] S Hubei, China
243. ?*Eurypterus tschernyschevi* Simorin, 1956 C Karaganda, Kazakh.
244. ?*Eurypterus yangi* Chang, 1957 [non eurypterid?] S Hubei, China
- † *Holmipterus* Kjellesvig-Waering, 1979 Silurian
245. *Holmipterus suecicus* Kjellesvig-Waering, 1979 S Gotland, Sweden
- † *Marsupipterus* Caster & Kjellesvig-Waering, 1955 Silurian
246. *Marsupipterus sculpturatus* Caster & Kjellesvig-Waering, 1955* S Herefordshire, Engl.
- † ?*Nanahughmilleria*
247. ?*Nanahughmilleria lanceolata* Salter, 1856 S Lesmahagow
- i. = *Eurypterus chartarius* Salter, 1859 S Lesmahagow
- ii. = *Eurypterus linearis* Salter, 1859 S Lesmahagow
- † ?*Salteropterus*
248. ?*Salteropterus longilabium* Kjellesvig-Waering, 1961b S Welsh Borderlands
- † ?*Stylonurus*
249. ?*Stylonurus perspicillum* Størmer, 1969 D Willwerath, Germany
- † *Unionopterus* Chernyshev, 1948 Carboniferous
250. *Unionopterus anastasiae* Chernyshev, 1948* C Kazakhstan

NOMINA DUBIA

1. *Bunodella horrida* Matthew, 1888 [non Xiphosura] S New Brunswick
2. ?*Dunsopterus wrightianus* Dawson 1881 D New York
3. *Eurypterella ornata* Matthew, 1888 C 'Fern Ledges'
4. *Eurypterus potens* Hall, 1884 C Pennsylvania
5. *Eurypterus pulicaris* Salter, 1863 D New Brunswick
6. *Hastimima sewardi* Strand, 1926 D South Africa
7. ?*Pterygotus formosus* Dawson, 1871 D Gaspé, Canada
8. *Pterygotus nobilis* Barrande, 1872 S Barrandian area
9. *Pterygotus siemiradzkii* Strand, 1926 D Podolia, Ukraine
10. *Pterygotus taurinus* Salter, 1868 S Ewyas Harold, Engl.
11. ?*Slimonia stylops* Salter in Huxley & Salter, 1859 S Herefordshire, Engl.

NOMINA NUDA

1. *Baltoeurypterus latus* Hanken & Størmer, 1975 S Ringerike, Norway

NOMINA VANA

1. *Pterygotus problematicus* Agassiz, 1844 S United Kingdom

MISIDENTIFICATIONS

1. *Buffalopterus verrucosus* Kjellesvig-Waering & Heubusch, 1962 [crustacean] ... O New York
2. *Carcinosoma ?logani* (Williams, 1915) [crustacean] S Ontario, Canada
3. *Eurypterus (Stylonurus?) maccarthyi* Kjellesvig-Waering, 1934 [cephalopod] D Ludlowville, New York
4. *Eurypterus pugio* Barrande, 1872 [crustacean] S Barrandian area
5. *Eurypterus thomasi* Walter, 1924 [aglaspidid] E Wisconsin
6. *Kockurus grandis* Chlupáč, 1995 [?aglaspidid] E central Bohemia
7. *Kodymirus vagans* Chlupáč & Havlíček, 1965 [?aglaspidid] E central Bohemia
8. *Mazonipterus cyclophthalmus* Kjellesvig-Waering, 1963b [plant] C Mazon Creek
9. *Melbournopterus crossotus* Caster & Kjellesvig-Waering, 1953 [brachiopod] ... S Melbourne, Australia
10. *Pterygotus expectatus* Barrande, 1872 [crustacean] S Barrandian area
11. *Pterygotus (Curvirostrum) elliotti* Ruedemann, 1935 [crustacean] D New York
12. *Pterygotus (Curvirostrum) montanensis* Ruedemann, 1935 [crustacean] D Montana
13. *Pterygotus (Leptocheles) leptodactylum* M'Coy, 1849 [crustacean] S Herefordshire, Engl.

PSEUDOFOSSILS

1. *Brachypterella magna* (Clarke & Ruedemann, 1912) O New York
2. ?*Carcinosoma linguata* (Clarke & Ruedemann, 1912) O New York
3. ?*Carcinosoma longiceps* (Clarke & Ruedemann, 1912) O New York
4. *Dolichopterus antiquus* Ruedemann, 1942 O New York
5. *Dolichopterus frankfortensis* (Clarke & Ruedemann, 1912) O New York
6. *Dolichopterus insolitus* Ruedemann, 1926 O New York
7. ?*Dolichopterus stellatus* (Clarke & Ruedemann, 1912) O New York
8. ?*Drepanopterus ruedemanni* (O'Connell, 1916) O New York
9. ?*Eocarcinosoma breviceps* (Ruedemann, 1926) O New York
10. *Eocarcinosoma ruedemanni* (Flower, 1945) O New York
11. *Eocarcinosoma triangulatus* (Clarke & Ruedemann, 1912) O New York
12. *Erettopterus walcotti* (Ruedemann, 1926) O New York
13. *Erieopterus chadwicki* (Clarke & Ruedemann, 1912) O New York
14. *Erieopterus hudsonicus* (Ruedemann, 1934) O New York
15. ?*Eurypterus deceptiens* (Ruedemann, 1942) O New York
16. *Eurypterus indicus* Dubey, 1985 pC M. Pradesh, India
17. ?*Eurypterus pristinus* (Clarke & Ruedemann, 1912) O New York
18. *Eurypterus vermai* Dubey, 1985 pC M. Pradesh, India
19. *Hughmilleria chiplonkari* Dubey, 1985 pC M. Pradesh, India
20. *Hughmilleria kilfoylei* Ruedemann, 1934 O New York

21. *Hughmilleria prisca* Ruedemann, 1934 O New York
22. *Hughmilleria uticana* Ruedemann, 1926 O New York
23. *Parastylonurus rusti* (Ruedemann, 1926) O New York
24. *Pterygotus deepkillensis* Ruedemann, 1934 O New York
25. *Pterygotus nasutus* Clarke & Ruedemann, 1912 O New York
26. ?*Pterygotus normanskilensis* Clarke & Ruedemann, 1912 O New York
27. *Ruedemannipterus breviceps* (Clarke & Ruedemann, 1912) O New York
28. *Ruedemannipterus latifrons* (Clarke & Ruedemann, 1912) O New York
29. *Styloceras modestus* (Clarke & Ruedemann, 1912) O New York
30. *Styloceras limbatus* (Clarke & Ruedemann, 1912) O New York
31. ?*Waeringopterus pristinus* (Ruedemann, 1942) O New York
32. *Waeringopterus prolificus* (Clarke & Ruedemann, 1912) O New York

no Recent species

SCORPIONES

118 currently valid species of fossil scorpion

SCORPIONES C. L. Koch, 1851 Silurian – Recent

† **Plesion** (Family) PROSCORPIIDAE Scudder, 1885 Silurian – Carbon.

- = † ARCHAEOCTONIDAE Petrunkevitch, 1949
- = † HYDROSCORPIONIDAE Kjellesvig-Waering, 1986
- = † LABRIOSCORPIONIDAE Kjellesvig-Waering, 1986
- = † STOERMEROSCORPIONIIDAE Kjellesvig-Waering, 1986
- = † WAERINGOSCORPIONIDAE Størmer, 1970

† **Archaeoctonus** Pocock, 1911 Carboniferous

- 1. *Archaeoctonus glaber* (Peach, 1883)* C Glencarholm

† **Hydroscorpius** Kjellesvig-Waering, 1986 Devonian

- 2. *Hydroscorpius denisoni* Kjellesvig-Waering, 1986* D Wyoming

† **Labriscorpio** Leary, 1980 Carboniferous

- 3. *Labriscorpio alliedensis* Leary, 1980* C Illinois

† **Proscorpius** Whitfield, 1885b Silurian

- = † *Archaeophonus* Kjellesvig-Waering, 1966b
- = † *Stoermeroscorpio* Kjellesvig-Waering, 1986
- 4. *Proscorpius osborni* (Whitfield, 1885a)* S ‘Bertie Waterlime’
 - i. = *Archaeophonus eurypterooides* Kjellesvig-Waering, 1966b* S ‘Bertie Waterlime’
 - ii. = *Stoermeroscorpio delicatus* Kjellesvig-Waering, 1986 S ‘Bertie Waterlime’

† **Pseudoarchaeoctonus** Kjellesvig-Waering, 1986 Carboniferous

- 5. *Pseudoarchaeoctonus denticulatus* Kjellesvig-Waering, 1986* C Glencarholm

† **Waeringoscorpio** Størmer, 1970 Devonian

- 6. *Waeringoscorpio hefteri* Størmer, 1970* D Alken an der Mosel
- 7. *Waeringoscorpio westerwaldensis* Poschmann, Dunlop, Kamenz & Scholtz, 2008 D Westerwald

† **BILOBOSTERNINA** Kjellesvig-Waering, 1986 (suborder) Silurian – Devonian

† **BRANCHIOSCORPINOIDEA** Kjellesvig-Waering, 1986 Devonian

† **BRANCHIOSCORPIONIIDAE** Kjellesvig-Waering, 1986 Devonian

† **Branchioscorpio** Kjellesvig-Waering, 1986 Devonian

- 8. *Branchioscorpio richardsoni* Kjellesvig-Waering, 1986* D Wyoming

† **DOLICHOPHONIIDAE** Petrunkevitch, 1953 Silurian

† **Dolichophonus** Petrunkevitch, 1949 Silurian

9. <i>Dolichophonus loudonensis</i> (Laurie, 1899)*	S Pentland Hills
† HOLOSTERNINA Kjellesvig-Waering, 1986	Devonian
† ACANTHOSCORPINOIDEA Kjellesvig-Waering, 1986	Devonian
† ACANTHOSCORPONIIDAE Kjellesvig-Waering, 1986	Devonian
† <i>Acanthocorpio</i> Kjellesvig-Waering, 1986	Devonian
10. <i>Acanthoscorpio mucronatus</i> Kjellesvig-Waering, 1986*	D Wyoming
† STENOSCORPONIIDAE Kjellesvig-Waering, 1986	Triassic
† <i>Stenoscorpio</i> Kjellesvig-Waering, 1986	Triassic
11. <i>Stenoscorpio gracilis</i> (Wills, 1910)*	Tr Keuper sandstone
12. <i>Stenoscorpio pseudogracilis</i> (Wills, 1947)	Tr Keuper sandstone
† ALLOPALAEOPHONOIDEA Kjellesvig-Waering, 1986	Silurian
† ALLOPALAEOPHONIDAE Kjellesvig-Waering, 1986	Silurian
† <i>Allopalaeophonus</i> Kjellesvig-Waering, 1986	Silurian
13. <i>Allopalaeophonus caledonicus</i> (Hunter, 1886)*	S Logan Water
i. = <i>Palaeophonus hunteri</i> Pocock, 1901	S Logan Water
† EOCTONOIDAE Kjellesvig-Waering, 1986	Carboniferous
† ALLOBUTHISCORPIIIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Aspiscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
14. <i>Aspiscorpio eageri</i> Kjellesvig-Waering, 1986*	C Sparth Bottoms
<i>Aspiscorpio</i> sp. in Poschmann (2009)	C Saar
† ANTHRACOSCORPIONIDAE Frič, 1904	Carboniferous
† <i>Allobuthus</i> Kjellesvig-Waering, 1986	Carboniferous
15. <i>Allobuthus pescei</i> (Vachon & Heyler, 1985)*	C Montceau-les-Mines
† Anthracoscorpio Kušta, 1885	Carboniferous
16. <i>Anthracoscorpio dunlopi</i> Pocock, 1911	C Airdrie
17. <i>Anthracoscorpio juvenis</i> Kušta, 1885*	C Rakovník
† BUTHISCORPIIIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Buthiscorpis</i> Petrunkevitch, 1953	Carboniferous
18. <i>Buthiscorpis lemayi</i> Kjellesvig-Waering, 1986	C Illinois
† EOCTONIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Eoconus</i> Petrunkevitch, 1913	Carboniferous
19. <i>Eoconus miniatus</i> Petrunkevitch, 1913*	C Mazon Creek
† GARNETTIIDAE Dubinin, 1962	Carboniferous
† <i>Garnettius</i> Petrunkevitch, 1953	Carboniferous

20. *Garnettius hungerfordi* (Elias, 1936)* C Garnett, Kansas
- † **GIGANTOSCORPIONOIDEA** Kjellesvig-Waering, 1986 Devonian – Carbon.
- † **GIGANTOSCORPIONIDAE** Kjellesvig-Waering, 1986 Devonian – Carbon.
= † *PETALOSCORPIONIDAE* Kjellesvig-Waering, 1986
- † **Gigantoscorpio** Størmer, 1963 Carboniferous
21. *Gigantoscorpio willsi* Størmer, 1963* C Glencarholm
- † **Petaloscorpio** Kjellesvig-Waering, 1986 Devonian
22. *Petaloscorpio bureaui* Kjellesvig-Waering, 1986* D Miguasha, Quebec
- † **MESOPHONOIDEA** Wills, 1910 Carbon. – Triassic
- † **CENTROMACHIDAE** Petrunkevitch, 1953 Carboniferous
- = † *ANTHRACOCHAERILIDAE* Kjellesvig-Waering, 1986
= † *PHOXISCORPIONIDAE* Kjellesvig-Waering, 1986
- † **Anthracochaerilus** Kjellesvig-Waering, 1986 Carboniferous
23. *Anthracochaerilus palustris* Kjellesvig-Waering, 1986* C Glencarholm
- † **Centromachus** Thorell & Lindström, 1885 Carboniferous
24. *Centromachus euglyptus* (Peach, 1883)* C Glencarholm
- † **Phoxiscorpio** Kjellesvig-Waering, 1986 Carboniferous
25. *Phoxiscorpio peachi* Kjellesvig-Waering, 1986* C Dalmeny, Edinburgh
- † **Pulmonoscorpio** Jeram, 1994a Carboniferous
26. *Pulmonoscorpius kirktonensis* Jeram, 1994a* C East Kirkton
- † **GALLIOSCORPIONIDAE** Lourenço & Gall, 2004 Triassic
- † **Gallioscorpio** Lourenço & Gall, 2004 Triassic
27. *Gallioscorpio voltzi* Lourenço & Gall, 2004* Tr Vosges, France
- † **HELOSCORPIONIDAE** Kjellesvig-Waering, 1986 Carboniferous
- † **Heloscorpio** Kjellesvig-Waering, 1986 Carboniferous
28. *Heloscorpio sutcliffei* (Woodward, 1907b)* C Sparth Bottoms
- † **MAZONIIDAE** Petrunkevitch, 1913 Carboniferous
- † **Mazonia** Meek & Worthen, 1868b Carboniferous
29. *Mazonia wardingleyi* (Woodward, 1907b) C Sparth Bottoms
30. *Mazonia woodiana* Meek & Worthen, 1868b* C Mazon Creek
- † **MESOPHONIDAE** Wills, 1910 Triassic
- † **Mesophonus** Wills, 1910 Triassic
31. *Mesophonus perornatus* Wills, 1910* Tr Keuper sandstone
i. = *Mesophonus opistophthalmus* Wills, 1947 Tr Keuper sandstone
32. ?*Mesophonus pulcherrimus* Wills, 1910 Tr Keuper sandstone
33. ?*Mesophonus pulcherrimus immaculatus* Wills, 1947 Tr Keuper sandstone

† WILLISCORPIONIDAE Kjellesvig-Waering, 1986	Triassic
† <i>Willsiscorpio</i> Kjellesvig-Waering, 1986	Triassic
34. <i>Willsiscorpio bromsgroviensis</i> (Wills, 1910)*	Tr Keuper sandstone
† PALAEOSCORPOIDEA Lehmann, 1944	Devonian – Triassic
† PALAEOSCORPIONIDAE Lehmann, 1944	Devonian
† <i>Palaeoscorpio</i> Lehmann, 1944	Devonian
35. <i>Palaeoscorpius devonicus</i> Lehmann, 1944*	D Hünsruckschiefer
[NB: Kühl et al. (2012) simply list the genus unplaced under Protoscorpionina.]	
† SPONGIOPHONOIDEA Kjellesvig-Waering, 1986	Devonian – Triassic
† PRAERCTURIDAE Kjellesvig-Waering, 1986	Devonian
† <i>Praearcturus</i> Woodward, 1871a	Devonian
36. <i>Praearcturus gigas</i> Woodward, 1871a*	D Rowlestone
† SPONGIOPHONIDAE Kjellesvig-Waering, 1986	Triassic
† <i>Spongiphonus</i> Wills, 1947	Triassic
37. <i>Spongiphonus pustulosus</i> Wills, 1947*	Tr Keuper sandstone
† MERISTOSTERNINA Kjellesvig-Waering, 1986	Carboniferous
† CYCLOPHTHALMOIDEA Thorell & Lindström, 1885	Carboniferous
† CYCLOPHTHALMIDAE Thorell & Lindström, 1885	Carboniferous
† <i>Cyclophthalmus</i> Corda, 1835	Carboniferous
38. <i>Cyclophthalmus senior</i> Corda, 1835*	C Cholme
39. <i>Cyclophthalmus robustus</i> Kjellesvig-Waering, 1986	C Coseley
40. ? <i>Cyclophthalmus sibiricus</i> Novojilov & Størmer, 1963	C Kemerov Region
† MICROLABIIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Microlabis</i> Corda, 1839	Carboniferous
41. <i>Microlabis sternbergii</i> Corda, 1839*	C Cholme
† PALAEOBUTHOIDEA Kjellesvig-Waering, 1986	Carboniferous
† PALAEOBUTHIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Palaeobuthus</i> Petrunkevitch, 1913	Carboniferous
= † <i>Mazoniscorpio</i> Wills, 1960	
42. <i>Palaeobuthus distinctus</i> Petrunkevitch, 1913*	C Mazon Creek
i. = <i>Mazoniscorpio mazonensis</i> Wills, 1960	C Mazon Creek
† LOBOSTERNINA Pocock, 1911	Silurian – Carbon.
† ISOBUTHOIDEA Petrunkevitch, 1913	Carboniferous
† EOBUTHIDAE Kjellesvig-Waering, 1986	Carboniferous

† <i>Eobuthus</i> Frič, 1904	Carboniferous
43. <i>Eobuthus cordai</i> Kjellesvig-Waering, 1986	C Kralupy Hill
44. <i>Eobuthus holti</i> Pocock, 1911	C Sparth Bottoms
45. <i>Eobuthus rakovicensis</i> Frič, 1904*	C Rakovník
 † EOSCORPIIDAE Scudder, 1884	Carboniferous
† <i>Eoscorpius</i> Meek & Worthen, 1868a	Carboniferous
= † <i>Alloscorpius</i> Petrunkevitch, 1949	
= † <i>Europhthalmus</i> Petrunkevitch, 1949	
= † <i>Lichnophthalmus</i> Petrunkevitch, 1949	
= † <i>Trigonoscorpio</i> Petrunkevitch, 1913	
= † <i>Typhloscorpius</i> Petrunkevitch, 1949	
46. <i>Eoscorpius bornaensis</i> Sterzel, 1918	C Chemnitz–Borna
47. <i>Eoscorpius carbonarius</i> Meek & Worthen, 1868a*	C Mazon Creek
i. = <i>Eoscorpius typicus</i> Petrunkevitch, 1913	C Mazon Creek
ii. = <i>Eoscorpius granulosus</i> Petrunkevitch, 1913	C Mazon Creek
iii. = <i>Trigonoscorpio americanus</i> Petrunkevitch, 1913	C Mazon Creek
48. <i>Eoscorpius casei</i> Kjellesvig-Waering, 1986	C Nova Scotia
49. <i>Eoscorpius distinctus</i> (Petrunkevitch, 1949)	C Coseley
50. <i>Eoscorpius mucronatus</i> Kjellesvig-Waering, 1986	C Barnsley
51. <i>Eoscorpius pulcher</i> (Petrunkevitch, 1949)	C Barnsley
i. = <i>Europhthalmus longimanus</i> Petrunkevitch, 1949	C Barnsley
52. <i>Eoscorpius sparthensis</i> Baldwin & Sutcliffe, 1904	C Sparth Bottoms
† <i>Eskioscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
53. <i>Eskiscorpio parvus</i> Kjellesvig-Waering, 1986*	C Glencarholm
† <i>Trachyscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
54. <i>Trachyscorpio squarrosum</i> Kjellesvig-Waering, 1986*	C Fouldon
 † ISOBUTHIDAE Petrunkevitch, 1913	Carbon. – Triassic
† <i>Boreoscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
55. <i>Boreoscorpio copelandi</i> Kjellesvig-Waering, 1986*	C Nova Scotia
† <i>Bromsgroviscorpio</i> Kjellesvig-Waering, 1986	Triassic
56. <i>Bromsgroviscorpio willsi</i> Kjellesvig-Waering, 1986*	Tr Keuper sandstone
† <i>Feistmantelia</i> Frič, 1904	Carboniferous
57. <i>Feistmantelia ornata</i> Frič, 1904*	C Studhoves
† <i>Isobuthus</i> Frič, 1904	Carboniferous
58. <i>Isobuthus kralupensis</i> (Thorell & Lindström, 1885)*	C Kralup
59. ? <i>Isobuthus nyranensis</i> Frič, 1904	C Nýřany
 † KRONOSCORPIONIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Kronoscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
60. <i>Kronoscorpio danielsi</i> (Petrunkevitch, 1913)*	C Mazon Creek

† PAREOBUTHIDAE Wills, 1959	Carboniferous
† <i>Pareobuthus</i> Wills, 1959	Carboniferous
61. <i>Pareobuthus salopiensis</i> Wills, 1959*	C Shropshire
† PARAISOBUTHOIDEA Kjellesvig-Waering, 1986	Carboniferous
† OPSIEOBUTHIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Opsieobuthus</i> Kjellesvig-Waering, 1986	Carboniferous
62. <i>Opsieobuthus pottsvilleensis</i> (Moore, 1923)*	C Indiana
† PARAISOBUTHIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Paraisobuthus</i> Kjellesvig-Waering, 1986	Carboniferous
63. <i>Paraisobuthus duobicarinatus</i> Kjellesvig-Waering, 1986	C Shipley
64. <i>Paraisobuthus frici</i> Kjellesvig-Waering, 1986	C Kralupy Hill
65. <i>Paraisobuthus prantli</i> Kjellesvig-Waering, 1986*	C Rakovník
66. <i>Paraisobuthus virginiae</i> Kjellesvig-Waering, 1986	C Mazon Creek
† SCOLOPOSCORPIONIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Benniescorpio</i> Wills, 1960	Carboniferous
67. <i>Benniescorpio tuberculatus</i> (Peach, 1883)*	C Dysart, Fife
† <i>Scoloposcorpio</i> Kjellesvig-Waering, 1986	Carboniferous
68. <i>Scoloposcorpio crumondensis</i> Kjellesvig-Waering, 1986*	C Cramond, Edinburgh
† TELMATOSCORPIONIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Telmatoscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
69. <i>Telmatoscorpio brevipectus</i> Kjellesvig-Waering, 1986*	C Mazon Creek
† LOBOARCHAEOTONOIDEA Kjellesvig-Waering, 1986	Carboniferous
† LOBOARCHAEOTONIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Loboarchaeoctonus</i> Kjellesvig-Waering, 1986	Carboniferous
70. <i>Loboarchaeoctonus squamosus</i> Kjellesvig-Waering, 1986*	C Glencarholm
† WATERSTONIIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Waterstonia</i> Kjellesvig-Waering, 1986	Carboniferous
71. <i>Waterstonia airdriensis</i> Kjellesvig-Waering, 1986*	C Airdrie
† PALAEOPHONOIDEA Thorell & Lindström, 1884	Silurian
† PALAEOPHONIDAE Thorell & Lindström, 1884	Silurian
† <i>Palaeophonus</i> Thorell & Lindström, 1884	Silurian
72. <i>Palaeophonus nuncius</i> Thorell & Lindström, 1884*	S Visby, Gotland
73. ? <i>Palaeophonus lightbodyi</i> Kjellesvig-Waering, 1954 [claw only!]	S Ludford Lane

ORTHOSTERNINA Pocock, 1911	Carbon. – Recent
Orthosternina incertae sedis	
† <i>Corniops</i> Jeram, 1994b	Carboniferous
74. <i>Corniops mapesii</i> Jeram, 1994b*	C Lone Star Lake
 SCORPIONIOIDEA Latreille, 1802	Carbon. – Recent
† PALAEOPISTHACANTHIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Cryptoscorpius</i> Jeram, 1994b	Carboniferous
75. <i>Cryptoscorpius americanus</i> Jeram, 1994b*	C Lone Star Lake
 † Palaeopisthacanthus Petrunkevitch, 1913	Carboniferous
76. <i>Palaeopisthacanthus schucherti</i> Petrunkevitch, 1913*	C Mazon Creek
77. <i>Palaeopisthacanthus vogelandurdeni</i> Jeram, 1994b	C Lone Star Lake
 family uncertain	
† Compsoscorpius Petrunkevitch 1949	Carboniferous
= † <i>Allobuthiscorpius</i> Kjellesvig-Waering, 1986	
= † <i>Coseleyscorpio</i> Kjellesvig-Waering, 1986	
= † <i>Leioscorpio</i> Kjellesvig-Waering, 1986	
= † <i>Lichnoscorpius</i> Petrunkevitch, 1949	
= † <i>Pseudobuthiscorpius</i> Kjellesvig-Waering, 1986	
= † <i>Typhlopisthacanthus</i> Petrunkevitch, 1949	
78. <i>Compsoscorpius buthiformis</i> (Pocock, 1911)*	C Coal Measures
i. = <i>Typhlopisthacanthus anglicus</i> Petrunkevitch, 1949 ... C Coseley	
ii. = <i>Lichnoscorpius minutus</i> Petrunkevitch, 1949 C Coseley	
iii. = <i>Compsoscorpius elegans</i> Petrunkevitch 1949 C Coseley	
iv. = <i>Compsoscorpius elongatus</i> Petrunkevitch, 1949 C Coseley	
v. = <i>Buthiscorpius major</i> Wills, 1960 C Kilburn Coal	
vi. = <i>Leioscorpio pseudobuthiformis</i> Kjellesvig-Waering, 1986 C Coseley	
vii. = <i>Pseudobuthiscorpius labiosus</i> Kjellesvig-Waering, 1986 C Coseley	
viii. = <i>Coseleyscorpio lanceolatus</i> Kjellesvig-Waering, 1986 C Coseley	
ix. = <i>Allobuthus macrostethus</i> Kjellesvig-Waering, 1986C Coseley	
 PSEUDOCHACTIDAE Gromov, 1998	Recent
no fossil record	
 BUTHOIDEA C. L. Koch, 1837	Triassic – Recent
family uncertain	
† Palaeoburmesebuthus Lourenço, 2002	Cretaceous
79. <i>Palaeoburmesebuthus grimaldii</i> Lourenço, 2002*	K Myanmar amber

† ARCHAEOBUTHIDAE Lourenço, 2001	Cretaceous
† <i>Archaeobuthus</i> Lourenço, 2001	Cretaceous
80. <i>Archaeobuthus estephani</i> Lourenço, 2001*	K Lebanese amber
† PROTOBUTHIDAE Lourenço & Gall, 2004	Triassic
† <i>Protobuthus</i> Lourenço & Gall, 2004	Triassic
81. <i>Protobuthus elegans</i> Lourenço & Gall, 2004*	Tr Vosges
BUTHIDAE C. L. Koch, 1837	Palaeogene – Recent
= ANDROCTONIDAE C. L. Koch, 1837	
= MICROCHARMIDAE Lourenço, 1996a	
Centruroides Marx, 1890a	Neogene – Recent
82. <i>Centruroides nitidus</i> (Thorell, 1876a) [Recent]	Ne Dominican amber
i. = <i>Centruroides beynai</i> Schawaller, 1979a	Ne Dominican amber
Microcharmus Lourenço, 1995	Quaternary – Recent
83. <i>Microcharmus henderickxi</i> (Lourenço, 2009a)	Qt Madagascar copal
Microtityus Kjellesvig-Waering, 1966c	Neogene – Recent
84. <i>Microtityus ambarensis</i> (Schawaller, 1982a)	Ne Dominican amber
† Palaeoakentrobuthus Lourenço & Weitschat, 2000	Palaeogene
85. <i>Palaeoakentrobuthus knodeli</i> Lourenço & Weitschat, 2000*	Pa Baltic amber
† Palaeoananteris Lourenço & Weitschat, 2001	Palaeogene
86. <i>Palaeoananteris ribnitiodamgartensis</i> Lourenço & Weitschat, 2001*	Pa Baltic amber
87. <i>Palaeoananteris ukrainensis</i> Lourenço & Weitschat, 2009	Pa Rovno amber
88. <i>Palaeoananteris wunderlichi</i> Lourenço, 2004	Pa Baltic amber
† Palaeoisometrus Lourenço & Weitschat, 2005a	Palaeogene
89. <i>Palaeoisometrus elegans</i> Lourenço & Weitschat, 2005a*	Pa Baltic amber
† Palaeogroosphus Lourenço, 2000a	Quaternary
90. <i>Palaeogroosphus copalensis</i> (Lourenço, 1996b)	Qt Copal
91. <i>Palaeogroosphus jacquesi</i> Lourenço & Henderickx, 2002	Qt Copal
† Palaeolychas Lourenço & Weitschat, 1996	Palaeogene
92. <i>Palaeolychas balticus</i> Lourenço & Weitschat, 1996*	Pa Baltic amber
93. <i>Palaeolychas weitschati</i> Lourenço, 2012	Pa Baltic amber
† Palaeoprotobuthus Lourenço & Weitschat, 2000	Palaeogene
94. <i>Palaeoprotobuthus pusillus</i> Lourenço & Weitschat, 2000*	Pa Baltic amber
† Palaeospinobuthus Lourenço, Henderickx & Weitschat, 2005	Palaeogene
95. <i>Palaeospinobuthus cenozoicus</i> Lourenço, Henderickx & Weitschat, 2005*	Pa Baltic amber
† Palaeotityobuthus Lourenço & Weitschat, 2000	Palaeogene
96. <i>Palaeotityobuthus longiaculeus</i> Lourenço & Weitschat, 2000*	Pa Baltic amber
Tityus C. L. Koch, 1836	?Palaeogene – Recent
97. <i>Tityus azari</i> Lourenço, 2013	Ne Dominican amber
98. 'Tityus' eogenus Menge, 1869 [presumably misplaced]	Pa Baltic amber

99. *Tityus geratus* Santiago-Blay & Poinar, 1988 Ne Dominican amber
 100. *Tityus (Brazilotityus) hartkorni* Lourenço, 2009b Ne Dominican amber
 101. *Tityus (Brazilotityus) knodeli* Lourenço, 2014 Ne Chiapas amber
 † ***Uintascorpio* Perry, 1995** **Palaeogene**
 102. *Uintascorpio halandrasorum* Perry, 1995* Pa Green River

BUTHIDAE incertae sedis

103. ‘*Scorpio*’ *schweiggeri* Holl, 1829 Qt Copal [not amber!]

BOTHRIURIDAE Simon, 1880 **Recent**

= TELEONIDAE Peters, 1861 [based on a generic homonym]
 = ACANTHOCHIROIDAE Karsch, 1880b

no fossil record

CHACTOIDEA Pocock, 1893 **Cretaceous – Recent**

- † **PALAOEUSCORPIDAE Lourenço, 2003** **Cretaceous**

- † ***Palaeoeuscorpius* Lourenço, 2003** **Cretaceous**

104. *Palaeoeuscorpius gallicus* Lourenço, 2003* K French amber

CHACTIDAE Pocock, 1893 **Cretaceous – Recent**

= BROTEIDAE Simon, 1879a [supressed for lack of usage]

- † ***Araripescorpius* Campos, 1986** **Cretaceous**

105. *Araripescorpius ligabuei* Campos, 1986* K Crato Formation

Chactas Gervais, 1844 **Subrecent – Recent**

106. *Chactas pleistocenicus* Lourenço & Weitschat, 2005b Qt Colombian copal

AKRAVIDAE Levy, 2007 **Recent**

no fossil record

CHAERILIDAE Pocock, 1893 **Cretaceous – Recent**

- Electrochaerilus* Santiago-Blay et al., 2004** **Cretaceous**

107. *Electrochaerilus buckleyi* Santiago-Blay et al., 2004 K Myanmar amber

DIPLOCENTRIDAE Karsch, 1880b **Recent**

no fossil record

EUSCORPIIIDAE Laurie, 1896 **Recent**

no fossil record

HETEROSCORPIONIDAE Kraepelin, 1905 **Recent**

no fossil record

HEMISCORPIIIDAE Pocock, 1893 **Cretaceous – Recent**

= ISCHNURIDAE Simon, 1879a

= LIOCHELIDAE Fet & Bechly, 2001	
= † PROTOISCHNURIDAE Carvalho & Lourenço, 2001	
† <i>Protoischnurus</i> Carvalho & Lourenço, 2001	Cretaceous
108. <i>Protoischnurus axelrodorum</i> Carvalho & Lourenço, 2001*	K Crato Formation
IURIDAE Thorell, 1876b	Recent
no fossil record	
SCORPIONIDAE Latreille, 1802	Neogene – Recent
= PANDINOIDAE Thorell, 1876b	
= HETEROMETRIDAE Simon, 1879a	
† <i>Mioscorpio</i> Kjellesvig-Waering, 1986	Neogene
109. <i>Mioscorpio zeuneri</i> (Hadži, 1931)*	Ne Swabian Alps
† <i>Sinoscorpius</i> Hong, 1983a	Neogene
110. <i>Sinoscorpius shandongensis</i> Hong, 1983a*	Ne Shandong, China
SUPERSTITIONIIDAE Stahnke, 1940	Recent
no fossil record	
TROGLOTAYOSICIDAE Lourenço, 1998	Recent
no fossil record	
VAEJOVIDAE Thorell, 1876b	Recent
no fossil record	
SCORPIONES <i>incertae sedis</i>	
Scorpiones <i>incertae sedis</i> in Dunlop & Selden (2013)	S Trecastle, Wales
† <i>Brontoscorpio</i> Kjellesvig-Waering, 1972	Devonian
111. <i>Brontoscorpio anglicus</i> Kjellesvig-Waering, 1972	D England
† <i>Gondwanascorpio</i> Gess, 2013	Devonian
112. <i>Gondwanascorpio emzantsiensis</i> Gess, 2013*	D Grahamstown
† <i>Gymnoscorpius</i> Jeram, 1994b	Carboniferous
113. <i>Gymnoscorpius mutillidigitatus</i> Jeram, 1994b*	C northern England
† <i>Hubeiscorpio</i> Walossek, Li & Brauckmann, 1990	Devonian
114. <i>Hubeiscorpio gracilitarsis</i> Walossek, Li & Brauckmann, 1990*	D Hubei, China
† <i>Liassoscorionides</i> Bode, 1951	Jurassic
115. <i>Liassoscorionides schmidti</i> Bode, 1951*	J Hondelage, Germany
† <i>Palaeomachus</i> Pocock, 1911	Carboniferous
116. <i>Palaeomachus anglicus</i> (Woodward, 1876)*	C Mansfield
† <i>Titanoscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
117. <i>Titanoscorpio douglassi</i> Kjellesvig-Waering, 1986	C Mazon Creek
† <i>Wattisonia</i> Wills, 1960	Carboniferous

118. *Wattisonia coseleyensis* Wills, 1960 C Coseley

MISIDENTIFICATIONS

1. ?*Waterstonia brachistodactyla* Kjellesvig-Waering, 1986 [plant fragment?] C Beith, Ayrshire
2. ?*Mesophonus maculatus* (Brauer, Redtenbacher & Ganglbauer, 1889)
[?insect: cockroach] J Siberia
3. *Tiphoscorpio hueberi* Kjellesvig-Waering, 1986 [myriapod: *Eoarthroleura*] D New York

c. 2,000 Recent species

OPILIONES

38 currently valid species of fossil harvestman

OPILIONES Sundevall, 1833	Devonian – Recent
CYPHOPHTHALMI Simon, 1879a (suborder)	Cretaceous – Recent
NEOGOVEIDAE Shear, 1980	Recent
no fossil record	
OGOVEIDAE Shear, 1980	Recent
no fossil record	
PETTALIDAE Shear, 1980	Recent
no fossil record	
SIRONIDAE Simon, 1879a	Palaeogene – Recent
Siro Latreille, 1796	Palaeogene – Recent
1. <i>Siro balticus</i> Dunlop & Mitov, 2011	Pa Baltic amber
2. <i>Siro platypedibus</i> Dunlop & Giribet, 2003	Pa Bitterfeld amber
STYLOCELLIDAE Hansen & Sørensen, 1904	Cretaceous – Recent
† <i>Palaeosiro</i> Poinar, 2008	Cretaceous – Recent
3. <i>Palaeosiro burmanicum</i> Poinar, 2008	K Myanmar amber
NB: Originally described as a sironid, but regarded as a stylocellid by Giribet <i>et al.</i> (2012)	
TROGLOSIRONIDAE Shear, 1993	Recent
no fossil record	
TETROPHTHALMI Garwood, Sharma, Dunlop & Giribet, 2014	
(suborder)	Devonian – Carbon.
† <i>Eophalangium</i> Dunlop, Anderson, Kerp & Hass, 2004	Devonian
4. <i>Eophalangium sheari</i> Dunlop, Anderson, Kerp & Hass, 2004*	D Rhynie chert
† <i>Hastocularis</i> Garwood, Sharma, Dunlop & Giribet, 2014	Devonian
5. <i>Hastocularis argus</i> Garwood, Sharma, Dunlop & Giribet, 2014*	D Montceau-les-Mines
EUPNOI Hansen & Sørensen, 1904 (suborder)	Devonian – Recent
plesion taxa	
† <i>Brigantibunum</i> Dunlop & Anderson, 2005	Carboniferous
6. <i>Brigantibunum listoni</i> Dunlop & Anderson, 2005*	C East Kirkton

† <i>Kustarachne</i> Scudder, 1890b	Carboniferous
7. <i>Kustarachne tenuipes</i> Scudder, 1890b*	C Mazon Creek
i. = <i>Kustarachne exstincta</i> Melander, 1903	C Mazon Creek
ii. = <i>Kustarachne conica</i> Petrunkevitch, 1913	C Mazon Creek
† <i>Macrogyion</i> Garwood et al., 2011	Carboniferous
8. <i>Macrogyion cronus</i> Garwood et al. 2011*	C Montceau-les-Mines
CADDOIDEA Banks, 1893	Palaeogene – Recent
CADDIDAE Banks, 1893	Palaeogene – Recent
Caddo Banks, 1892a	Palaeogene – Recent
9. <i>Caddo dentipalpus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic / Bitter. amber
PHALANGIOIDEA Latreille, 1802	Palaeogene – Recent
family uncertain	
† <i>Petrunkewitchiana</i> Mello-Leitão, 1937 [genus <i>incertae sedis</i>]	Palaeogene
10. <i>Petrunkewitchiana oculata</i> (Petrunkewitch, 1922)*	Pa Florissant
MONOSCUTIDAE Forster, 1948	Recent
no fossil record	
NEOPILIONIDAE Lawrence, 1931	Recent
no fossil record	
PHALANGIIDAE Latreille, 1802	Palaeogene – Recent
Amilenus Martens, 1969	Palaeogene – Recent
11. <i>Amilenus deltshevi</i> Dunlop & Mitov, 2009	Pa Bitterfeld amber
Dicranopalpus Doleschall, 1852	Palaeogene – Recent
12. <i>Dicranopalpus ramiger</i> (C. L. Koch & Berendt, 1854)	Pa Baltic / Bitter. amber
i. = <i>Opilio corniger</i> Menge, 1854	Pa Baltic amber
ii. = <i>Dicranopalpus palmnickensis</i> Roewer, 1939	Pa Baltic amber
† <i>Lacinius</i> Thorell, 1876	Palaeogene – Recent
13. ? <i>Lacinius erinaceus</i> Staręga, 1966 [Recent]	Pa Bitterfeld amber
† <i>Stephanobunus</i> Dunlop & Mammitzsch, 2010	Palaeogene
14. <i>Stephanobunus mitovi</i> Dunlop & Mammitzsch, 2010*	Pa Baltic amber
?Phalangiidae	
15. <i>Opilio ovalis</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
[probably misplaced at genus level]	
SCLEROSOMATIDAE Simon, 1879a	Jurassic – Recent
† <i>Amauropilio</i> Mello-Leitão, 1937	Palaeogene
16. <i>Amauropilio atavus</i> (Cockerell, 1907)	Pa Florissant

17. *Amauropilio lacoei* (Petrunkewitch, 1922) Pa Florissant
- Leiobunum* C. L. Koch, 1839a** **Jurassic – Recent**
18. *Leiobunum longipes* Menge, 1854 Pa Baltic /Bitter. amber
- i. = *Leiobunum saparum* Menge, 1854 [?lapsus] Pa Baltic amber
- ii. = *Leiobunum inclusum* Roewer, 1939 Pa Baltic amber
- † ***Mesobunus* Huang, Selden & Dunlop, 2009** **Jurassic**
19. *Mesobunus dunlopi* Giribet, Tourhino, Shih & Ren, 2012 J Daohugou
20. *Mesobunus martensi* Huang, Selden & Dunlop, 2009* J Daohugou
- Family uncertain**
- † ***Daohugopilio* Huang, Selden & Dunlop, 2009** **Jurassic**
21. *Daohugopilio sheari* Huang, Selden & Dunlop, 2009* J Daohugou
- DYSPNOI Hansen & Sørensen, 1904 (suborder)** **Carbon. – Recent**
- family uncertain
- † ***Ameticos* Garwood et al., 2011** **Carboniferous**
22. *Ameticos scolos* Garwood et al. 2011* C Montceau-les-Mines
- † ***Echinopustulatus* Dunlop, 2004** **Carboniferous**
23. *Echinopustulatus samuelnelsoni* Dunlop, 2004* C Missouri
- ISCHYROPSALIDOIDEA Simon, 1879a** **Palaeogene – Recent**
- Tentative assignment, family uncertain
- † ***Piankhi* Dunlop, Bartel & Mitov, 2012** **Palaeogene**
24. *Piankhi steineri* Dunlop, Bartel & Mitov, 2012* Pa Baltic amber
- CERATOLASMATIDAE Shear, 1986** **Recent**
- no fossil record
- ISCHYROPSALIDIDAE Simon, 1879a** **Recent**
- no fossil record
- SABAONIDAE Dresco, 1970** **Palaeogene – Recent**
- Sabacon* Simon, 1879a** **Palaeogene – Recent**
25. *Sabacon claviger* (Menge, 1854) Pa Baltic amber
- i. = *Sabacon bachofeni* Roewer, 1939 Pa Baltic amber
- TROGULOIDEA Sundevall, 1833** **Cretaceous – Recent**
- [family uncertain; Shear (2010) suggested it is not an ortholasmatine, but may represent a new family]
- † ***Halitherses* Giribet & Dunlop, 2005** **Cretaceous**
26. *Halitherses grimaldii* Giribet & Dunlop, 2005* K Myanmar amber
- DICRANOLASMATIDAE Simon, 1879a** **Recent**
- no fossil record

† EOTROGULIDAE Petrunkevitch, 1955a	Carboniferous
† <i>Eotrogulus</i> Thevenin, 1901	Carboniferous
27. <i>Eotrogulus fayoli</i> Thevenin, 1901*	C Commentry
 NEMASTOMATIDAE Simon, 1879a	Palaeogene – Recent
<i>Histicostoma</i> Kratochvíl, 1958	Palaeogene – Recent
28. ? <i>Histicostoma tuberculatum</i> (C. L. Koch & Berendt, 1854)	Pa Baltic/Bitter. amber
<i>Mitostoma</i> Roewer, 1951	Palaeogene – Recent
29. ? <i>Mitostoma denticulatum</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
i. = <i>Nemastoma succineum</i> Roewer, 1939	Pa Baltic amber
30. ? <i>Mitostoma gruberi</i> Dunlop & Mitov, 2009	Pa Bitterfeld amber
<i>Nemastoma</i> C. L. Koch, 1836	Palaeogene – Recent
31. ? <i>Nemastoma incertum</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
 † NEMASTOMOIDIDAE Petrunkevitch, 1955a	Carboniferous
† <i>Nemastomoides</i> Thevenin, 1901	Carboniferous
= † <i>Protopilio</i> Petrunkevitch, 1913	
32. <i>Nemastomoides elaveris</i> Thevenin, 1901*	C Commentry
33. <i>Nemastomoides longipes</i> (Petrunkevitch, 1913)	C Mazon Creek
 NIPPONOSALIDIDAE Martens, 1976	Recent
no fossil record	
 TROGULIDAE Sundevall, 1833	Palaeogene – Recent
<i>Trogulus</i> Latreille, 1802	Palaeogene – Recent
34. <i>Trogulus longipes</i> Haupt, 1956	Pa Geiseltal
 LANIATORES Thorell, 1876c (suborder)	Palaeogene – Recent
family uncertain	
<i>Philacarus</i> Sørensen, 1932	Neogene – Recent
35. <i>Philacarus hispaniolensis</i> Cokendolpher & Poinar, 1992	Ne Dominican amber
 INSIDIATORES Loman, 1900 (infraorder)	Palaeogene – Recent
TRAVUNIOIDEA Absolon & Kratochvíl, 1932	Palaeogene – Recent
CLADONYCHIDAE Hadži, 1935	Palaeogene – Recent
† <i>Proholoscotolemon</i> Ubick & Dunlop, 2005	Palaeogene
36. <i>Proholoscotolemon nemastomoides</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
? <i>Proholoscotolemon</i> sp. in Ubick & Dunlop (2005)	Pa Baltic amber
 PENTANYCHIDAE Briggs, 1971	Recent
no fossil record	

TRAVUNIIDAE Absolon & Kratochvíl, 1932	Recent
no fossil record	
TRIAENONYCHOIDEA Sørensen, 1886	Recent
SYNTHETONYCHIIDAE Forster, 1954	Recent
no fossil record	
TRIAENONYCHIDAE Sørensen, 1886	Recent
no fossil record	
GRASSATORES Kury, 2002 (infraorder)	Neogene – Recent
SAMOIDEA Sørensen, 1886	Neogene – Recent
BIANTIDAE Thorell, 1889	Recent
no fossil record	
ESCADABIIDAE Kury & Pérez González <i>in</i> Kury, 2003	Recent
no fossil record	
KIMULIDAE Pérez González, Kury & Alonso-Zarazaga <i>in</i> Pérez González & Kury, 2007	Neogene – Recent
<i>Kimula</i> Goodnight & Goodnight, 1942	Neogene – Recent
<i>Kimula</i> sp. <i>in</i> Cokendolpher & Poinar (1992)	Ne Dominican amber
PODOCTIDAE Roewer, 1912	Recent
no fossil record	
SAMOIDAE Sørensen, 1886	Neogene – Recent
<i>Hummelinckiolus</i> Šilhavý, 1979	Neogene – Recent
37. <i>Hummelinckiolus silhavyi</i> Cokendolpher & Poinar, 1998	Ne Dominican amber
Pellobunus Banks, 1905	Neogene – Recent
38. <i>Pellobunus proavus</i> Cokendolpher, 1987	Ne Dominican amber
STYGNOMMATIDAE Roewer, 1923	Recent
no fossil record	
ASSAMIOIDEA Sørensen, 1884	Recent
ASSAMIIDAE Sørensen, 1884	Recent
no fossil record	
EPEDANIDAE Sørensen, 1886	Recent
no fossil record	

PETROBUNIDAE Sharma & Giribet, 2011	Recent
no fossil record	
PYRAMIDOPHIIDAE Sharma, Prieto & Giribet, 2011	Recent
no fossil record	
STYGNOPSIDAE Sørensen, 1932	Recent
no fossil record	
TITHAEIDAE Sharma & Giribet, 2011	Recent
no fossil record	
GONYLEPTOIDEA Sundevall, 1833	Recent
AGORISTENIDAE Šilhavý, 1973	Recent
no fossil record	
COSMETIDAE C. L. Koch, 1839a	Recent
no fossil record	
CRANAIDAE Roewer, 1913	Recent
no fossil record	
GONYLEPTIDAE Sundevall, 1833	Recent
no fossil record	
MANAOSSIIDAE Roewer, 1943	Recent
no fossil record	
STYGNIDAE Simon, 1879b	Recent
no fossil record	
PHALANGODOIDEA Simon, 1879a	Recent
ONCOPODIDAE Thorell, 1876c	Recent
no fossil record	
PHALANGODIDAE Simon, 1879a	Recent
no fossil record	
ZALMOXOIDEA Sørensen, 1886	Recent
FISSIPHALLIIDAE Martens, 1988	Recent
no fossil record	
GUASINIIDAE González-Sponga, 1997	Recent
no fossil record	

ICALEPTIDAE Kury & Pérez González, 2002 Recent

no fossil record

ZALMOXIDAE Sørensen, 1886 Recent

no fossil record

OPILIONES *incertae sedis*

unnamed specimen *in* Jell & Duncan (1986) K Koonwarra

NOMINA DUBIA

1. *Cheiromachus coriaceus* Menge, 1854 Pa Baltic amber
2. *Phalangium succineum* Presl, 1822 Pa Baltic amber

MISIDENTIFICATIONS

1. *Hasseltides primigenius* Weyenbergh, 1869 [crinoid] J Solnhofen
2. *Phalangites multipes* Münster *in* Roth, 1851 [crustacean] J Solnhofen
3. *Phalangites priscus* Münster, 1839 [crustacean] J Solnhofen
4. *Rhabdotarachnoides simoni* Haupt, 1957 [plant fragment] P Rotliegend

6,491 Recent species according to Kury (2011)

PHALANGIOTARBIDA

31 currently valid species of fossil phalangiotarbid

† PHALANGIOTARBIDA Haase, 1890	Devonian – Permian
= † ARCHITARBIDA Petrunkevitch, 1945a	
† DEVONOTARBIDAe Poschmann & Dunlop, 2012	Devonian
† Devonotarbus Poschmann, Anderson & Dunlop, 2005	Devonian
1. <i>Devonotarbus hombachensis</i> Poschmann, Anderson & Dunlop, 2005* D Germany	
† ANTHRACOTARBIDAe Kjellesvig-Waering, 1969	Carboniferous
† Anthracotarbus Kjellesvig-Waering, 1969	Carboniferous
2. <i>Anthracotarbus hintoni</i> Kjellesvig-Waering, 1969* C Oklahoma	
† ARCHITARBIDAe Karsch, 1882	Carboniferous
= † PHALANGIOTARBIDAe Haase, 1890	
† Architarbus Scudder, 1868	Carboniferous
3. <i>Architarbus hoffmanni</i> Guthörl, 1934 C Saar basin	
i. = <i>Opiliotarbus klicheri</i> Waterlot, 1935 C Saar basin	
ii. = <i>Goniatarbus sarana</i> Guthörl, 1965 C Saar basin	
4. <i>Architarbus minor</i> Petrunkevitch, 1913 C Mazon Creek	
5. <i>Architarbus rotundatus</i> Scudder, 1868* C Mazon Creek	
† Bornatarbus Rößler & Schneider, 1997	Carboniferous
6. <i>Bornatarbus mayasii</i> (Haupt in Nindel, 1955)* C Germany / UK	
† Discotarbus Petrunkevitch, 1913	Carboniferous
7. <i>Discotarbus deplanatus</i> Petrunkevitch, 1913* C Mazon Creek	
† Geratarbus Scudder, 1890b	Carboniferous
8. <i>Geratarbus lacoei</i> Scudder, 1890b* C Mazon Creek	
9. <i>Geratarbus bohemicus</i> Petrunkevitch, 1953 C Nýřany	
† Goniatarbus Petrunkevitch, 1949	Carboniferous
10. <i>Goniatarbus angulatus</i> (Pocock, 1911) C Coseley	
11. <i>Goniatarbus tuberculatus</i> (Pocock, 1911)* C Coseley	
i. = <i>Goniatarbus tuberculatus</i> Petrunkevitch, 1949 C Coseley	
† Hadrachne Melander, 1903	Carboniferous
12. <i>Hadrachne horribilis</i> Melander, 1903* C Mazon Creek	
† Leptotarbus Petrunkevitch, 1945a	Carboniferous
13. <i>Leptotarbus torpedo</i> (Pocock, 1911)* C Coseley	
† Mesotarbus Petrunkevitch, 1949	Carboniferous
14. <i>Mesotarbus angustus</i> (Pocock, 1911) C Coseley	

15. *Mesotarbus eggintoni* (Pocock, 1911) C Coseley
16. *Mesotarbus hindi* (Pocock, 1911) C Coseley
17. *Mesotarbus intermedius* Petrunkevitch, 1949* C Coseley
18. *Mesotarbus peteri* Dunlop & Horrocks, 1997 C Westhoughton
- † ***Metatarbus* Petrunkevitch, 1913** **Carboniferous**
19. *Metatarbus triangularis* Petrunkevitch, 1913* C Mazon Creek
- † ***Otarbus* Petrunkevitch, 1945a** **Carboniferous**
20. *Otarbus pulcher* Petrunkevitch, 1945a* C Mazon Creek
21. *Otarbus ovatus* Petrunkevitch, 1945a C Mazon Creek
- † ***Orthotarbus* Petrunkevitch, 1945a** **Carboniferous**
22. *Orthotarbus longipes* Simon, 1971 C Halleschen Mulde
23. *Orthotarbus minutus* (Petrunkevitch, 1913)* C Mazon Creek
24. *Orthotarbus robustus* Petrunkevitch, 1945a C Mazon Creek
25. *Orthotarbus nyranensis* Petrunkevitch, 1953 C Nýřany
- † ***Paratarbus* Petrunkevitch, 1945a** **Carboniferous**
26. *Paratarbus carbonarius* Petrunkevitch, 1945a* C Mazon Creek
- † ***Phalangiotarbus* Haase, 1890** **Carboniferous**
27. *Phalangiotarbus subovalis* (Woodward, 1872b)* C Burnley
- † ***Pycnotarbus* Darber, 1990** **Carboniferous**
28. *Pycnotarbus verrucosus* Darber, 1990* C Oelsnitz
- † ***Triangulotarbus* Patrick, 1989** **Carboniferous**
29. *Triangulotarbus terrehautesis* Patrick, 1989* C Indiana
- † **HETEROTARBIDAE Petrunkevitch, 1913** **Carboniferous**
- † ***Heterotarbus* Petrunkevitch, 1913** **Carboniferous**
30. *Heterotarbus ovatus* Petrunkevitch, 1913* C Mazon Creek
- † **OPILIOTARBIDAE Petrunkevitch, 1945a** **Carb. – Permian**
- † ***Opiliotarbus* Pocock, 1910** **Carb. – Permian**
31. *Opiliotarbus elongatus* (Scudder, 1890b)* C – P USA / Germany

NOMINA DUBIA

1. *Eotarbus litoralis* Kušta, 1888 C Rakovník
2. *Nemastomoides depressus* Petrunkevitch, 1913 C Mazon Creek

no Recent species

PSEUDOSCORPIONES

45 currently valid species of fossil pseudoscorpion

PSEUDOSCORPIONES De Geer, 1778	Devonian – Recent
= CHERNETES Simon, 1879a		
† DRACOCHELIDAE Schawaller, Shear & Bonamo, 1991 (plesion family)	Devonian
† <i>Dracochela</i> Schawaller, Shear & Bonamo, 1991	Devonian
1. <i>Dracochela deprehendor</i> Schawaller, Shear & Bonamo, 1991*	D Gilboa
CHELONETHI Thorell, 1882	Cretaceous – Recent
EPIOCHIERATA Harvey, 1992	Cretaceous – Recent
CTHONOIDEA Daday, 1888	Palaeogene – Recent
CTHONIIDAE Daday, 1888	Palaeogene – Recent
<i>Chthonius</i> C. L. Koch, 1843a	Palaeogene – Recent
2. <i>Chthonius (Chthonius) mengei</i> Beier, 1937	Pa Baltic amber
3. <i>Chthonius (Chthonius) pristinus</i> Schawaller, 1978	Pa Baltic amber
<i>Pseudochthonius</i> Balzan, 1892	Neogene – Recent
4. <i>Pseudochthonius squamosus</i> Schawaller, 1980a	Ne Dominican amber
<i>Tyrannchthonius</i> Chamberlin, 1929	Quaternary – Recent
<i>Tyrannchthonius</i> sp. in Judson (2010)	Qt Madagascan copal
LECHYTIDAE Chamberlin, 1929	Neogene – Recent
<i>Lechyta</i> Balzan, 1892	Neogene – Recent
5. <i>Lechyta tertaria</i> Schawaller, 1980a	Ne Dominican amber
TRIDENCHTHONIIDAE Balzan, 1892	Palaeogene – Recent
= DITHIDAE Chamberlin, 1929		
† <i>Chelignathus</i> Menge, 1854	Palaeogene
6. <i>Chelignathus kochii</i> Menge, 1854*	Pa Baltic amber
FEAELLOIDEA Ellingsen, 1906	Palaeogene – Recent
FEAELLIDAE Ellingsen, 1906	Recent
† <i>Feaella (Tetrafeaella)</i> Beier, 1955	Palaeogene – Recent
7. <i>Feaella (Tetrafeaella) groehni</i> Henderickx in Henderickx & Boone, 2014	Pa	Baltic amber
PSEUDOGARYPIDAE Chamberlin, 1923a	Palaeogene – Recent
<i>Pseudogarypus</i> Ellingsen, 1909	Palaeogene – Recent

8. <i>Pseudogarypus extensus</i> Beier, 1937	Pa	Baltic amber
9. <i>Pseudogarypus hemprichii</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
10. <i>Pseudogarypus minor</i> Beier, 1947a	Pa	Baltic/Rovno amber
11. <i>Pseudogarypus pangaea</i> Henderickx <i>in</i> Henderickx <i>et al.</i> , 2006	Pa	Baltic amber
12. <i>Pseudogarypus synchrotron</i> Henderickx <i>in</i> Henderickx <i>et al.</i> , 2012	Pa	Baltic amber
IOCHIERATA Harvey, 1992		Cretaceous – Recent
HEMICTENATA Balzan, 1892		Cretaceous – Recent
NEOBISIOIDEA Chamberlin, 1930		Cretaceous – Recent
BOCHICIDAE Chamberlin, 1930		Recent
= VACHONIIDAE Chamberlin, 1947		
no fossil record		
GYMNOBISIIDAE Beier, 1947b		Recent
no fossil record		
HYIDAE Chamberlin, 1930		Recent
no fossil record		
IDEORONCIDAE Chamberlin, 1930		Recent
no fossil record		
NEOBISIIDAE Chamberlin, 1930		Cretaceous – Recent
= OBISIIDAE Sundevall, 1833		
† Electrobisium Cockerell, 1917		Cretaceous
13. <i>Electrobisium acutum</i> Cockerell, 1917a*	K	Myanmar amber
Microcreagris Balzan, 1892		Palaeogene – Recent
14. <i>Microcreagris koellnerorum</i> Schawaller, 1978	Pa	Baltic amber
Neobisium Chamberlin, 1930		Palaeogene – Recent
15. <i>Neobisium (Neobisium) exstinctum</i> Beier, 1955	Pa	Baltic amber
16. <i>Neobisium henderickxi</i> Judson, 2003	Pa	Baltic amber
Roncus L. Koch, 1873		Palaeogene – Recent
17. <i>Roncus succineus</i> Beier, 1955	Pa	Baltic amber
PARAHYIDAE Harvey, 1992		Recent
no fossil record		
SYARINIDAE Chamberlin, 1930		Recent
no fossil record		
PANCTENATA Balzan, 1892		Cretaceous – Recent
GARYPOIDEA Simon, 1879a		Cretaceous – Recent
GARYPIDAE Simon, 1879a		Recent

= SYNSPHRONIDAE Beier, 1932a	
no fossil record	
GARYPINIDAE Daday, 1888	Cretaceous – Recent
Amblyolpium Simon, 1898b	Cretaceous – Recent
18. <i>Amblyolpium burmiticum</i> (Cockerell, 1920)	K Myanmar amber
Garypinus Daday, 1888	Palaeogene – Recent
19. <i>Garypinus electri</i> Beier, 1937	Pa Baltic amber
GEOGARYPIDAE Chamberlin, 1930	Palaeogene – Recent
Geogarypus Chamberlin, 1930	Palaeogene – Recent
20. <i>Geogarypus gorskii</i> Henderickx, 2005	Pa Baltic/Rovno amber
21. <i>Geogarypus macrodactylus</i> Beier, 1937	Pa Baltic amber
22. <i>Geogarypus major</i> Beier, 1937	Pa Baltic amber
LARCIDAE Harvey, 1992	Recent
no fossil record	
MENTHIDAE Chamberlin, 1930	Recent
no fossil record	
OLPIIDAE Banks, 1895	Palaeogene – Recent
no fossil record	
STERNOPHOROIDEA Chamberlin, 1923b	Neogene – Recent
STERNOPHORIDAE Chamberlin, 1923b	Neogene – Recent
Idiogaryops Hoff, 1963	Neogene – Recent
23. <i>Idiogaryops pumilus</i> (Hoff, 1963) [Recent]	Ne–R Dominican amber
CHEIRIDIOIDEA Hansen, 1894	Palaeogene – Recent
CHEIRIDIIDAE Hansen, 1894	Palaeogene – Recent
Cheiridium Menge, 1855	Palaeogene – Recent
24. <i>Cheiridium hartmanni</i> (Menge, 1854)	Pa Baltic amber
Cryptocheiridium Chamberlin, 1931a	Neogene – Recent
25. <i>Cryptocheiridium (Cryptocheiridium) antiquum</i> Schawaller, 1981	Ne Dominican amber
PSEUDOCHIRIDIIDAE Chamberlin, 1923b	Neogene – Recent
Pseudochiridium With, 1906	Neogene – Recent
26. <i>Pseudochiridium lindae</i> Judson, 2007	Ne Dominican amber
CHELIFEROIDEA Risso, 1826	Cretaceous – Recent
ATEMNIDAE Kishida, 1929	Palaeogene – Recent
Atemninae indet. <i>in</i> Judson (2010)	Qt Dominican amber

Paratemnoides Harvey, 1991	Quaternary – Recent
27. <i>Paratemnoides nidificator</i> (Balzan, 1888) [Recent]	Qt–R Colombian copal
† Progonatemnus Beier, 1955	Palaeogene
28. <i>Progonatemnus succineus</i> Beier, 1955*	Pa Baltic amber
 CHELIFERIDAE Riso, 1826	 Cretaceous – Recent
Cheliferidae? indet. <i>in</i> Judson (2009)	K Archingeay amber
† Dicella Menge, 1854	Palaeogene
= † <i>Oligochelifer</i> Beier, 1937	
29. <i>Dicella berendtii</i> Menge, 1954*	Pa Baltic amber
30. <i>Dicella gracilis</i> (Beier, 1937)	Pa Baltic amber
31. <i>Dicella granulatus</i> (Beier, 1937)	Pa Baltic amber
32. <i>Dicella serratidentatus</i> (Beier, 1937)	Pa Baltic amber
† Electrochelifer Beier, 1937	Palaeogene
33. <i>Electrochelifer bachofeni</i> Beier, 1947a	Pa Baltic amber
34. <i>Electrochelifer balticus</i> Beier, 1955	Pa Baltic amber
35. <i>Electrochelifer mengei</i> Beier, 1937*	Pa Baltic amber
36. <i>Electrochelifer rapulitarsatus</i> Beier, 1947a	Pa Baltic amber
† Heurtaultia Judson, 2009 [tentative referral to family]	Cretaceous
37. <i>Heurtaultia rossiorum</i> Judson, 2009	K Archingeay amber
† Pycnochelifer Beier, 1937	Palaeogene
38. <i>Pycnochelifer kleemannii</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
i. = <i>Obisium rathkii</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† Trachychelifer Hong, 1983b	Palaeogene
39. <i>Trachychelifer liaoningense</i> Hong, 1983b*	Pa Chinese amber
 CHERNETIDAE Menge, 1855	 Cretaceous – Recent
Chernetidae gen. et sp. indet <i>in</i> Schawaller (1991)	K Canadian amber
Chernetidae gen. et sp. indet <i>in</i> Schawaller (1982b)	Ne Chiapas amber
† Oligochernes Beier, 1937	Palaeogene
40. <i>Oligochernes bachofeni</i> Beier, 1937	Pa Baltic amber
41. <i>Oligochernes wigandi</i> (Menge, 1854)	Pa Baltic amber
Pachychernes Beier, 1932b	Neogene – Recent
42. <i>Pachychernes effossus</i> Schawaller, 1980b	Ne Dominican amber
43. <i>Pachychernes aff. subrobustus</i> (Balzan, 1892) [Recent]	Qt–R Colombian copal
 WITHIIDAE Chamberlin, 1931b	 Palaeogene – Recent
† Beierowithius Mahnert, 1979	Palaeogene
44. <i>Beierowithius sieboldtii</i> (Menge, 1854)*	Pa Baltic amber
Withius Kew, 1911	Quaternary – Recent
45. <i>Chelifer eucarpus</i> Dalman, 1826	Qt East African opal

NOMINA DUBIA

1. *Chelifer ehrenbergii* C. L. Koch & Berendt, 1854 Pa Baltic amber

NOMINA NUDA

1. *Chelifer fossilis* Weyenbergh, 1874 J Solnhofen

3,385 Recent species according to Harvey (2009)

SOLIFUGAE

5 currently valid species of camel spider

- *Schneidarachne* appears to show some solifuge-like features and was tentatively assigned to the stem-lineage of this order; for convenience it is listed here alongside the camel spiders
- a family name *Protosolpugidae* has been proposed for *Protosolpuga*, but was not recognised in most of the subsequent literature – cf. Selden & Shear's (1996) revision

stem-lineage?

† *Schneidarachne* Dunlop & Rössler, 2003 Carboniferous
 1. *Schneidarachne saganii* Dunlop & Rössler, 2003* C Kamienna Góra

SOLIFUGAE Sundevall, 1833 Carbon. – Recent

† *Protosolpuga* Petrunkevitch, 1913 Carboniferous
 2. *Protosolpuga carbonaria* Petrunkevitch, 1913* C Mazon Creek

AMMOTRECHIDAE Roewer, 1934 Neogene – Recent

† *Happlodontus* Poinar & Santiago-Blay, 1989 Neogene
 3. *Happlodontus proterus* Poinar & Santiago-Blay, 1989* Ne Dominican amber

CEROMIDAE Roewer, 1933 Cretaceous – Recent

† *Cratosolpuga* Selden in Selden & Shear, 1996 Cretaceous
 4. *Cratosolpuga wunderlichi* Selden in Selden & Shear, 1996* K Crato Formation

DAESIIDAE Kraepelin, 1899 Palaeogene – Recent

† *Palaeoblossia* Dunlop, Wunderlich & Poinar, 2004 Palaeogene
 5. *Palaeoblossia groehni* Dunlop, Wunderlich & Poinar, 2004* Pa Baltic amber

EREMOBATIDAE Kraepelin, 1901 Recent

no fossil record

GALEODIDAE Sundevall, 1833 Recent

no fossil record

GYLIPPIDAE Roewer, 1933 Recent

no fossil record

HEXISOPODIDAE Pocock, 1897 Recent

no fossil record

KARSCHIIDAE Kraepelin, 1899 **Recent**

no fossil record

MELANOBLOSSIDAE Roewer, 1933 **Recent**

no fossil record

MUMMUCIIDAE Roewer, 1934 **Recent**

no fossil record

RHAGODIDAE Pocock, 1897 **Recent**

no fossil record

SOLPUGIDAE Leach, 1815 **Recent**

no fossil record

1,075 Recent species according to Harvey (2003)

PALPIGRADI

1 currently valid species of fossil palpigrade

PALPIGRADI Thorell, 1888 Neogene – Recent

= MICROTHELYPHONIDA Grassi & Calandruccio, 1885

family uncertain

† **Paleokoenenia** Rowland & Sissom, 1980 Neogene

1. *Paleokoenenia mordax* Rowland & Sissom, 1980* Ne Onyx Marble

EUKOENENIIDAE Petrunkevitch, 1955a Recent

no fossil record

PROKOENENIIDAE Condé, 1996 Recent

no fossil record

MISIDENTIFICATIONS

1. *Sternarthron zitteli* Haase, 1890 [insect] J Solnhofen

2. *Sternarthron zitteli* var. *minor* (Oppenheim, 1887) [insect] J Solnhofen

78 Recent species according to Harvey (2003)

ACARI: PARASITIFORMES

16 currently valid species of fossil parasitiform mite

- higher systematics and sequence of taxa follows the third edition of *A Manual of Acarology* (Krantz & Walter, eds, 2009), except that their orders are listed here as suborders, and suborders as infraorders to achieve some degree of consistency with other arachnid higher taxa throughout this list

PARASITIFORMES Reuter, 1909 Cretaceous – Recent

= ANACTINOTRICHIDA author, date?

OPILIOACARIDA Zachvatkin, 1952 (suborder) Palaeogene – Recent

= NOTOSTIGMATA author, date?

OPILIOACAROIDEA Vitzthum, 1931 Cretaceous – Recent

OPILIOACARIDAE Vitzthum, 1931 Cretaceous – Recent

= NEOACARIDAE Chamberlin & Mulaik, 1942

Opilioacarus With, 1902 ?Cretaceous – Recent

1. ?*Opilioacarus aenigmus* Dunlop, Sempf & Wunderlich, 2010 Pa Baltic amber
2. ?*Opilioacarus groehni* Dunlop & Bernardi, 2014 K Myanmar amber

Paracarus Chamberlin & Mulaik, 1942 Palaeogene – Recent

3. *Paracarus pristinus* Dunlop, Wunderlich & Poinar, 2004 Pa Baltic amber

HOLOTHYRIDAE Thorell, 1882 (suborder) Recent

= TETRASTIGMATA author, date?

HOLOTYHROIDEA Thorell, 1882 Recent

ALLOTHYRIDAE van der Hammen, 1972 Recent

no fossil record

HOLOTHYRIDAE Thorell, 1882 Recent

no fossil record

NEOTHYRIDAE Lehtinen, 1981 Recent

no fossil record

IXODIDA Leach, 1815 (suborder) Cretaceous – Recent

= METASTIGMATA author, date?

IXODOIDEA Banks, 1907 Cretaceous – Recent

ARGASIDAE Murray, 1877 Cretaceous – Recent

Carios Latreille, 1796 Cretaceous – Recent

4. *Carios jerseyi* Klompen & Grimaldi, 2001 K New Jersey amber

<i>Ornithodoros</i> C. L. Koch, 1844	Neogene – Recent
5. <i>Ornithodoros antiquus</i> Poinar, 1995	Ne Dominican amber
IXODIDAE Banks, 1907	Cretaceous – Recent
Amblyomma C. L. Koch, 1844	Neogene – Recent
6. <i>Amblyomma</i> near <i>argentinae</i> Neumann, 1905 [Recent] (as <i>testudinis</i>) in Lane & Poinar (1986)	Ne–R Dominican amber
7. <i>Amblyomma</i> near <i>dissimile</i> C. L. Koch, 1844 [Recent] in Kierens et al. (1986)	Ne–R Dominican amber
† Compluriscutata Poinar & Buckley, 2008	Cretaceous
8. <i>Compluriscutata</i> <i>vetulum</i> Poinar & Buckley, 2008*	K Myanmar amber
† Cornupalpatum Poinar & Brown, 2003	Cretaceous
9. <i>Cornupalpatum</i> <i>burmanicum</i> Poinar & Brown, 2003*	K Myanmar amber
Dermacentor C. L. Koch, 1844	Neogene – Recent
10. <i>Dermacentor</i> nr. <i>reticulatus</i> (Fabricius, 1794) [Recent] (in Kulczyński in Schille 1916)	Ne–R in a Rhino's ear
Hyalomma C. L. Koch, 1844	Palaeogene – Recent
Hyalomma spp.	Pa Baltic amber
Ixodes Latreille, 1795	Palaeogene – Recent
11. <i>Ixodes</i> <i>sigelos</i> Keirans, Clifford & Corwin, 1976 [Recent]	Qt Argentina
12. <i>Ixodes</i> <i>succineus</i> Weidner, 1964	Pa Baltic amber
NUTALLIELLIDAE Schulze, 1935	Recent
no fossil record	
MESOSTIGMATA G. Canestrini, 1891 (suborder)	Palaeogene – Recent
= GAMASIDA Leach, 1815	
SEJIDA Kramer, 1885 (infraorder)	Palaeogene – Recent
= LIROASPINA author, date?	
= TRICHOPYGIDIINA author, date?	
SEJOIDEA Berlese, 1885	Palaeogene – Recent
ICHTHYOSTOMATOGASTERIDAE Sellnick, 1953	Recent
no fossil record	
SEJIDAE Berlese, 1885	Palaeogene – Recent
= LIROASPIDIDAE Trägårdh, 1946	
Sejus C. L. Koch, 1836 [NB: <i>Seius</i> in an invalid emendation]	Palaeogene – Recent
13. <i>Sejus</i> <i>bdelloides</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
UROPODELLIDAE Camin, 1955	Recent
no fossil record	

TRIGYNASPIDA Camin & Gorirossi, 1955 (infraorder)	Recent
CERCOMEGISTINA Camin & Gorirossi, 1955 (cohort)	Recent
CERCOMEGISTOIDEA Trägårdh, 1937	Recent
ASTERNOSEIIDAE Vale, 1955	Recent
no fossil record	
CERCOMEGISTIDAE Trägårdh, 1937	Recent
no fossil record	
DAVACARIDAE Kethley, 1979	Recent
no fossil record	
PYROSEJIDAE Lindquist & Moraza, 1993	Recent
no fossil record	
SALTISEIIDAE Walter, 2000	Recent
no fossil record	
SEIODIDAE Kethley, 1979	Recent
no fossil record	
ANTENNOPHORINA Berlese, 1882 (cohort)	Recent
ANTENNOPHOROIDEA Berlese, 1892	Recent
ANTENNOPHORIDAE Berlese, 1892	Recent
no fossil record	
CELAENOPSIDEOA Berlese, 1892	Recent
CELAENOPSIDAE Berlese, 1892	Recent
no fossil record	
COSTACARIDAE Hunter, 1993	Recent
no fossil record	
DIPLOGYNIIDAE Trägårdh, 1941	Recent
no fossil record	
EUZERCONIDAE Trägårdh, 1938	Recent
no fossil record	
MEGACELAENOPSIDAE Funck, 1975	Recent
no fossil record	
MEINERTULIDAE Trägårdh, 1950	Recent

no fossil record

NEOTENOGENIIDAE Kethley, 1974 Recent

no fossil record

SCHIZOGENIIDAE Trägårdh, 1950 Recent

no fossil record

TRIPOLOGENIIDAE Funck, 1977 Recent

no fossil record

PARAMEGISTOIDEA Trägårdh, 1946 Recent

PARAMEGISTIDAE Trägårdh, 1946 Recent

no fossil record

FEDRIZZIOIDEA Trägårdh, 1937 Recent

FEDRIZZIIDAE Trägårdh, 1937 Recent

no fossil record

KLINCKOWSTROEMIIDAE Camin & Gorirossi, 1955 Recent

no fossil record

PROMEGISTIDAE Kethley, 1979 Recent

no fossil record

MEGISTHANOIDEA Berlese, 1914 Recent

HOPLOMEGISTIDAE Camin & Gorirossi, 1955 Recent

no fossil record

MEGISTHANIDAE Berlese, 1914 Recent

no fossil record

PARANTENNULOIDEA Willmann, 1940 Recent

PARANTENNULIDAE Willmann, 1940 Recent

no fossil record

PHIODANIDAE Kethley, 1977b Recent

no fossil record

AENICTEQUOIDEA Kethley, 1979 Recent

AENICTEQUIDAE Kethley, 1979 Recent

no fossil record

EUPHYSALOZERCONIDAE Kim, 2008	Recent
no fossil record	
MESSORACARIDAE Kethley, 1977	Recent
no fossil record	
PHYSALOZERCONIDAE Kethley, 1977	Recent
no fossil record	
PTOCHACARIDAE Kethley, 1979	Recent
no fossil record	
MONOGYNASPIDA Camin & Gorirossi, 1955 (infrorder)	Palaeogene – Recent
MICROGYNIINA Trägårdh, 1942 (cohort)	Palaeogene – Recent
MICROGYNIOIDEA Trägårdh, 1942	Palaeogene – Recent
Microgynoidea sp. <i>in</i> Dunlop <i>et al.</i> (2013)	Pa Baltic amber
MICROGYNIIDAE Trägårdh, 1942	Recent
= MICROSEJIDAE Trägårdh, 1942	
no fossil record	
NOTHOGYNIDAE Walter & Kranz, 1999	Recent
no fossil record	
HEATHERELLINA author, date? (cohort)	Recent
HEATHERELLOIDEA Walter, 1997	Recent
HEATHERELLIDAE Walter, 1997	Recent
no fossil record	
UROPODOIDEA Kramer, 1881 (cohort)	Palaeogene – Recent
UROPODIAE Kramer, 1881 (subcohort)	Palaeogene – Recent
PROTODINYCHOIDEA Evans, 1957	Recent
PROTODINYCHIDAE Evans, 1957	Recent
no fossil record	
THINOZERCONOIDEA Halbert, 1915	Recent
THINOZERCONIDAE Halbert, 1915	Recent
no fossil record	
POLYASPIDOIDEA Berlese, 1913	Recent
DITHINOZERCONIDAE Ainscough, 1979	Recent
no fossil record	

POLYASPIDIDAE Berlese, 1913	Recent
no fossil record	
TRACHYTIDAE Trägårdh, 1938	Recent
no fossil record	
UROPODOIDEA Kramer, 1881	Palaeogene – Recent
BALOGHKASZABIIDAE Hirschmann, 1979	Recent
no fossil record	
BRASILUROPODIDAE Hirschmann, 1979	Recent
no fossil record	
CILLIBIDAE Trägårdh, 1944	Recent
no fossil record	
CLAUSIADINYCHIDAE Hirschmann, 1979	Recent
no fossil record	
CIRCOCYLLIBAMIDAE Sellnick, 1926	Recent
no fossil record	
CYLLIBULIDAE Hirschmann, 1979	Recent
no fossil record	
DERAIOPHORIDAE Trägårdh, 1952	Recent
no fossil record	
DINYCHIDAE Berlese, 1916	Recent
no fossil record	
DISCOURELLIDAE Baker & Wharton, 1952	Recent
no fossil record	
EUTRACHYTIDAE Trägårdh, 1944	Recent
no fossil record	
HUTUFEIDERIIDAE Hirschmann, 1979	Recent
no fossil record	
KASZABJBALOGHIIDAE Hirschmann, 1979	Recent
no fossil record	
MACRODINYCHIDAE Hirschmann, 1979	Recent

no fossil record

METAGYNURIDAE Balogh, 1943 Recent

no fossil record

NENTERIIDAE Hirschmann, 1979 Recent

no fossil record

OPLITIDAE Johnston, 1968 Recent

no fossil record

PHYMATODISCIDAE Hirschmann, 1979 Recent

no fossil record

PRODINYCHIDAE Berlese, 1917 Recent

no fossil record

ROTUNDABALOGHIIDAE Hirschmann, 1979 Recent

no fossil record

TERASEJASPIDAE Hirschmann, 1979 Recent

no fossil record

TREMATURIDAE Berlese, 1917 ?Palaeogene – Recent

= TREMATURELLIDAE Trägårdh, 1944

?Trematuridae *in* Lyubarsky & Perkovsky (2012) Pa Rovno amber

Trichouropoda Berlese, 1916 ?Palaeogene – Recent

?*Trichouropoda* sp. [as *Oodinychus* sp.] *in* Ramsay (1960) Qt New Zealand

TRICHOCYLLIBIDAE Hirschmann, 1979 Recent

no fossil record

TRICOPODELLIDAE Hirschmann, 1979 Recent

no fossil record

TRIGONUROPODIDAE Hirschmann *in* Wisniewski, 1979 Recent

no fossil record

UROACTINIIDAE Hirschmann & Zirngiebl-Nicol, 1964 Recent

no fossil record

URODIASPIDIDAE Trägårdh, 1944 Recent

no fossil record

URODINYCHIDAE Berlese, 1917	Palaeogene – Recent
<i>Uroobovella</i> Berlese, 1903	?Palaeogene – Recent
? <i>Uroobovella</i> sp. in Dunlop et al. (2013)	Pa Baltic amber
UROPODIDAE Kramer, 1881	Recent
no fossil record	
TRACHYUROPODOIDEA Berlese, 1917	Recent
TRACHYUROPODIDAE Berlese, 1917	Recent
no fossil record	
DIARTHROPHALLIAE Trägårdh, 1946 (subcohort)	Recent
DIARTHROPHALLOIDEA Trägårdh, 1946	Recent
DIARTHROPHALLIDAE Trägårdh, 1946	Recent
no fossil record	
HETEROZERCONINA author, date? (cohort)	Recent
HETEROZERCONOIDEA Berlese, 1892	Recent
DISCOZERCONIDAE Berlese, 1910	Recent
no fossil record	
HETEROZERCONIDAE Berlese, 1892	Recent
no fossil record	
GAMASINA Kramer, 1881 (cohort)	Palaeogene – Recent
Gamasina indet in Perkovsky et al. (2007)	Pa Rovno amber
EPICRIIAE Vitzthum, 1938 (subcohort)	Neogene – Recent
EPICRIOIDEA Berlese, 1885	Recent
EPICRIIDAE Berlese, 1885	Recent
no fossil record	
ZERCONOIDEA Berlese, 1892	Neogene – Recent
COPROZERCONIDAE Moraza & Lindquist, 1999	Recent
no fossil record	
ZERCONIDAE Berlese, 1892	Neogene – Recent
† <i>Paleozercon</i> Błaszk, Cokendolpher & Polyak, 1995	Neogene
14. <i>Paleozercon caverniculus</i> Błaszk, Cokendolpher & Polyak, 1995	Ne New Mexico
ARCTACARIAE Johnston, 1982 (subcohort)	Recent
ARCTACAROIDEA Evans, 1955	Recent

ARCTACARIDAE Evans, 1955	Recent
no fossil record	
PARASITIAE Reuter, 1909 (subcohort)	Palaeogene – Recent
PARASITOIDEA Oudemans, 1901	Palaeogene – Recent
PARASITIDAE Oudemans, 1901	Palaeogene – Recent
?Parasitidae indet. <i>in</i> Dunlop & Falkenhagen (2014)	Qt Germany
Aclerogamasus Athias, 1971	Palaeogene – Recent
15. <i>Aclerogamasus stenocornis</i> Witaliński, 2000	Pa Baltic amber
DERMANYSSIAE Evans & Till, 1997 (subcohort)	Palaeogene – Recent
VEIGAOIDEA Oudemans, 1939	Recent
VEIGAIIDAE Oudemans, 1939	Recent
= GAMASOLAELOAPTIDAE Oudemans, 1939	
no fossil record	
RHODACAROIDEA Oudemans, 1902	Palaeogene – Recent
DIGAMASELLIDAE Evans, 1954 ...[or 57?].	Palaeogene – Recent
Digamasellidae sp. <i>in</i> Perkovsky <i>et al.</i> (2007)	Pa Rovno amber
Dendrolaelaps Halbert, 1915	Neogene – Recent
16. <i>Dendrolaelaps fossilis</i> Hirschman, 1971	Ne Chiapas amber
EURYPARASITIDAE d'Antony, 1987	Recent
no fossil record	
GAMASIPHIDAE author, date?	Recent
no fossil record	
LAELOPTONYSSIDAE Womersley, 1956	Recent
no fossil record	
OLOGAMASIDAE Ryke, 1962	Recent
no fossil record	
PANTENIPHIDIDAE d'Antony, 1987	Recent
no fossil record	
RHODACARIDAE Oudemans, 1902	Recent
no fossil record	
TERANYSSIDAE Halliday, 2006	Recent
no fossil record	

EVIPHIDOIDEA Berlese, 1913	Quaternary–Recent
EVIPHIDIDAE Berlese, 1913	Recent
no fossil record	
MACROCHELIDAE Vitzthum, 1930	Quaternary–Recent
Macrocheles Latreille, 1829	Quaternary–Recent
<i>Macrocheles</i> sp. <i>in</i> Ramsay (1960)	Qt New Zealand
MEGALOELAPIDAE author, date?	Recent
no fossil record	
PACHYLAELAPIDAE Berlese, 1913	Recent
= NEOPARASITIDAE Oudemans, 1939	
= BULBOGAMASIDAE Gu, Wang & Duan, 1991	
no fossil record	
PARHOLASPIDIDAE Evans, 1956	Recent
no fossil record	
ASCOIDEA Oudemans, 1905	Palaeogene – Recent
AMEROSEIIDAE Evans <i>in</i> Hughs, 1961	Recent
no fossil record	
ASCIDAE Voigts & Oudemans, 1905	?Palaeogene – Recent
? <i>Ascidae</i> sp. <i>in</i> Dunlop <i>et al.</i> (2013)	Pa Baltic amber
HALOELAPIDAE Karg, 1965	Recent
no fossil record	
MELICHARIDAE Hirschmann, 1962	Recent
no fossil record	
PODOCINIDAE Berlese, 1913	Quaternary – Recent
<i>Podocinidae</i> sp. <i>in</i> Aoki (1974)	Qt Mizunami copal
PHYTOSEIOIDEA Berlese, 1916	Recent
BLATTISCOIIDAE Garman, 1948	Recent
no fossil record	
OTOPHEIDOMENIDAE Treat, 1955	Recent
no fossil record	

PHYTOSEIIDAE Berlese, 1916	Recent
no fossil record	
DERMANYSSOIDEA Kolenati, 1859	Palaeogene – Recent
DASYPONYSSIDAE Fonseca, 1940	Recent
no fossil record	
DERMANYSSIDAE Kolenati, 1859	Recent
no fossil record	
ENTONYSSIDAE Ewing, 1922	Recent
no fossil record	
HAEMOGAMASIDAE Oudemans, 1939	Recent
no fossil record	
HALARACHNIDAE Oudemans, 1906	Recent
no fossil record	
HIRSTONYSSIDAE Evans & Till, 1966	Recent
no fossil record	
HYSTRICHONYSSIDAE Keegan, Yunker & Baker, 1960	Recent
no fossil record	
IPIHOSSIDAE Kramer, 1886	Recent
no fossil record	
IXODORHYNCHIDAE Ewing, 1923	Recent
no fossil record	
LAE LAPIDAE Berlese, 1892	Palaeogene – Recent
Myrmozercon Berlese, 1902	Palaeogene – Recent
<i>Myrmozercon</i> sp. in Dunlop et al. (2014)	Pa Baltic amber
LARVAMIMIDAE Elzinga, 1993	Recent
no fossil record	
LEPTOLAE LAPIDAE Karg, 1978	Recent
no fossil record	
MACRONYSSIDAE Oudemans , 1936	Recent
no fossil record	

MANITHERIONYSSIDAE Radovsky & Yunker, 1971 Recent
no fossil record

OMENTOLAELEPTIDAE Fain, 1961 Recent
no fossil record

PNEUMOPHIONYSSIDAE Fonseca, 1940 Recent
no fossil record

RAILLIETIIDAE Vitzthum, 1942 Recent
no fossil record

RHINONYSSIDAE Trouessart, 1895 Recent
no fossil record

SPELAEORHYNCHIDAE Oudemans, 1902 Recent
no fossil record

SPINTURNICIDAE Oudemans, 1902 Recent
no fossil record

TRICOASPIDIDAE Gu, Wang & Li, 1991 Recent
no fossil record

VARROIDAE Delfinado & Baker, 1974 Recent
no fossil record

nomum dubium

1. *Ixodes tertiaris* Scudder, 1885 Pa Wyoming

c. 12,500 Recent species

ACARIFORMES

297 currently valid species of fossil acariform mite

- higher systematics and sequence of taxa follows the third edition of *A Manual of Acarology* (Krantz & Walter, eds, 2009), except that their orders are listed here as suborders, and suborders as infraorders to achieve some degree of consistency with other arachnid higher taxa throughout this list
- a putative Ordovician mite assigned to the derived Brachypylina group of the oribatids remains controversial and is not formally listed below

ACARIFORMES Zachvatkin, 1952 Devonian – Recent

= ACTINOTRICHIDA author, date?

TROMBIDIIFORMES Reuter, 1909 (suborder)..... Devonian – Recent

SPHAEROLICHIDA OConnor, 1984 (infraorder)..... Recent

LORDALYCOIDEA Grandjean, 1939 Recent

LORDALYCHIDAE Grandjean, 1939 Recent

= HYBALICIDAE Theron, 1974

no fossil record

SPHAEROLICOIDEA Berlese, 1913 Recent

SPHAEROLICHIDAE Berlese, 1913 Recent

no fossil record

PROSTIGMATA Kramer, 1877 (infraorder) Devonian – Recent

LABIDOSTOMMATIDES Lindquist, Krantz & Walter, 2009 (s.cohort) .. Palaeogene – Recent

LABIDOSTOMMATOIDEA Oudemans, 1906 Palaeogene – Recent

LABIDOSTOMMATIDAE Oudemans, 1906 Palaeogene – Recent

= NICOLETIELLIDAE Canestrini, 1891

Labidostomatidae sp. *in* Sidorchuk & Bertrand (2013) Pa Rovno amber

Labidostomatidae sp. *in* Sidorchuk & Bertrand (2013) Pa Bitterfeld amber

***Labidostomma* Kramer, 1879** Palaeogene – Recent

1. *Labidostomma (Nicoletiella) paleoluteum* Dunlop & Bertrand, 2011 Pa Baltic amber

2. *Labidostomma (Pseudocornutella) electri* Sidorchuk & Bertrand, 2013 .. Pa Baltic amber

***Sellnickiella* Feider & Vasiliu, 1969** Palaeogene – Recent

3. *Sellnickiella balticae* Sidorchuk & Bertrand, 2013 Pa Baltic amber

EUPODIDES Krantz, 1978 (supercohort) Devonian – Recent

BDELLIOIDEA Dugès, 1834 Cretaceous – Recent

BDELLIDAE Dugès, 1834 Cretaceous – Recent

Bdellidae sp. <i>in</i> Aoki (1974)	Qt	Mizunami copal
<i>Bdella</i> Latreille, 1795		Cretaceous – Recent
4. <i>Bdella bicincta</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
5. <i>Bdella bombycinia</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
6. <i>Bdella obconica</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
7. <i>Bdella vetusta</i> Ewing, 1937	K	Manitobian amber
<i>Bdelloides</i> Oudemans, 1937		Palaeogene – Recent
8. <i>Bdelloides lata</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
CUNAXIDAE Thor, 1902		Recent
no fossil record		
HALACAROIDEA Murray, 1877		Recent
HALACARIDAE Murray, 1877		Recent
no fossil record		
PEZIDAE Harvey, 1990		Recent
no fossil record		
EUPODOIDEA C. L. Koch, 1842		Palaeogene – Recent
COCC-EUPODIDAE Jesionowska, 2010		Recent
no fossil record		
DENDOCHAETIDAE Oliver, 2008		Recent
no fossil record		
EUPODIDAE C. L. Koch, 1842		Recent
no fossil record		
ERIORHYNCHIDAE Qin & Halliday, 1997		Recent
no fossil record		
PENTAPALPIDAE Oliver & Theron, 2000		Recent
no fossil record		
PENTHALEIDAE Oudemans, 1931		Recent
no fossil record		
PENTHALODIDAE Thor, 1933		Palaogene – Recent
<i>Penthalodes</i> Murray, 1877		Palaeogene – Recent
9. <i>Penthalodes tristiculus</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber

PROTERORHAGIIDAE Lindquist & Palacios-Vargas, 1991	Recent
no fossil record	
 RHAGIDIIDAE Oudemans, 1922	Paleogene – Recent
Rhagidiidae indet. <i>in</i> Judson & Wunderlich (2003)	Pa Baltic amber
 <i>Poecilophysis</i> O. P.-Cambridge, 1876	Paleogene – Recent
? <i>Poecilophysis</i> sp. <i>in</i> Judson & Wunderlich (2003)	Pa Baltic amber
+ <i>Zachardia</i> Judson & Wunderlich, 2003	Paleogene
10. <i>Zachardia flexipes</i> Judson & Wunderlich, 2003	Pa Baltic amber
 STRANDTMANNIIDAE Zacharda, 1979	Recent
no fossil record	
 TYDEOIDEA Kramer, 1877	Devonian – Recent
EREYNETIDAE Oudemans, 1931	Recent
= MICROEREUNETIDAE Bottazzi, 1950	
no fossil record	
 IOLINIDAE Pritchard, 1956	Recent
no fossil record	
 TRIOPHTYDEIDAE Andrè, 1980	Recent
= MEYERELLIDAE André, 1979	
no fossil record	
 TYDEIDAE Kramer, 1877	Devonian – Recent
+ <i>Palaeotydeus</i> Dubinin, 1962	Devonian – Recent
11. <i>Palaeotydeus devonicus</i> Dubinin, 1962	D Rhynie chert
+ <i>Parapotacarus</i> Dubinin, 1962	Devonian – Recent
12. <i>Paraprotacarus hirsti</i> Dubinin, 1962	D Rhynie chert
 TETRAPODILI sensu Oudemans, 1923	Triassic – Recent
TRIASACAROIDAE Lindquist & Sidorchuk <i>in</i> Sidorchuk et al., 2014	Triassic
TRIASACARIDAE Lindquist & Sidorchuk <i>in</i> Sidorchuk et al., 2014	Triassic
+ <i>Ampezzoa</i> Linquist & Grimaldi <i>in</i> Schmidt et al., 2012,	Triassic
13. <i>Ampezzoa triassica</i> Lindquist & Grimaldi <i>in</i> Schmidt et al., 2012*	Tr Italian amber
+ <i>Cheirolepidoptus</i> Sidorchuk & Lindquist <i>in</i> Sidorchuk et al. 2014	Triassic
14. <i>Cheirolepidoptus dolomiticus</i> Sidorchuk & Lindquist <i>in</i> Sidorchuk et al., 2014*	Tr Italian amber
+ <i>Minyacarus</i> Sidorchuk & Lindquist <i>in</i> Sidorchuk et al., 2014	Triassic
15. <i>Minyacarus aderces</i> Sidorchuk & Lindquist <i>in</i> Sidorchuk et al., 2014* ...	Tr Italian amber
+ <i>Triasacarus</i> Linquist & Grimaldi <i>in</i> Schmidt et al., 2012,	Triassic – Recent

16. <i>Triasacarus fedelei</i> Lindquist & Grimaldi <i>in Schmidt et al.</i> , 2012*	Tr	Italian amber
ERIOPHYOIDEA Nalepa, 1898		?Palaeogene – Recent
DIPTILOMIOPIDAE Keifer, 1944		Recent
no fossil record		
ERIOPHYIDAE Nalepa, 1898		?Palaeogene – Recent
Aculops Keifer, 1966		? Palaeogene – Recent
17. <i>Aculops keiferi</i> Southcott & Lange, 1971	?Pa	Australia
PHYTOOPTIDAE Murray, 1877		Neogene – Recent
= NALEPELLIDAE Roivainen, 1953		
no fossil record		
ANYSTIDES van der Hammen, 1972 (supercohort)		Cretaceous – Recent
ANYSTINA van der Hammen, 1972 (cohort)		Cretaceous – Recent
CAECULOIDEA Berlese, 1883		Paleogene – Recent
CAECULIDAE Berlese, 1883		Paleogene – Recent
Procaeculus Jacot, 1936		Paleogene – Recent
18. <i>Procaeculus dominicensis</i> Coineau & Poinar, 2001	Ne	Dominican amber
19. <i>Procaeculus eridanosae</i> Coineau & Magowski, 1994	Pa	Baltic amber
ADAMYSTOIDEA Cunliffe, 1957		Recent
ADAMYSTIDAE Cunliffe, 1957		Recent
= SAXIDROMIDAE Coineau, 1974		
no fossil record		
ANYSTOIDEA Oudemans, 1902		Cretaceous – Recent
ANYSTIDAE Oudemans, 1902		Cretaceous – Recent
Anystidae sp. <i>in</i> Aoki (1974)	Qt	Mizunami copal
Anystis von Heyden, 1826		Cretaceous – Recent
20. <i>Anystis malleator</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa	Baltic amber
21. <i>Anystis subnuda</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa	Baltic amber
22. <i>Anystis venustula</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
+ Mesoanystis Zacharda, 1985		Cretaceous
23. <i>Mesoanystis taymirensis</i> Zacharda, 1985*	K	Siberian amber
+ Palaeoerythracarus Zacharda, 1985		Palaeogene
24. <i>Palaeoerythracarus sachalinensis</i> Zacharda, 1985*	Pa	Sachalin amber
PSEUDOCHEYLIDAE Oudemans, 1909		Recent
= STIGMOCHEYLIDAE Kethley, 1990		
no fossil record		

TENERIFFIIDAE Thor, 1911b	Paleogene – Recent
Teneriffiidae sp. indet <i>in</i> Sayre et al. (1992)	Pa Baltic amber
PARATYDEOIDEA Baker, 1949	Recent
PARATYDEIDAE Baker, 1949	Recent
no fossil record	
STIGMOCHEYLIDAE Kethley, 1990	Recent
no fossil record	
POMERANTZIOIDEA Baker, 1949	Recent
POMERANTZIIDAE Baker, 1949	Recent
no fossil record	
PARASITENGONA Oudemans, 1909 (cohort)	Cretaceous – Recent
ERYTHRAIAE author, date? (subcohort)	Cretaceous – Recent
CALYPTOSTOMATOIDEA Oudemans, 1923	Recent
CALYPTOSTOMATIDAE Oudemans, 1923	Recent
no fossil record	
ERYTHRAEOIDEA Grandjean, 1947a	Cretaceous – Recent
larval Erythraeoidea <i>in</i> Zacharda & Krivolutskij (1985)	K Siberian amber
† Pararainbowia Dunlop, 2007	Cretaceous
25. <i>Pararainbowia martilli</i> Dunlop, 2007*	K Crato Formation
ERYTHRAEIDAE Robineau-Desvoidy, 1828	Paleogene – Recent
= LEPTIDAE Billberg, 1820	
= BALUSTIIDAE Grandjean, 1947	
Erythraeidae sp. <i>in</i> Aoki (1974)	Qt Mizunami copal
† Arytaena Menge, 1854 <i>in</i> C. L. Koch & Berendt, 1854	Paleogene
26. <i>Arytaena troguloides</i> Menge <i>in</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
Balaustium von Heyden, 1826	Paleogene – Recent
27. <i>Balaustium illustris</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
Erythraeus Latrielle, 1806	Paleogene – Recent
28. <i>Erythraeus bifrons</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa Baltic amber
29. <i>Erythraeus foveolatus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
30. <i>Erythraeus hirsutus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
31. <i>Erythraeus lagopus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
32. <i>Erythraeus longipes</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
33. <i>Erythraeus proavus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
34. <i>Erythraeus procerus</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa Baltic amber
35. <i>Erythraeus raripilus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber

36. *Erythraeus rostratus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
37. *Erythraeus saccatus* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- Leptus** Latrielle, 1796 Paleogene – Recent
38. *Leptus incertus* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- † **PROTERYTHRAEIDAE** Vercammen-Grandjean, 1973 Cretaceous
- † **Proterythraeus** Vercammen-Grandjean, 1973 Cretaceous
39. *Proterythraeus southcotti* Vercammen-Grandjean, 1973* K Manitoba amber
- SMARIDIDAE** Vitzthum, 1929 Paleogene – Recent
- Smarididae in Kulicka (1990) Pa Baltic amber
- TROMBIDIAE** author, date? (subcohort) Creteaceous – Recent
- trombidiid mites?**
40. *Megameropsis aquensis* Gourret, 1887 Pa Aix-en-Provence
41. *Pseudopachygnathus maculatus* Gourret, 1887 Pa Aix-en-Provence
- AMPHOTROMBIOIDEA** Zhang, 1998 Recent
- AMPHOTROMBIIDAE**, Zhang, 1998 Recent
- no fossil record
- ALLOTANAUPODOIDAE** Zhang & Fan, 2007 Recent
- ALLOTANAUPODIDAE** Zhang & Fan, 2007 Recent
- no fossil record
- TANAUPODOIDEA** Thor, 1935 Creteaceous – Recent
- TANAUPODIDAE** Thor, 1935 Creteaceous – Recent
- = ?AMPHOTROMBIIDAE Zhang, 1998
- = TANAUPODASTRIDAE Feider, 1959
- † **Atanaupodus** Judson & Mąkol, 2009 Cretaceous
42. *Atanaupodus bakeri* Judson & Mąkol, 2009 K Archingeay amber
- CHYZERIOIDEA** Womersley, 1954 Recent
- CHYZERIIDAE** Womersley, 1954 Recent
- no fossil record
- TROMBIDIIOIDEA** Leach, 1815 Paleogene – Recent
- ACHAEMENOTHROMBIIDAE** Saboori, Wohltmann & Hakimitabar, 2010 Recent
- no fossil record
- EUTROMBIDIIDAE** Thor, 1935 Recent

no fossil record

MICROTROMBIDIIDAE Thor, 1935 **Recent**

no fossil record

NEOTHROMBIIDAE Feider, 1955 **Recent**

no fossil record

TROMBIDIIDAE Leach, 1815 **Paleogene – Recent**

= PARATHROMBIIDAE Feider, 1959

Allothrombiidae Berlese, 1903 **Paleogene – Recent**

43. *Allothrombium clavipes* (C. L. Koch & Berendt, 1854) Pa Baltic amber

Paratrombiidae Bruyant, 1910 **Paleogene – Recent**

44. *Paratrombium rovniense* Konikiewicz & Mąkol, 2014 Pa Rovno amber

Trombidium Fabricius, 1775 **Paleogene – Recent**

45. *Trombidium crassipes* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

46. *Trombidium granulatum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

47. *Trombidium heterotrichum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

48. *Trombidium scrobiculatum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

NB: the next two families may be synonyms

WALCHIIDAE Ewing, 1946 **Recent**

no fossil record

TROMBICULOIDEA Ewing, 1929 **Recent**

AUDYANIDAE Southcott, 1987 **Recent**

no fossil record

JOHNSTONIANIDAE Thor, 1935 **Recent**

= NOTOTHROMBIIDAE Feider, 1959

no fossil record

NEOTROMBIDIIDAE Feider, 1959 **Recent**

no fossil record

LEEUWENHOEKIIDAE Womersley, 1944 **Recent**

no fossil record

TROMBELLIDAE Leach, 1815 **Recent**

no fossil record

TROMBICULIDAE Ewing, 1929 **Recent**

= VATACARIDAE Southcott, 1957

no fossil record

YUREBILLOIDEA Southcott, 1966 Recent

YUREBILLIDAE Southcott, 1996 Recent

no fossil record

HYDRACARNIDIAE van der Hoeven, 1849 (subcohort) Neogene – Recent

= HYDRACHNIDIA author, date?

= HYDRACHNELLAE author, date?

Undetermined water mites

Hygrobatoidea, Arrenuroidea or Lebertioidae *in* Poinar (1985) Ne Dominican amber

HYDRYPHANTOIDEA Piersig, 1896 Recent

CTENOTHYADIDAE Lundblad, 1936 Recent

no fossil record

EUPATRELLIDAE Viets, 1935 Recent

no fossil record

HYDRODROMIDAE Viets, 1936 Recent

= DIPLODONTIDAE Lundblad, 1927

no fossil record

HYDRYPHANTIDAE Piersig, 1896 Recent

= PROTZIIDAE Viets, 1926

no fossil record

MALGASACARIDAE Tuzovskij, Gerecke & Goldschmidt, 2007 Recent

no fossil record

RHYNCHOHYDRACARIDAE Lundblad, 1936 Recent

= CHATHROSPERCHONIDAE Lundblad, 1936

no fossil record

TERATOHYADIDAE Viets, 1929 Recent

no fossil record

THERMACARIDAE Sokolow, 1927 Recent

no fossil record

ZELANDOTHYADIDAE Cook, 1983 Recent

no fossil record

EYLAOIDEA Leach, 1815	Recent
APHEVIDERULICIDAE Gerecke, Smith & Cook, 1999.....	Recent
no fossil record	
EYLAIDAE Leach, 1815	Recent
no fossil record	
LIMNOCHARIDAE Grube, 1859	Recent
no fossil record	
PIERSIGIIDAE Oudemans, 1902	Recent
no fossil record	
HYDROVOLZIOIDEA Thor, 1905	Recent
ACHERONTACARIDAE Cook, 1967	Recent
no fossil record	
HYDROVOLZIIDAE Thor, 1905	Recent
= POLYXOHALACARIDAE Motas, 1972	
no fossil record	
HYDRACHNOIDEA Leach, 1815	Recent
HYDRACHNIDAE Leach, 1815	Recent
no fossil record	
LEBERTOIDEA Thor, 1900	Recent
ACUCAPITIDAE Wiles, 1996	Recent
no fossil record	
ANISITSIELLIDAE Koenicke, 1910	Recent
= MAMERSOPSIDAE Viets, 1914	
no fossil record	
BANDAKIOPSIDAE Panesar, 2004	Recent
no fossil record	
LEBERTIIDAE Thor, 1900	Recent
no fossil record	
NILOTONIIDAE Viets, 1929	Recent
no fossil record	

- OXIDAE Viets, 1926** **Recent**
no fossil record
- RUTRIPALPIDAE Solokow, 1834** **Recent**
no fossil record
- SPERCHONTIDAE Thor, 1900** **Recent**
no fossil record
- STYGOTONIIDAE Cook, 1992** **Recent**
no fossil record
- TEUTONIDAE Koenike, 1910** **Recent**
no fossil record
- TORRENTICOLIDAE Piersig, 1902** **Recent**
= ATRACTIDEIDAE Thor, 1902
no fossil record
- HYGROBATOIDEA C. L. Koch, 1842** **Recent**
ASTACOCROTONIDAE Thor, 1927 **Recent**
no fossil record
- ATURIDAE Thor, 1900** **Recent**
= BRADYPODIDAE Thor, 1900 [preoccupied]
= AXONOPSIDAE Viets, 1929
= LJANIIDAE Thor, 1929
no fossil record
- FELTRIIDAE Viets, 1926** **Recent**
no fossil record
- FERRADASIIDAE Cook, 1980** **Recent**
no fossil record
- FRONTIPODOPSIDAE Viets, 1931** **Recent**
no fossil record
- HYGROBATIDAE C. L. Koch, 1842b** **Recent**
no fossil record
- LETHAXONIDAE Cook, Smith & Harvey, 2000** **Recent**
no fossil record

LIMNESIIDAE Thor, 1900	Recent
= NEOTORRENTICOLIDAE Lundblad, 1936	
= EPALLAGOPODIDAE Viets, 1953	
no fossil record	
OMARTACARIDAE Cook, 1963	Recent
no fossil record	
PIONIDAE Thor, 1900	Recent
= CURVIPEDIDAE Thor, 1900	
= ACERCIDAE Thor, 1909	
= FORELIIDAE Thor, 1923	
= NAUTARACHNIDAE Walter, 1925	
= HYDROCHOREUTIDAE Viets, 1942	
no fossil record	
PONTARACHNIDAE Koenicke, 1910	Recent
no fossil record	
UNIONICOLIDAE Oudemans, 1909	Recent
= ATRACIDAE Thor, 1900	
= NEUMANIIDAE Thor, 1923	
no fossil record	
WETTINIDAE Cook, 1956	Recent
no fossil record	
ARRENUROIDEA Thor, 1900	Neogene – Recent
Family uncertain	
† <i>Protoarrenurus</i> Cook in Palmer, 1957	Neogene – Recent
49. <i>Protoarrenurus convergens</i> Cook in Palmer, 1957*	Ne Mojave Desert
ACALYPTONOTIDAE Walter, 1911	Recent
no fossil record	
AMOENACARIDAE Smith & Cook, 1997	Recent
no fossil record	
ARENOHYDRACARIDAE Cook, 1974	Recent
no fossil record	
ARRENURIDAE Thor, 1900	Recent
no fossil record	

- ATHIENEMANNIIDAE Viets, 1922** Recent
 = CHELOMIDEOPSIDAE Lundblad, 1962
 no fossil record
- BOGATIIDAE Motas & Tanasachi, 1938** Recent
 no fossil record
- CHAPPUISIDIDAE Motas & Tanasachi, 1946** Recent
 no fossil record
- GRETACARIDAE Viets, 1978** Recent
 no fossil record
- HARPAGOPALPIDAE Viets, 1924** Recent
 no fossil record
- HUNGAROHYDRACACARIDAE Motas & Tanasachi, 1959** Recent
 no fossil record
- KANTACARIDAE Imamura, 1959** Recent
 no fossil record
- KRENDOWSKIIDAE Viets, 1926** Recent
 no fossil record
- LAVERSIIDAE Cook, 1955** Recent
 no fossil record
- MIDEIDAE Thor, 1911a** Recent
 no fossil record
- MIDEOPSIDAE Koenicke, 1910** Recent
 no fossil record
- MOMONIIDAE Viets, 1926** Recent
 = STYGOMOMONIDAE Szalay, 1943
 no fossil record
- NEOACARIDAE Motas & Tanasachi, 1947** Recent
 no fossil record
- NIPPONACARIDAE Imamura, 1959** Recent
 no fossil record

NUDOMIDEOPSIDAE Smith, 1990	Recent
no fossil record	
UCHIDASTYGACARIDAE Imamura, 1956	Recent
no fossil record	
STYGOTHROMBIAE Thor, 1935 (subcohort)	Recent
STYGOTHROMBOIDEA Thor, 1935	Recent
STYGOTHROMBIIDAE Thor, 1935	Recent
ELEUTHERENCONIDES Oudemans, 1909 (supercohort)	Cretaceous – Recent
RAPHIGNATHINA Kethley, 1982 (cohort)	Cretaceous – Recent
MYOBIOIDEA Mégnin, 1877	Recent
MYOBIIDAE Mégnin, 1877	Recent
no fossil record	
PTERYGOSOMATOIDEA Oudemans, 1910	Recent
PTERYGOSOMATIDAE Oudemans, 1910	Recent
no fossil record	
RAPHIGNATHOIDEA Kramer, 1877	Paleogene – Recent
BARBUTIIDAE Robaux, 1975	Recent
no fossil record	
CALIGONELLIDAE Grandjean, 1944	Recent
no fossil record	
CAMEROBIIDAE Southcott, 1957	Paleogene – Recent
<i>Neophyllobius</i> Berlese, 1886	Paleogene – Recent
50. <i>Neophyllobius succineus</i> Bolland & Magowski, 1990	Pa Baltic amber
CRYPTOGNATHIDAE Oudemans, 1902	Paleogene – Recent
no fossil record	
DASYTHREIDAE Walter & Gerson, 1998	Recent
no fossil record	
EUPALOPSELLIDAE Willmann, 1952	Recent
no fossil record	
HOMOCALIGIDAE Wood, 1969	Recent
no fossil record	

MECOGNATHIDAE Gerson & Walter, 1998	Recent
no fossil record	
RAPHIGNATHIDAE Kramer, 1877	Recent
no fossil record	
STIGMAEIDAE Oudemans, 1931	Paleogene – Recent
Mediolata Canestrini, 1890	Paleogene – Recent
51. <i>Mediolata eocenia</i> Kuznetsov, Khaustov & Perkovsky, 2010.....	Pa Rovno amber
XENOCALIGONELLIDAE Gonzalez, 1978	Recent
no fossil record	
TETRANYCHOIDEA Donnadiet, 1876	Palaeogene – Recent
ALLOCHAETOPHORIDAE Reck, 1959	Recent
no fossil record	
LINOTETRANIDAE Baker & Pritchard, 1953	Recent
no fossil record	
TENUIPALPIDAE Berlese, 1913	Recent
no fossil record	
TETRANYCHIDAE Donnadiet, 1876	Palaeogene – Recent
= BRYOBIIDAE Berlese, date?	
Metatetranychus Oudemans, 1931	Palaeogene – Recent
52. <i>Metatetranychus gibbus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
Schizotetranychus Trägårdh, 1915	Palaeogene – Recent
53. <i>Schizotetranychus brevipes</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
TUCKERELLIDAE Baker & Pritchard, 1953	Recent
no fossil record	
CHEYLETOIDEA Leach, 1815	Cretaceous – Recent
CHEYLETIDAE Leach, 1815	Cretaceous – Recent
Chelytidae sp. indet <i>in</i> Bradley (1931)	Pa Green River
Cheyletus Latreille, 1796	Cretaceous – Recent
54. <i>Cheyletus burmiticus</i> Cockerell, 1917b.....	K Myanmar amber
55. <i>Cheyletus portentosus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
DEMODECIDAE Nicolet, 1855	Recent

no fossil record

HARPIRHYNCHIDAE Dubinin, 1957 Recent

no fossil record

OPHOPTIDAE Southcott, 1956 Recent

no fossil record

PSORERGATIDAE Dubinin *in* Bregatova et al., 1955 Recent

no fossil record

SYRINGOPHILIDAE Laviopierre, 1953 Recent

no fossil record

HETEROSTIGMATINA Berlese, 1899 (cohort) Cretaceous – Recent

TARSOCHYELOIDEA Atyeo & Baker, 1964 Recent

TARSOCHEYLIDAE Atyeo & Baker, 1964 Recent

no fossil record

HETEROCHYELOIDEA Trägårdh, 1950 Recent

HETEROCHEYLIDAE Trägårdh, 1950 Recent

no fossil record

DOLICHOCYBOIDEA Mahunka, 1970 Recent

CROTALOMORPHIDAE Lindquist & Kranz, 2002 Recent

no fossil record

DOLICHOCYBIDAE Mahunka, 1970 Recent

no fossil record

TROCHOMETRIDIOIDEA Mahunka, 1970 Recent

ATHYREACARIDAE Lindquist Kaliszewski & Rack, 1990 Recent

= BEMBIDIACARIDAE Khuastov, 2000

no fossil record

TROCHOMETRIDIIDAE Mahunka, 1970 Recent

no fossil record

SCUTACAROIDEA Oudemans, 1916 Recent

MICRODISPIDAE Cross, 1965 Recent

no fossil record

SCUTACARIDAE Oudemans, 1916 Recent

no fossil record

PYGEMEPhOROIDEA Cross, 1965 Palaeogene – Recent
Pygmephoroidae sp. in Magowski (1995) Pa Baltic amber

NEOPYGMEPHORIDAE Cross, 1965 Recent
no fossil record

PYGMEPHORIDAE Cross, 1965 Recent
no fossil record

SITROPTIDAE Mahunka, 1970 Recent
no fossil record

PYEMOTOIDEA Oudemans, 1937 Cretaceous – Recent
ACAROPHENACIDAE Cross, 1965 Cretaceous – Recent
† *Protophenax* Magowski, 1994 Cretaceous
56. *Protophenax kotejii* Magowski, 1994* K Russian amber

CARABOACARIDAE Mahunka, 1970 Recent
no fossil record

PYEMOTIDAE Oudemans, 1937 Recent
= *TROCHOMETRIDAE* Mahunka, 1970
Pyemotes Amerling, 1862 Palaeogene – Recent
57. *Pyemotes primus* Khaustov & Perkovsky, 2010 Pa Rovno amber

RESINACARIDAE Mahunka, 1975 Cretaceous – Recent
Protoresinacris Khaustov & Poinar, 2010 Cretaceous
58. *Protoresinacris brevipedis* Khaustov & Poinar, 2010* K Myanmar amber

TARSONEMOIDEA Canestrini & Fanzago, 1877 Quaternary – Recent
PODAPOLIPIDAE Ewing, 1922 Recent
no fossil record

TARSONEMIDAE Canestrini & Fanzango, 1877 Quaternary – Recent
Tarsonemidae sp. in Aoki (1974) Qt Mizunami copal

Cohort *incertae sedis*
CLOACAROIDEA Camin, Moss, Oliver & Singer, 1967 Recent
CLOACARIDAE Camin, Moss, Oliver & Singer, 1967 Recent
no fossil record

EPIMYODICIDAE Fain, Lukoschus & Rosmalen, 1982	Recent
no fossil record	
SARCOPTIFORMES author, date? (suborder)	Devonian – Recent
ENDEOSTIGMATA author, date? (infraorder)	Devonian – Recent
= PACHYGNATHINA author, date?	
ALYCINA author, date? (cohort)	
ALYCOIDEA Canestrini & Fanzago, 1877	Devonian – Recent
ALYCIDAE Canestrini & Fanzago, 1877	Devonian – Recent
= PACHYGNATHIDAE Kramer, 1877	
= BIMICHAELIIDAE Womersley, 1944	
† Protacarus Hirst, 1923	Devonian
59. <i>Protacarus crani</i> Hirst, 1923*	D Rhynie chert
GRANDJEANICIDAE Kethley, 1977a	Recent
no fossil record	
MICROPSAMMIDAE Coineau & Theorn, 1983	Recent
no fossil record	
NANORCHESTIDAE Grandjean, 1937	Devonian – Recent
† Protospeleorchestes Dubinin, 1962	Devonian – Recent
60. <i>Protospeleorchestes pseudoprotacarus</i> Dubinin, 1962*	D Rhynie chert
NEMATALYCINA author, date? (cohort)	Recent
NEMATALYCOIDEA Strenke, 1954	Recent
NEMATALYCIDAE Strenke, 1954	Recent
no fossil record	
PROTONEMATALYCIDAE Kethley, 1989 [superfamily correct?]	Recent
no fossil record	
TERPNACARINA author, date? (cohort)	Recent
OEHSERCHESTOIDEA Kethley, 1977a	Recent
OEHSERCHESTIDAE Kethley, 1977a	Recent
no fossil record	
TERPNACAROIDEA Grandjean, 1939	Recent
TERPNACARIDAE Grandjean, 1939	Recent
no fossil record	

ALICORHAGIINA author, date? (cohort)	Devonian – Recent
ALICORHAGIOIDEA Grandjean, 1939	Devonian – Recent
ALICORHAGIIDAE Grandjean, 1939	Devonian – Recent
† <i>Archaeacarus</i> Kethley & Norton in Kethley et al., 1989	Devonian
61. <i>Archaeacarus dubinini</i> Kethley & Norton in Kethley et al., 1989*	D Gilboa
† <i>Pseudoprotacarus</i> Dubinin, 1962	Devonian
62. <i>Pseudoprotacarus scoticus</i> Dubinin, 1962*	D Rhynie chert
ORIBATIDA Dugès, 1834 (infraorder)	Devonian – Recent
= CRYPTOSTIGMATA author, date?	
NB: see remarks on the Ordovician fossil above	
PALAEOSOMATA Grandjean, 1969 (supercohort)	Devonian–Recent
family uncertain	
† <i>Marcvipeda</i> Pérez-DA, 1988	Palaeogene
63. <i>Marcvipeda magallanes</i> Pérez-DA, 1988* [Acari incerte sedis?].....	Pa Patagonia, Chile
ACARONYCHOIDEA Grandjean, 1932	Recent
ACARONYCHIDAE Grandjean, 1932b	Recent
no fossil record	
ARCHAEONOTHRIDAE Grandjean, 1932	Recent
no fossil record	
CTENACAROIDEA Grandjean, 1954c	Devonian – Recent
ADELPHACARIDAE Grandjean, 1954c	Carbon. – Recent
† <i>Monoaphelacarus</i> Subías & Arillo, 2002	Carboniferous
64. <i>Monoaphelacarus carboniferus</i> Subías & Arillo, 2002*	C County Antrim
APHELACARIDAE Grandjean, 1954c	Recent
no fossil record	
CTENACARIDAE Grandjean, 1954b	Devonian – Recent
† <i>Ctenacaronychus</i> Subías & Arillo, 2002	Devonian
65. <i>Ctenacaronychus nortoni</i> Subías & Arillo, 2002*	D New York
† <i>Palaeoctenacarus</i> Subías & Arillo, 2002	Carboniferous
66. <i>Palaeoctenacarus simmsoi</i> Subías & Arillo, 2002*	C County Antrim
PALAEACAROIDEA Grandjean, 1932b	Recent
PALAEACARIDAE Grandjean, 1932b	Recent
no fossil record	

ENARTHRONOTA Grandjean, 1947b (supercohort)	Devonian – Recent
superfamily uncertain	
† DEVONACARIDAE Norton <i>in Norton et al.</i> , 1988	Devonian – Recent
† <i>Devonacarus</i> Norton <i>in Norton et al.</i> , 1988	Devonian – Recent
67. <i>Devonacarus sellnicki</i> Norton <i>in Norton et al.</i> , 1988*	D Gilboa
† PROTOCHTHONIIDAE Norton <i>in Norton et al.</i> , 1988	Devonian – Recent
† <i>Protochthonius</i> Norton <i>in Norton et al.</i> , 1988	Devonian – Recent
68. <i>Protochthonius gilboa</i> Norton <i>in Norton et al.</i> , 1988*	D Gilboa
BRACHYCHTHONIOIDEA Thor, 1934	Recent
BRACHYCHTHONIIDAE Thor, 1934	Recent
no fossil record	
ATOPOCHTHONIOIDEA Grandjean, 1948	Recent
ATOPOCHTHONIIDAE Grandjean, 1948	Recent
no fossil record	
PHYLLOCHTHONIIDAE Travé, 1967	Recent
no fossil record	
PTEROCHTHONIIDAE Grandjean, 1950	Recent
no fossil record	
HYPOCHTHONIOIDEA Berlese, 1910	Carbon. – Recent
ENIOCHTHONIIDAE Grandjean, 1947b	Recent
no fossil record	
HYPOCHTHONIIDAE Berlese, 1910	Carbon. – Recent
<i>Hypochthonius</i> C. L. Koch, 1835	Quaternary – Recent
69. <i>Hypochthonius rufulus</i> C. L. Koch, 1835 [Recent]	Qt Finland
† <i>Palaeohypochthonius</i> Subías & Arillo, 2002	Carboniferous
70. <i>Palaeohypochthonius jerami</i> Subías & Arillo, 2002*	C County Antrim
LOHMANNIIDAE Berlese, 1916	Recent
= XENOLOHMANNIDAE Balogh & Mahunka, 1969	
no fossil record	
MESOPLOPHORIDAE Ewing, 1917	Recent
= ARCHOPLOPHORIDAE Grandjean, 1965	
no fossil record	

PROTOLOPHOROIDEA Ewing, 1917	Carbon. – Recent
COSMOCHTHONIIDAE Grandjean, 1947b	Carbon. – Recent
† Carbochthonius Subías & Arillo, 2002	Carboniferous
71. <i>Carbochthonius antrimensis</i> Subías & Arillo, 2002*	C County Antrim
HAPLOCHTHONIIDAE van der Hammen, 1959	Recent
no fossil record	
PEDICULOCHELIDAE Lavoipierre, 1946	Recent
no fossil record	
PROTHOLOPHORIDAE Ewing, 1917	Carbon. – Recent
= APOLOPHORIDAE Niedbala, 1984	
† Archaeolophora Subías & Arillo, 2002	Carboniferous
72. <i>Archaeolophora bella</i> Subías & Arillo, 2002*	C County Antrim
SPHAEROCHTHONIIDAE Grandjean, 1947b	Recent
no fossil record	
HETEROCHTHONOIDEA Grandjean, 1954b	Recent
ARBORICHTHONIIDAE Balogh & Balogh, 1992	Recent
no fossil record	
HETEROCHTHONIIDAE Grandjean, 1954b	Recent
no fossil record	
TRICHTOCHTHONIIDAE Lee, 1982	Recent
no fossil record	
PARHYPOSOMATA Grandjean, 1969 (supercohort)	Carbon. – Recent
PARHYPOCHTHONIOIDEA Grandjean, 1932b	Carbon. – Recent
ELLIPTOCHTHONIIDAE Norton, 1975	Recent
no fossil record	
GEHYPOCHTHONIIDAE Strenzke, 1963	Carbon. – Recent
† Gehypochthonimimus Subías & Arillo, 2002	Carboniferous
73. <i>Gehypochthonimimus hibernicus</i> Subías & Arillo, 2002*	C County Antrim
PARHYPOCHTHONIIDAE Grandjean, 1932b	Recent
no fossil record	

MIXONOMATA Grandjean, 1969(supercohort)	Palaeogene – Recent
NEHYPOCHTHONOIDEA Norton & Metz, 1980	Recent
NEHYPOCHTHONIIDAE Norton & Metz, 1980	Recent
no fossil record	
EULOHMANNOIDEA Grandjean, 1931	Recent
EULOHMANNIIDAE Grandjean, 1931	Recent
no fossil record	
PERLOHMANNIOIDEA Grandjean, 1954b	Recent
PERLOHMANNIIDAE Grandjean, 1954b	Recent
no fossil record	
EPILOHMANNIOIDEA Oudemans, 1923	Recent
EPILOHMANNIIDAE Oudemans, 1923	Recent
= <i>LESSIRIIDAE</i> Oudemans, 1916	
no fossil record	
COLLOHMANNIOIDEA Grandjean, 1958a	Paleogene – Recent
COLLOHMANNIIDAE Grandjean, 1958a	Paleogene – Recent
<i>Collohmnia</i> Sellnick, 1922	Paleogene – Recent
74. <i>Collohmnia schusteri</i> Norton, 2006	Pa Baltic amber
† <i>Embolacarus</i> Sellnick, 1919	Palaeogene – Recent
75. <i>Embolacarus pergratus</i> Sellnick, 1919*	Pa Baltic amber
EUPYCTIMA Grandjean, 1967	Palaeogene – Recent
NB: Eupyctima is listed here as a mixonomatid clade, but is not recognised in all classifications, or else is removed from this group and given equal rank	
EUPHTHIRACAROIDEA Jacot, 1930	Palaeogene – Recent
EUPHTHIRACARIDAE Jacot, 1930	Palaeogene – Recent
<i>Microtritia</i> Märkel, 1964	Quaternary – Recent
76. <i>Microtritia minima</i> (Berlese, 1904) [Recent]	Qt Germany
<i>Rhysotritia</i> Märkel & Meyer, 1959	Quaternary – Recent
77. <i>Rhysotritia ardua</i> (C. L. Koch, 1841) [Recent]	Qt Germany
78. <i>Rhysotritia duplicata</i> (Grandjean, 1953) [Recent]	Qt Germany
ORIBOTRITIIDAE Grandjean, 1954b	Palaeogene – Recent
= <i>SABAHTRITIIDAE</i> Mahunka, 1987	
<i>Oribotritia</i> Jacot, 1924	Palaeogene – Recent
79. <i>Oribotritia pyropus</i> (Sellnick, 1919)	Pa Baltic amber
80. <i>Oribotritia translucida</i> Sellnick, 1931	Pa Baltic amber

SYNICHOTRITIIDAE Walker, 1965	Recent
no fossil record	
PHTHIRACAROIDEA Perty, 1841	Palaeogene – Recent
PHTHIRACARIDAE Perty, 1841	Palaeogene – Recent
= STEGANACARIDAE Niedbala, 1986	
Hoplophthiacarus Jacot, 1933	Quaternary – Recent
81. <i>Hoplophthiacarus pavidus</i> (Berlese, 1913) [Recent]	Qt Karelia, Russia
Phthiacarus Perty, 1841	Palaeogene – Recent
82. <i>Phthiacarus borealis</i> Trägårdh, date? [Recent]	Qt Karelia, Russia
83. <i>Phthiacarus multipunctus</i> (Sellnick, 1919)	Pa Baltic amber
Steganacarus Ewing, 1917	Quaternary – Recent
84. <i>Steganacarus applicatus</i> (Sellnick, 1920) [Recent]	Qt Denmark
85. <i>Steganacarus carinatus</i> (C. L. Koch, 1841) [Recent]	Qt Finland
86. <i>Steganacarus striculus</i> (C. L. Koch, 1835) [Recent]	Qt Europe
<i>Steganacarus</i> sp.	Qt Finland
DESMONOMATA Woodley, 1873 (supercohort)	Jurassic – Recent
NOTHRINA van der Hammen, 1982 (cohort)	Jurassic – Recent
= HOLOSOMATA author, date?	
CROTONIOIDEA Thorell, 1876	Jurassic – Recent
CAMISIIDAE Oudemans, 1900	Cretaceous – Recent
Camisia von Heyden, 1826	Paleogene – Recent
87. <i>Camisia foveolata</i> Hammer, 1955 [Recent]	Qt western Norway
88. <i>Camisia horrida</i> [Recent] <i>fossilis</i> Sellnick, 1919	Pa Baltic amber
i. = <i>Nothrus kuehli</i> Karsch, 1884	Pa Baltic amber
NB: unclear why the older name is the synonym	
89. <i>Camisia invenusta</i> (Michael, 1888) [Recent]	Qt western Norway
90. <i>Camisia lapponica</i> Trägårdh, 1910 [Recent]	Qt Karelia, Russia
† Eocamisia Bulanova-Zachvatkina, 1974	Cretaceous
91. <i>Eocamisia sukatshvae</i> Bulanova-Zachvatkina, 1974*	K Siberian amber
Platynothrus Berlese, 1913	Quaternary – Recent
92. <i>Platynothrus peltifer</i> (C. L. Koch, 1839) [Recent]	Qt Greenland
93. <i>Platynothrus punctatus</i> (L. Koch, 1879) [Recent]	Qt northern Europe
CROTONIIDAE Thorell, 1876	Neogene – Recent
= HOLONOTHRIDAE Wallwork, 1963	
Crotonia Thorell, 1876	Neogene – Recent
94. <i>Crotonia ramus</i> (Womersley, 1957)	Ne Australian retinite
HERMANNIIDAE Sellnick, 1928	Palaeogene – Recent
= GALAPAGACARIDAE P. Balogh, 1985	

<i>Hermannia</i> Nicolet, 1855	Palaeogene – Recent
95. <i>Hermannia gibba</i> (C. L. Koch, 1839) [Recent]	Qt Finland
96. <i>Hermannia reticulata</i> Thorell, 1871 [Recent]	Qt Subarctic – Arctic
97. <i>Hermannia scabra</i> (L. Koch, 1879) [Recent]	Qt Greenland
98. <i>Hermannia sellnicki</i> Norton, 2006	Pa Baltic amber
 MALACONOTHRIDAE Berlese, 1916	 Quaternary – Recent
<i>Malaconothrus</i> Berlese, 1904	Quaternary – Recent
99. <i>Malaconothrus monodactylus</i> (Michael, 1888) [Recent]	Qt Europe
<i>Trimalaconothrus</i> Berlese, 1916	Quaternary – Recent
100. <i>Trimalaconothrus maior</i> (Berlese, 1910) [Recent]	Qt northern Europe
 NANHERMANNIIDAE Sellnick, 1928	 Quaternary – Recent
<i>Nanhermannia</i> Berlese, 1913	Quaternary – Recent
101. <i>Nanhermannia coronata</i> Berlese, 1913 [Recent]	Qt Karelia, Russia
102. <i>Nanhermannia elegantula</i> Berlese, 1913 [Recent]	Qt Germany
 NOTHRIDAE Berlese, 1896	 Paleogene – Recent
<i>Nothrus</i> C. L. Koch, 1836	Paleogene – Recent
103. <i>Nothrus illautus</i> Sellnick, 1919	Pa Baltic amber
104. <i>Nothrus punctulum</i> Karsch, 1884	Pa Baltic amber
105. <i>Nothrus silvestris</i> Nicolet, 1855 [Recent]	Qt Europe
 TRHYPOCHTHONIIDAE Willmann, 1931	 Jurassic – Recent
= <i>ALLONOTHRIDAE</i> Lee, 1985	
= <i>MUCRONOTHRIDAE</i> Kunst, 1972	
= <i>PARALLONOTHRIDAE</i> Badejo, Woas & Beck, 2002	
= <i>TRHYPOCHTHONIELLIDAE</i> Knüller, 1957	
<i>Allonothrus</i> van der Hammen, 1953	Neogene – Recent
<i>Allonothrus</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
† <i>Juracarus</i> Krivolutsky in Krivolutsky & Krasilov, 1977	Jurassic – Recent
106. <i>Juracarus serratus</i> Krivolutsky in Krivolutsky & Krasilov, 1977	J Russian far east
<i>Mucronothrus</i> Trägårdh, 1931	Quaternary – Recent
107. <i>Mucronothrus nasalis</i> (Willmann, 1929) [Recent]	Qt Karelia, Russia
† <i>Palaeochthonius</i> Krivolutsky in Krivolutsky & Krasilov, 1977	Jurassic – Recent
108. <i>Palaeochthonius krasilovi</i> Krivolutsky in Kriv. & Krasilov, 1977	J Russian far east
<i>Trhypochthonius</i> Berlese, 1904	Palaeogene – Recent
109. <i>Trhypochthonius badiformis</i> Sellnick, 1931	Pa Baltic amber
110. <i>Trhypochthonius cladonicola</i> (Willmann, 1919) [Recent]	Qt Germany
111. <i>Trhypochthonius corniculatus</i> Sellnick, 1931	Pa Baltic amber
112. <i>Trhypochthonius tectorum</i> (Berlese, 1896) [Recent]	Qt Karelia, Russia

BRACHYPSYLINA Hull, 1918 (cohort)	Jurassic – Recent
= CIRCUMDEHISCENTIAE Grandjean, 1954b	
= PORONOTA Grandjean, 1954b [in part; taxon used for seven brachypyline superfamilies]	
 superfamily uncertain	
ARIBATIDAE Aoki, Takaku & Ito, 1994	Recent
no fossil record	
 HERMANNIELLOIDEA Grandjean, 1934	Paleogene – Recent
HERMANNIELLIDAE Grandjean, 1934	Paleogene – Recent
Hermannella Berlese, 1908	Paleogene – Recent
113. <i>Hermannella concamerata</i> Sellnick, 1931	Pa Baltic amber
114. <i>Hermannella tuberculata</i> Sellnick, 1919	Pa Baltic amber
Sacculobates Grandjean, 1962	Neogene – Recent
<i>Sacculobates</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
 PLASMOBATIDAE Grandjean, 1961a	Recent
no fossil record	
 NEOLIODOIDEA Sellnick, 1928	Palaeogene – Recent
= LIODOIDEA Grandjean, 1954b	
NEOLIODIDAE Sellnick, 1928	Palaeogene – Recent
= LIODIDAE Grandjean, 1954b	
Neoliodes Berlese, 1888	Palaeogene – Recent
= <i>Liodes</i> von Heyden, 1826 [preoccupied]	
115. <i>Neoliodes brevitarus</i> (Woolley, 1971)	Ne Chiapas amber
116. <i>Neoliodes dominicus</i> Heethoff, Helfen & Norton, 2009	Ne Dominican amber
117. <i>Neoliodes quadriscutatus</i> Sellnick, 1919	Pa Baltic amber
<i>Neoliodes</i> sp. in Norton & Poinar (1993) [as <i>Liodes</i>]	Ne Dominican amber
Platyliodes Berlese, 1917	Palaeogene – Recent
118. <i>Platyliodes ensigerus</i> (Sellnick, 1919)	Pa Baltic amber
Teleoliodes author, date?	Neogene – Recent
<i>Teleoliodes</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
 PLATEREMAEAOIDEA Trägårdh, 1926	Cretaceous – Recent
= GYMNODAMAEAOIDEA Grandjean, 1954a	
ALEURODAMAEIDAE Paschoal & Johnston, 1985	Recent
no fossil record	
 GYMNODAMAEIDAE Grandjean, 1954a	Paleogene – Recent
Gymnodamaeus Kulczynski, 1902	Paleogene – Recent
119. <i>Gymnodamaeus sepotisus</i> Sellnick, 1919	Pa Baltic amber

IDIODAMEIDAE Paschoal, 1987	Recent
no fossil record	
LICNOBELBIDAE Grandjean, 1965a	Recent
no fossil record	
LICNODAMEIDAE Grandjean, 1954b	Recent
= NACUNANSELLIDAE author, date	
no fossil record	
LYRIFISSIELLIDAE Paschoal, 1987	Recent
no fossil record	
PEDROCORTESELLIDAE Paschoal, 1987	Recent
no fossil record	
PHEROLIODIDAE Paschoal, 1987	Recent
= HAMMERIELLIDAE Paschoal, 1987	
= NOOLIODIDAE Paschoal, 1987	
no fossil record	
PLATEREMAEIDAE Trägårdh, 1926	Cretaceous – Recent
Rasnitsynella Krivoluckij, 1976	Cretaceous
120. <i>Rasnitsynella punctulata</i> Krivoluckij, 1976	K Taymir amber
DAMAEOIDEA Berlese, 1896	Paleogene – Recent
DAMAEIDAE Berlese, 1896	Paleogene – Recent
Damaeidae sp. in Aoki (1974)	Qt Mizunami copal
Belba von Heyden, 1826	Quaternary – Recent
121. <i>Belba compta</i> (Kulczynski, 1902) [Recent]	Qt western Norway
122. <i>Belba cornyops</i> (Hermann, 1804)* [Recent]	Qt Finland
† Belbites Pampaloni, 1902	Neogene
123. <i>Belbites disodilis</i> Pampaloni, 1902*	Ne? Sicily
Damaeobelba Sellnick, 1928	Quaternary – Recent
124. <i>Damaeobelba minutissima</i> (Sellnick, 1920) [Recent]	Qt Germany
Damaeus C. L. Koch, 1835	Paleogene – Recent
125. <i>Damaeus auritus</i> C. L. Koch, 1835* [Recent]	Qt Finland
126. <i>Damaeus genadensis</i> Sellnick, 1931	Pa Baltic amber
Spatiodamaeus Bulanova-Zachvatkina, 1967	Quaternary – Recent
127. <i>Spatiodamaeus verticillipes</i> (Nicolet, 1855)* [Recent]	Qt Finland

CEPHEOIDEA Berlese, 1896	Cretaceous – Recent
= EUTEGOIDEA Balogh, 1965	
ANDEREMAEIDAE Balogh, 1972	Recent
no fossil record	
CEPHEIDAE Berlese, 1896	Cretaceous – Recent
= COMPATOZETIDAE Luxton, 1988	
Cepheus C. L. Koch, 1835	Paleogene – Recent
128. <i>Cepheus cepheiformis</i> (Nicolet, 1855) [Recent]	Qt Finland
129. <i>Cepheus dentatus</i> (Michael, 1888) [Recent]	Qt Finland
130. <i>Cepheus implicatus</i> (Sellnick, 1919)	Pa Baltic amber
131. <i>Cepheus latus</i> C. L. Koch, 1835* [Recent]	Qt Finland
Epterotegaeus Berlese, 1916	Cretaceous – Recent
132. <i>Epterotegaeus bitranslammellatus</i> Arillo & Subías, 2002	K Álava amber
Ommatocepheus Berlese, 1913	Cretaceous – Recent
133. <i>Ommatocepheus nortoni</i> Arillo, Subías & Shtanchaeva, 2008	K Álava amber
CEROCEPHEIDAE Mahunka, 1986	Recent
no fossil record	
EUTEGAEIDAE Balogh, 1965	Recent
= PTEROZETIDAE Luxton, 1988	
no fossil record	
MICROTEGEIDAE Balogh, 1972	Recent
no fossil record	
NODOCEPHEIDAE Piffl, 1972	Recent
no fossil record	
NOSYBEIDAE Mahunka, 1994	Recent
no fossil record	
PTEROBATIDAE Balogh & Balogh, 1992	Recent
no fossil record	
POLYPTEROZETOIDEA Grandjean, 1959	Recent
PODOPTEROTEGAEIDAE Piffl, 1972	Recent
no fossil record	
POLYPTEROZETIDAE Grandjean, 1959	Recent
no fossil record	

TUMEROZETIDAE Hammer, 1966	Recent
no fossil record	
MICROZETOIDEA Grandjean, 1936a	Recent
MICROZETIDAE Grandjean, 1936a	Recent
no fossil record	
AMEROIDEA Bulanova-Zachvatkina, 1957	Palaeogene – Recent
= AMEROBELBOIDEA Grandjean, 1954b	
= CALEREMEIOIDEA Grandjean, 1965c	
AMERIDAE Bulanova-Zachvatkina, 1957	Recent
no fossil record	
AMEROBELBIDAE Grandjean, 1961b	Recent
no fossil record	
BASILOBELBIDAE Balogh, 1961	Recent
no fossil record	
CALEREMAEIDAE Grandjean, 1965c	Palaeogene – Recent
Caleremaeus Berlese, 1910	Palaeogene – Recent
134. <i>Caleremaeus gleso</i> Sellnick, 1931	Pa Baltic amber
CTENOSELBIDAE Grandjean, 1965b	Recent
no fossil record	
DAMEOLIDAE Grandjean, 1965b	Recent
no fossil record	
EREMOBELBIDAE Balogh, 1961	Recent
no fossil record	
EREMULIDAE Grandjean, 1965b	Recent
no fossil record	
HETEROSELBIDAE Balogh, 1961	Recent
no fossil record	
HUNGAROBELBIDAE Miko & Travé, 1996	Recent
no fossil record	
STAUROBATIDAE Grandjean, 1966	Recent

no fossil record

ZETORCHESTOIDEA Michael, 1898 Cretaceous – Recent

- = EREMAEOIDEA Oudeman, 1900
- = NIPHOCEPHOIDEA Travé, 1959 [a separate superfamily in some studies]

† ARCHAORCHESTIDAE Arillo & Subías, 2000 Cretaceous

† Plategeocranus Sellnick, 1919 Palaeogene

- 135. *Plategeocranus sulcatus* (Karsch, 1884)*

† Strieremaeus Sellnick, 1919 Cretaceous – Recent

- = † Archaeorchestes Arillo & Subías, 2000

- 136. *Strieremaeus illibatus* Sellnick, 1919

- 137. *Strieremaeus minguezae* (Arillo & Subías, 2000)

Pa Baltic amber

K Álava amber

EREMAEIDAE Oudemans, 1900 Paleogene – Recent

Eremaeus C. L. Koch, 1836 Paleogene – Recent

- 138. *Eremaeus hepaticus* C. L. Koch, 1835* [Recent]

Qt Germany

- 139. *Eremaeus oblongus* [Recent] *fossilis* Sellnick, 1919

Pa Baltic amber

Eueremaeus Mihelcic, 1963 Quaternary – Recent

- 140. *Eueremaeus silvestris* (Forsslund, 1956) [Recent]

Qt Finland

† Gradidorsum Sellnick, 1919 Palaeogene – Recent

- 141. *Gradidorsum asper* Sellnick, 1919*

Pa Baltic amber

MEGEREMAEIDAE Woolley & Higgins, 1968 Recent

no fossil record

NIPHOCEPHEIDAE Travé, 1959 Recent

no fossil record

ZETORCHESTIDAE Michael, 1898 Palaeogene – Recent

- Zetorchestidae spp. *in* Sidorshuk & Norton (2011)

Pa Rovno amber

GUSTAVIOIDEA Oudemans, 1900 Jurassic – Recent

- = LIACAROIDEA Sellnick, 1928

ASTEGISTIDAE Balogh, 1961 Jurassic – Recent

Astegistes Hull, 1916 Quaternary – Recent

- 142. *Astegistes pilosus* (C. L. Koch, 1840) [Recent]

Qt Karelia, Russia

Cultroribula Berlese, 1908 Jurassic – Recent

- 143. *Cultroribula jurassica* Krivolutsky *in* Krivolutsky & Krasilov, 1977

J Russian far east

- 144. *Cultroribula lauta* Sellnick, 1931

Pa Baltic amber

- 145. *Cultroribula superba* Sellnick, 1931

Pa Baltic amber

GUSTAVIIDAE Oudemans, 1900	Quaternary – Recent
<i>Gustavia</i> Kramer, 1879	Quaternary – Recent
146. <i>Gustavia microcephala</i> (Nicolet, 1855) [Recent]	Qt Finland
 KODIAKELLIDAE Hammer, 1967	Recent
no fossil record	
 LIACARIDAE Sellnick, 1928	Quaternary – Recent
= XENILLIDAE Woolley & Higgins, 1966	
<i>Adoristes</i> Hull, 1916	Quaternary – Recent
147. <i>Adoristes ovatus</i> (C. L. Koch, 1839)* [Recent]	Qt northern Europe
<i>Liacarus</i> Michael, 1898	Quaternary – Recent
148. <i>Liacarus coracinus</i> (C. L. Koch, 1841) [Recent]	Qt Finland
<i>Xenillus</i> Robineau-Desvoidy, 1839	Paleogene – Recent
149. <i>Xenillus tegeocraniformis</i> (Sellnick, 1919)	Pa Baltic amber
 MULTORIBULIDAE Balogh, 1972	Recent
no fossil record	
 PELOPPIIDAE Balogh, 1943	Paleogene – Recent
<i>Ceratoppia</i> Berlese, 1908	Paleogene – Recent
150. <i>Ceratoppia bipilis fossilis</i> Sellnick, 1919	Pa Baltic amber
ii. = <i>Oribates politus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
151. <i>Ceratoppia quadridentata</i> (Haller, 1882) [Recent]	Qt Finland
 TENUIALIDAE Jacot, 1929	Quaternary – Recent
<i>Hafenrefferia</i> Oudemans, 1906	Quaternary – Recent
152. <i>Hafenrefferia gilvipes</i> (C. L. Koch, 1839)* [Recent]	Qt Finland
 CARABODOIDEA C. L. Koch, 1843b	Palaeogene – Recent
= OCTOCEPHOIDEA Balogh, 1961	
CARABOCEPHEIDAE Mahunka, 1986	Recent
no fossil record	
 CARABODIDAE C. L. Koch, 1843b	Palaeogene – Recent
<i>Carabodes</i> C. L. Koch, 1835	Palaeogene – Recent
153. <i>Carabodes areolatus</i> Berlese, 1916 [Recent]	Qt Karelia, Russia
154. <i>Carabodes coriaceus</i> C. L. Koch, 1835* [Recent]	Qt Finland
155. <i>Carabodes coriaceus</i> [Recent] <i>fossilis</i> Sellnick, 1931	Pa Baltic amber
156. <i>Carabodes dissonus</i> Sellnick, 1931	Pa Baltic amber
157. <i>Carabodes gerberi</i> Sellnick, 1931	Pa Baltic amber
158. <i>Carabodes labyrinthicus</i> (Michael, 1879) [Recent]	Qt Europe

159. *Carabodes labyrinthicus* [Recent] *fossilis* Sellnick, 1931 Pa Baltic amber
160. *Carabodes marginatus* (Michael, 1884) [Recent] Qt Finland
161. *Carabodes minusculus* Berlese, 1923 [Recent] Qt Germany
162. *Carabodes ornatus* Storkan, 1925 [Recent] Qt Finland
163. *Carabodes subarcticus* Trägårdh, 1902 [Recent] Qt Finland
164. *Carabodes willmanni* Bernini, 1975 [Recent] Qt western Norway
? *Carabodes* sp. in Norton & Poinar (1993) Ne Dominican amber
- † *Carabodites* Pampaloni, 1902 Neogene?
165. *Carabodites pavesii* Pampaloni, 1902* Ne? Sicily
- Odontocepheus Berlese, 1913** Quaternary – Recent
166. *Odontocepheus elongatus* (Michael, 1879)* [Recent] Qt Finland
- DAMPFIELLIDAE Balogh, 1961** Recent
- no fossil record
- HEXOPPIIDAE Balogh, 1983** Recent
- no fossil record
- LUXTONIIDAE Mahunka, 2001** Recent
- no fossil record
- NIPPOBODIDAE Aoki, 1959** Recent
- no fossil record
- OTOCEPHEIDAE Balogh, 1961** Paleogene – Recent
- Dolicheremaeus* Jacot, 1938 Neogene – Recent
Dolicheremaeus sp. in Norton & Poinar (1993) Ne Dominican amber
- Otocepheus Berlese, 1905** Paleogene – Recent
167. *Otocepheus niger* Sellnick, 1931 Pa Baltic amber
168. *Otocepheus praesignis* Sellnick, 1931 Pa Baltic amber
- TOKUNOCEPHEIDAE Aoki, 1966a** Recent
- no fossil record
- OPPIOIDEA Grandjean, 1951** Palaeogene – Recent
- = EREMELLOIDEA Balogh, 1961 [in part]
= TRIZETOIDEA Ewing, 1917 [in part]
- AUTOGNETIDAE Grandjean, 1960b** Quaternary – Recent
- Conchogneta Grandjean, 1963** Quaternary – Recent
169. *Conchogneta traegardhi* (Forsslund, 1947) [Recent] Qt Finland
- ARCEREMAEIDAE Balogh, 1972** Recent

no fossil record

BORHIDIIDAE Balogh, 1983 **Recent**

no fossil record

CHAVINIIDAE Balogh, 1983 **Recent**

no fossil record

ENANTIOOPPIIDAE Balogh, 1983 **Recent**

no fossil record

EPIMERELLIDAE Ayyildiz & Luxton, 1989 **Recent**

no fossil record

GRANULOPPIIDAE Balogh, 1983 **Recent**

no fossil record

MACHADOBELBIDAE Balogh, 1972 **Recent**

no fossil record

MACHUELLIDAE Balogh, 1893 **Recent**

no fossil record

NOSYBELBIDAE Mahunka, 1994 **Recent**

no fossil record

OPPIIDAE Grandjean, 1951 **Palaeogene – Recent**

Dissorrhina Hull, 1916 **Quaternary – Recent**

170. *Dissorrhina ornata* (Oudemans, 1900)* **[Recent]** Qt Germany

Oppia C. L. Koch, 1836 **Palaeogene – Recent**

171. *Oppia angustum* (Sellnick, 1931) Pa Baltic amber

172. *Oppia cervicornu* (Sellnick, 1919) Pa Baltic amber

173. *Oppites hurdi* Woolley, 1971 Ne Chiapas amber

174. *Oppia longilamellata* **[Recent]** *fossilis* (Sellnick, 1931) Pa Baltic amber

175. *Oppia medium* (Sellnick, 1931) Pa Baltic amber

176. *Oppia mexicana* (Woolley, 1971) Ne Chiapas amber

177. *Oppia setigera* (Woolley, 1971) Ne Chiapas amber

178. *Oppia sucinum* (Sellnick, 1931) Pa Baltic amber

?*Oppia* sp. in Norton & Poinar (1993) Ne Dominican amber

Oppiella Jacot, 1937 **Quaternary – Recent**

179. *Oppiella nova* (Oudemans, 1902)* **[Recent]** Qt northern Europe

180. *Oppiella ornata* (Oudemans, 1900) **[Recent]** Qt western Norway

181. *Oppiella splendens* (C. L. Koch, 1841) **[Recent]** Qt western Norway

182. <i>Oppiella subpectinata</i> (Oudemans, 1900) [Recent]	Qt	northern Europe
183. <i>Oppiella translamellata</i> (Willmann, 1923) [Recent]	Qt	northern Europe
† Oppites Pampaloni, 1902		Neogene
184. <i>Oppites melilli</i> Pampaloni, 1902*	Ne?	Sicily
Ramusella Hammer, 1962		Quaternary – Recent
185. <i>Ramusella clavipectinata</i> (Michael, 1885) [Recent]	Qt	Germany
OXYAMERIDAE Aoki, 1965		Recent
no fossil record		
PAPILLONOTIDAE Balogh, 1983		Recent
no fossil record		
PLATYAMERIDAE Balogh & Balogh, 1983		Recent
no fossil record		
QUADROPPIIDAE Balogh, 1983		Recent
no fossil record		
RHYNCHORIBATIDAE Balogh, 1961		Recent
no fossil record		
SPINOZETIDAE Balogh, 1972		Recent
no fossil record		
STERNOPPIIDAE Balogh & Mahunka, 1969		Recent
no fossil record		
SUCTOBELBIDAE Jacot, 1938		Palaeogene – Recent
Suctobelbella Jacot, 1937		Palaeogene – Recent
186. <i>Suctobelbella falcata</i> (Forsslund, 1941) [Recent]	Qt	Germany
187. <i>Suctobelbella latirostris</i> (Strenzke, 1950) [Recent]	Qt	Germany
188. <i>Suctobelbella longirostris</i> (Forsslund, 1941) [Recent]	Qt	western Norway
189. <i>Suctobelbella sarekensis</i> (Forsslund, 1941) [Recent]	Qt	Europe
190. <i>Suctobelbella similis</i> (Forsslund, 1941) [Recent]	Qt	Germany
191. <i>Suctobelbella subcornigera</i> (Forsslund, 1941) [Recent]	Qt	Germany
192. <i>Suctobelbella subtrigona</i> (Oudemans, 1916) [Recent]	Qt	Europe
193. <i>Suctobelbella subtrigona</i> [Recent] <i>fossilis</i> (Sellnick, 1931)	Pa	Baltic amber
TERATOPPIIDAE Balogh, 1983		Recent
no fossil record		
TETRACONDYLIDAE Aoki, 1961		Recent

no fossil record

THYRISOMIDAE Grandjean, 1954b	Quaternary – Recent
Banksinoma Oudemans, 1930	Quaternary – Recent
194. <i>Banksinoma lanceolata</i> (Michael, 1885)* [Recent]	Qt Europe

TRIZETIDAE Ewing, 1917	Recent
no fossil record	

TUPAREZETIDAE Balogh, 1972	Recent
no fossil record	

TECTOCEPHEOIDEA Grandjean, 1954b	Paleogene – Recent
TECTOCEPHEIDAE Oudemans, 1900	Paleogene – Recent
<i>Tectocepheus</i> Berlese, 1895	Paleogene – Recent
195. <i>Tectocepheus minor</i> Berlese, 1903 [Recent]	Qt western Norway
196. <i>Tectocepheus similis</i> Sellnick, 1931	Pa Baltic amber
197. <i>Tectocepheus velatus</i> (Michael, 1880)* [Recent]	Qt northern Europe

HYDROZETOIDEA Grandjean, 1954b	Jurassic – Recent
HYDROZETIDAE Grandjean, 1954b	Jurassic – Recent
<i>Hydrozetes</i> Berlese, 1902	Jurassic – Recent
198. <i>Hydrozetes confervae</i> (Schrank, 1791) [Recent]	Qt western Norway
199. <i>Hydrozetes lacustris</i> (Michael, 1882)* [Recent]	Qt northern Europe
200. <i>Hydrozetes oryktosis</i> Woolley, 1969	Qt Michigan
<i>Hydrozetes</i> sp. in Sivhead & Wallwork (1978)	J Sweden

LIMNOZETIDAE Thor, 1937	Quaternary – Recent
<i>Limnozetes</i> Hull, 1916	Quaternary – Recent
201. <i>Limnozetes ciliatus</i> (Schrank, 1803)* [Recent]	Qt northern Europe
202. <i>Limnozetes rugosus</i> (Sellnick, 1923) [Recent]	Qt northern Europe

AMERONOTHROIDEA Willmann, 1931	Quaternary – Recent
AMERONOTHRIDAE Willmann, 1931	Quaternary – Recent
<i>Ameronothrus</i> Berlese, 1896	Quaternary – Recent
203. <i>Ameronothrus lineatus</i> (Thorell, 1871)* [Recent]	Qt Europe / Greenland
204. <i>Ameronothrus maculatus</i> (Michael, 1882) [Recent]	Qt western Norway

FORTUYNIIDAE van der Hammen, 1963	Recent
no fossil record	

SELENORIBATIDAE Schuster, 1963	Recent

no fossil record

TEGEOCRANELLIDAE Balogh, 1987 **Recent**

no fossil record

CYMBAEREMAEOIDEA Sellnick, 1928 **Jurassic – Recent**

CYMBAEREMAEIDAE Sellnick, 1928 **Jurassic – Recent**

= AMETROPROCTIDAE Subías, 2004

= SCAPHEREMAEIDAE Subías, 2004

Ametroproctus Higgins & Woolley, 1968 **Cretaceous – Recent**

205. *Ametroproctus valeriae* Arillo, Subías & Shtanchaeva, 2009 K San Just amber

Cymbaeremaeus Berlese, 1896 **Paleogene – Recent**

206. *Cymbaeremaeus cymba* (Nicolet, 1855)* [Recent] Qt northern Europe

† **Jureremeus Krivolutsky in Krivolutsky & Krasilov, 1977** **Jurassic**

207. *Jureremeus foveolatus* Krivolutsky in Krivolutsky & Krasilov, 1977* J Russian far east

208. *Jureremeus phippsi* Selden, Baker & Phipps, 2008 J Yorkshire, UK

Scapheremaeus Berlese, 1910 **Paleogene – Recent**

209. *Scapheremaeus undosus* Sellnick, 1919 Pa Baltic amber

† **Tectocymba Sellnick, 1919** **Paleogene – Recent**

210. *Tectocymba rara* Sellnick, 1919* Pa Baltic amber

EREMAEZOZETOIDEA Piffl, 1972 **Paleogene – Recent**

= IDIOZETOIDEA Aoki, 1976

EREMAEZETIDAE Piffl, 1972 **Paleogene – Recent**

Eremaezetes Berlese, 1913 **Paleogene – Recent**

= † *Scutoribates* Sellnick, 1919

Eremaezetes sp. in Norton & Poinar (1993) Ne Dominican amber

IDIOZETIDAE Aoki, 1976 **Recent**

no fossil record

LICNEREMAEOIDEA Grandjean, 1931 **Palaeogene – Recent**

= CHARASSOBATOIDEA Grandjean, 1958b

ADHAESOZETIDAE Hammer, 1973 **Recent**

no fossil record

CHARASSOBATIDAE Grandjean, 1958b **Recent**

no fossil record

DENDEROEREMAEIDAE Behan-Pelletier, Eamer & Clavton, 2005 **Recent**

no fossil record

EREMELLIDAE Balogh, 1961	Recent
no fossil record	
LAMELLAREIDAE Balogh, 1972	Recent
no fossil record	
LICNEREMAEIDAE Grandjean, 1931	Palaeogene – Recent
Licneremaeus Paoli, 1908	Palaeogene – Recent
211. <i>Licneremaeus fritschi</i> Sellnick, 1931	Pa Baltic amber
212. <i>Licneremaeus licnophorus</i> (Michael, 1882) [Recent]	Qt Germany
MICREREMIDAE Grandjean, 1954b	Jurassic – Recent
Micreremus Grandjean, 1954b[not Berlese 1908?]	Paleogene – Recent
213. <i>Micreremus brevipes</i> (Michael, 1888)* [Recent]	Qt northern Europe
214. <i>Micreremus reticulatus</i> Sellnick, 1931	Pa Baltic amber
215. <i>Micreremus scrobiculatus</i> Sellnick, 1931	Pa Baltic amber
PASSALOZETIDAE Grandjean, 1954b	Quaternary – Recent
Passalozetes Grandjean, 1932a	Quaternary – Recent
216. <i>Passalozetes africanus</i> Grandjean, 1932a [Recent]	Qt Finland
SCUTOVERTICIDAE Grandjean, 1954b	Neogene – Recent
Arthrovertex Balogh, 1970	Neogene – Recent
217. <i>Arthrovertex hurdi</i> (Woolley, 1971)	Ne Chiapas amber
<i>Arthrovertex</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
Scutovertex Michael, 1879	Quaternary – Recent
218. <i>Scutovertex minutus</i> (C. L. Koch, 1835) [Recent]	Qt Germany
PHENOPELOPOIDEA Petrunkevitch, 1955a	Palaeogene – Recent
PHENOPELOPIDAE Petrunkevitch, 1955a	Palaeogene – Recent
= PELOPIDAE author, date?	
Eupelops Ewing, 1917	Palaeogene – Recent
219. <i>Eupelops acromios</i> (Hermann, 1804) [Recent]	Qt Finland
220. <i>Eupelops curtipilus</i> (Berlese, 1916) [Recent]	Qt Germany
221. <i>Eupelops occultus</i> (C. L. Koch, 1835) [Recent]	Qt Kerelia, Russia
222. <i>Eupelops plicatus</i> (C. L. Koch, 1835) [Recent]	Qt northern Europe
223. <i>Eupelops punctulatus</i> (Sellnick, 1931)	Pa Baltic amber
224. <i>Eupelops uraceus</i> (C. L. Koch, 1839)* [Recent]	Qt Kerelia, Russia
<i>Eupelops</i> sp. in Karppinen & Koponen (1974)	Qt Finland
Peloptulus Berlese, 1908	Quaternary – Recent
225. <i>Peloptulus phaenotus</i> (C. L. Koch, 1844)* [Recent]	Qt Germany

UNDULORIBATIDAE Kunst, 1971	Palaeogene – Recent
Scutoribates Sellnick, 1918	Palaeogene – Recent
226. <i>Scutoribates perornatus</i> Sellnick, 1918	Pa Baltic amber
Unduloribates Balogh, 1943	?Palaeogene – Recent
227. <i>Unduloribates parvus</i> (Sellnick, 1931)	Pa Baltic amber
[generic affinities need clarification]	
ACHIPTERIOIDEA Thor, 1929	?Jurassic – Recent
ACHIPTERIIDAE Thor, 1929	?Jurassic – Recent
Achipteria Berlese, 1885	?Jurassic – Recent
228. <i>Achipteria coleoptera</i> (Linnaeus, 1757) [Recent]	Qt Finland / Greenland
229. ? <i>Achipteria obscura</i> Krivolutsky in Krivolutsky & Krasilov, 1977	J Russian far east
[An <i>incertae sedis</i> taxon?]	
Parachipteria van der Hammen, 1952	Quaternary – Recent
230. <i>Parachipteria punctata</i> (Nicolet, 1855) [Recent]	Qt northern Europe
231. <i>Parachipteria willmanni</i> van der Hammen, 1952 [Recent]	Qt Germany
EPACTOZETIDAE Grandjean, 1936b	Recent
no fossil record	
TEGORIBATIDAE Grandjean, 1954b	Quaternary – Recent
Tegoribates Ewing, 1917	Quaternary – Recent
232. <i>Tegoribates latirostris</i> (C. L. Koch, 1844) [Recent]	Qt Finland
ORIBATELLOIDEA Jacot, 1925	Palaeogene – Recent
ORIBATELLIDAE Jacot, 1925	Palaeogene – Recent
Oribatella Banks, 1895	Palaeogene – Recent
233. <i>Oribatella berlesei</i> (Michael, 1898) [Recent]	Qt Finland
234. <i>Oribatella calcarata</i> (C. L. Koch, 1835) [Recent]	Qt Kerelia, Russia
235. <i>Oribatella mirabilis</i> Sellnick, 1931	Pa Baltic amber
ORIPODOIDEA Jacot, 1925	Palaeogene – Recent
CALOPPIIDAE Balogh, 1960	Recent
= ?CRASSORIBATULIDAE author, date?	
no fossil record	
CAMPBELLLOBATIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
CHAUNOPROCTIDAE Balogh, 1961	Recent
no fossil record	

DRYMOBATIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
HAPLOZETIDAE Grandjean, 1936c	Palaeogene – Recent
= PROTORIBATIDAE J. Balogh & P. Balogh, 1984	
= XLOBATIDAE J. Balogh & P. Balogh, 1984	
Protoribates Berlese, 1908	Palaeogene – Recent
236. <i>Protoribates longipilis</i> Sellnick, 1931	Pa Baltic amber
LAMELLAREIDAE Balogh, 1972	Recent
no fossil record	
MAUDHEIMIIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
MOCHLOZETIDAE Grandjean, 1960a	Neogene – Recent
Mochlozetidae sp. <i>in</i> Norton & Poinar (1993)	Ne Dominican amber
<i>Mochloribatula</i> Mahunka, 1978	Neogene – Recent
237. <i>Mochloribatula smithi</i> (Woolley, 1971)	Ne Chiapas amber
<i>Mochlozetes</i> Grandjean, 1930	Neogene – Recent
<i>Mochlozetes</i> sp. <i>in</i> Norton & Poinar (1993)	Ne Dominican amber
NASOBATIDAE Balogh, 1972	Recent
no fossil record	
NEOTRICHZETIDAE Balogh, 1965	Recent
no fossil record	
NESOZETIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
ORIBATULIDAE Thor, 1929	Palaeogene – Recent
Oribatulidae sp. <i>in</i> Aoki (1974)	Qt Mizunami copal
<i>Lucoppia</i> Berlese, 1908	Palaeogene – Recent
238. <i>Lucoppia simplex</i> Sellnick, 1919	Pa Baltic amber
<i>Oribatula</i> Berlese, 1895	Quaternary – Recent
239. <i>Oribatula tibialis</i> (Nicolet, 1855)* [Recent]	Qt Europe
<i>Phauloppia</i> Berlese, 1908	Palaeogene – Recent
240. <i>Phauloppia lucorum</i> (C. L. Koch, 1841) [Recent]	Qt northern Europe
241. <i>Phauloppia pellucida</i> (Sellnick, 1931)	Pa Baltic amber
† <i>Sachalinella</i> Rjabinin <i>in</i> Krivolutzkii & Rjabinin, 1976	Palaeogene – Recent
May be a homonym of a bivalve genus	
242. <i>Sachalinella zherichini</i> Rjabinin <i>in</i> Krivolutzkii & Rjabinin, 1976*	Pa Sachalin amber

Zygoribatula Berlese, 1916	Quaternary – Recent
243. <i>Zygoribatula exilis</i> (Nicolet, 1855) [Recent]	Qt northern Europe
ORIPODIDAE Jacot, 1925	Palaeogene – Recent
= BIROBATIDAE J. Balogh & P. Balogh, 1984	
Benoibates Balogh, 1958	Neogene – Recent
244. <i>Benoibates chiapasensis</i> (Woolley, 1971)	Ne Chiapas amber
Oripoda Banks, 1904	Palaeogene – Recent
245. <i>Oripoda baltica</i> Sellnick, 1931	Pa Baltic amber
<i>Oripoda</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
Parapirnodus Balogh & Mahunka, 1968	Neogene – Recent
246. <i>Parapirnodus denaius</i> (Woolley, 1971)	Ne Chiapas amber
PARAKALUMMIDAE Grandjean, 1936b	Palaeogene – Recent
Neoribates Berlese, 1914	Palaeogene – Recent
247. <i>Neoribates borussicus</i> Sellnick, 1931	Pa Baltic amber
SCHELORIBATIDAE Grandjean, 1933	Palaeogene – Recent
Liebstadia Oudemans, 1906	Palaeogene – Recent
248. <i>Liebstadia similiformis</i> Sellnick, 1931	Pa Baltic amber
249. <i>Liebstadia similis</i> (Michael, 1888)* [Recent]	Qt Europe / Greenland
Scheloribates Berlese, 1908	Palaeogene – Recent
250. <i>Scheloribates apterus</i> Sellnick, 1931	Pa Baltic amber
251. <i>Scheloribates areatus</i> Sellnick, 1931	Pa Baltic amber
252. <i>Scheloribates durhami</i> (Woolley, 1971)	Ne Chiapas amber
253. <i>Scheloribates initialis</i> (Berlese, 1908) [Recent]	Qt Europe
254. <i>Scheloribates laevigatus</i> (C. L. Koch, 1835) [Recent]	Qt northern Europe
255. <i>Scheloribates latipes</i> (C. L. Koch, 1844) [Recent]	Qt Europe
256. <i>Scheloribates pallidulus</i> (C. L. Koch, 1841) [Recent]	Qt Germany
257. <i>Scheloribates setatus</i> Sellnick, 1931	Pa Baltic amber
SELLNICKIIDAE Balogh & Balogh, 1984	Recent
no fossil record	
STELECHOBATIDAE Grandjean, 1965b	Recent
no fossil record	
SYMBIORIBATIDAE Aoki, 1966b	Recent
no fossil record	
TUBULOZETIDAE Balogh, 1989	Quaternary – Recent
<i>Grandjeanobates</i> Ramsay, 1967	Quaternary – Recent

? <i>Grandjeanobates</i> sp.	Qt New Zealand
ZETOMOTRICHIDAE Grandjean, 1954b	Paleogene – Recent
Zetomotrichidae sp. <i>in</i> Sidorchuk & Norton (2011)	P Baltic amber
CERATOZETOIDEA Jacot, 1925	Paleogene – Recent
CERATOKALUMMIDAE Balogh, 1970	Recent
no fossil record		
CERATOZETIDAE Jacot, 1925	Paleogene – Recent
Ceratozetes Berlese, 1908	Quaternary – Recent
258. <i>Ceratozetes gracilis</i> (Michael, 1884)* [Recent]	Qt Finland
259. <i>Ceratozetes minimus</i> Sellnick, 1928 [Recent]	Qt Germany
260. <i>Ceratozetes parvulus</i> Sellnick, 1922 [Recent]	Qt Germany
Diapterobates Grandjean, 1936b	Quaternary – Recent
261. <i>Diapterobates notatus</i> (Thorell, 1871) [Recent]	Qt Europe / Greenland
Edwardzetes Berlese, 1914	Quaternary – Recent
262. <i>Edwardzetes edwardsi</i> (Nicolet, 1855)* [Recent]	Qt western Norway
Fuscozetes Sellnick, 1928	Quaternary – Recent
263. <i>Fuscozetes fuscipes</i> (C. L. Koch, 1844)* [Recent]	Qt western Norway
Melanozetes Hull, 1916	Paleogene – Recent
264. <i>Melanozetes federatus</i> Sellnick, 1931	Pa Baltic amber
265. <i>Melanozetes mollicomnus</i> [Recent] <i>fossilis</i> Sellnick, 1931	Pa Baltic amber
266. <i>Melanozetes meridianus</i> Sellnick, 1928 [Recent]	Qt Greenland
<i>Melanozetes</i> sp. <i>in</i> Karppinen et al. (1979)	Qt Karelia, Russia
Oromucia Thor, 1930	Quaternary – Recent
267. <i>Oromucia bicuspidata</i> Thor, 1930* [Recent]	Qt western Norway
268. <i>Oromucia lucens</i> (C. L. Koch, date?) [Recent]	Qt Greenland
Sphaerozetes Berlese, 1885	Paleogene – Recent
269. <i>Sphaerozetes convexulus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
270. <i>Sphaerozetes pirifomis</i> (Nicolet, 1855) [Recent]	Qt Finland
271. <i>Sphaerozetes primus</i> Sellnick, 1931	Pa Baltic amber
Trichoribates Berlese, 1910	Quaternary – Recent
272. <i>Trichoribates biarea</i> Gjelstrup & Solhøy, 1994 [Recent]	Qt western Norway
273. <i>Trichoribates incisellus</i> (Kramer, 1897) [Recent]	Qt Europe
274. <i>Trichoribates monticola</i> (Trägårdh, 1902) [Recent]	Qt western Norway
275. <i>Trichoribates setiger</i> (Trägårdh, 1910) [Recent]	Qt western Norway
276. <i>Trichoribates trimaculatus</i> (C. L. Koch, 1835)* [Recent]	Qt northern Europe
CHAMOBATIDAE Thor, 1937	Paleogene – Recent
Chamobates Hull, 1916	Paleogene – Recent

277. *Chamobates borealis* (Trägårdh, 1902) [Recent] Qt western Norway
278. *Chamobates cuspidatus* (Michael, 1884) [Recent] Qt Finland
279. *Chamobates difficilis* Sellnick, 1931 Pa Baltic amber
- EUZETIDAE Grandjean, 1954b** Quaternary – Recent
- Euzetes* Berlese, 1908 Quaternary – Recent
280. *Euzetes globulus* (Nicolet, 1855) [Recent] Qt Finland
- HUMEROBATIDAE Grandjean, 1970** Recent
- no fossil record
- MYCOBATIDAE Grandjean, 1954b** Quaternary – Recent
- Mycobates* Hull, 1916 Quaternary – Recent
281. *Mycobates consimilis* Hammer, 1952 [Recent] Qt Greenland
282. *Mycobates parmeliae* (Michael, 1884) [Recent] Qt Karelia, Russia
283. *Mycobates sarekenis* (Trägårdh, 1910) [Recent] Qt western Norway
- Puncoribates* Berlese, 1908 Quaternary – Recent
284. *Puncoribates punctum* (C. L. Koch, 1839) [Recent] Qt Karelia, Russia
285. *Puncoribates sellnicki* Willmann, 1928 [Recent] Qt Europe
- Puncoribates* sp. in Karppinen & Koponen (1973) Qt Finland
- ONYCHOBATIDAE Luxton, 1985** Recent
- no fossil record
- RAMSAYELLIDAE Luxton, 1985** Recent
- no fossil record
- ZETOMIMIDAE Shaldybina, 1966** Quaternary – Recent
- Zetomimus* author, date? Quaternary – Recent
286. *Zetomimus furcatus* (Pearce & Warburton, 1906)* [Recent] Qt Karelia, Russia
- GALUMNOIDEA Jacot, 1925** Palaeogene – Recent
- GALUMNELLIDAE Piffl, 1970** Quaternary – Recent
- Galumnella* Berlese, 1917 Quaternary – Recent
- Galumnella* sp. in Aoki (1974) Qt Mizunami copal
- GALUMNIDAE Jacot, 1925** Palaeogene – Recent
- Galumnidae spp. in Norton & Poinar (1993) Pa Baltic amber
- Acrogalumna Grandjean, 1956b** Quaternary – Recent
287. *Acrogalumna longipluma* (Berlese, 1904)* [Recent] Qt Karelia, Russia
- Galumna von Heyden, 1826** Palaeogene – Recent
288. *Galumna clavata* Sellnick, 1931 Pa Baltic amber

289. <i>Galumna diversa</i> Sellnick, 1931	Pa	Baltic amber
290. <i>Galumna lanceata</i> (Oudemans, 1900) [Recent]	Qt	Karelia, Russia
291. <i>Galumna obvia</i> (Berlese, 1915) [Recent]	Qt	Finland
<i>Galumna</i> sp. in Karppinen & Koponen (1974)	Qt	Finland
<i>Pergalumna</i> Grandjean, 1936b		Quaternary – Recent
292. <i>Pergalumna dorsalis</i> (C. L. Koch, 1835) [Recent]	Qt	Finland
293. <i>Pergalumna nervosa</i> (Berlese, 1914)* [Recent]	Qt	northern Europe
<i>Pilogalumna</i> Grandjean, 1956b		Quaternary – Recent
294. <i>Pilogalumna tenuiclava</i> (Berlese, 1908) [Recent]	Qt	Germany
ASTIGMATA G. Canestrini, 1891 (cohort)		Palaeogene – Recent
= ACARIDIDA author, date?		
SCHIZOGLYPHOIDEA Mahunka, 1978		Recent
SCHIZOGLYPHIDAE Mahunka, 1978		Recent
no fossil record		
HISTIOSTOMATOIDEA Berlese, 1897		?Palaeogene – Recent
GUANOLICHIDAE Fain, 1968		Recent
no fossil record		
HISTIOSTOMATIDAE Berlese, 1897		?Palaeogene – Recent
Hististomatidae? [alternatively Acaridae] in Dunlop et al. (2012)	Pa	Baltic amber
CANESTRINIOIDEA Berlese, 1884		Recent
CANESTRINIIDAE Berlese, 1884		Recent
no fossil record		
CHETOCHELACARIDAE Fain, 1987		Recent
no fossil record		
HETEROCHOPTIDAE Fain, 1967b		Recent
no fossil record		
LEMANNIELLIIDAE Wurst, 2001		Recent
no fossil record		
Superfamily?		
[NB: Sidorchuk & Klimov (2011) discussed the problems in placing this extinct family.]		
† GLAESACARIDAE Klimov & Sidorchuk in Sidorchuk & Klimov, 2011		Palaeogene
† <i>Glaesacarus</i> Klimov & Sidorchuk in Sidorchuk & Klimov, 2011		Palaeogene – Recent
295. <i>Glaesacarus rhombeus</i> (C. L. Koch & Berendt, 1854)*	Pa	Baltic amber

HEMISCARPOCTOIDEA Oudemans, 1908	Neogene – Recent
ALGOPHAGIDAE Fain, 1974	Recent
no fossil record	
CARPOGLYPHIDAE Oudemans, 1923	Recent
no fossil record	
CHAETODACTYLIDAE Zachvatkin, 1941	Recent
no fossil record	
HEMISARCOPTIDAE Oudemans, 1908	Recent
no fossil record	
HYADESIIDAE Halbert, 1915	Recent
no fossil record	
MELIPONOCOPTIDAE Fain & Rosa, 1983	Recent
no fossil record	
WINTERSCHMIDTIIDAE Oudemans, 1923	Neogene – Recent
† Amphicalvolia Türk, 1963	Neogene – Recent
296. <i>Amphicalvolia hurdi</i> Türk, 1963*	Ne Chiapas amber
GLYCOPHAGOIDEA Berlese, 1897	Recent
AEROGLYPHIDAE Zachvatkin, 1941	Recent
no fossil record	
CHORTOGLYPHIDAE Berlese, 1897	Recent
no fossil record	
ECHIMYOPODIDAE Fain, 1967a	Recent
no fossil record	
EUGLYCYPHAGIDAE Fain & Phillips, 1977	Recent
no fossil record	
GLYCYPHAGIDAE Berlese, 1897	Recent
no fossil record	
PEDETOPODIDAE Fain, 1969	Recent
no fossil record	
ROSENSTEINIIDAE Coorman, 1954	Recent

= LOPHONOTACARIDAE Fain, 1987	
= TROGLOTACARIDAE Fain, 1977	
no fossil record	
ACAROIDEA Latreille, 1802	Neogene – Recent
ACARIDAE Latreille, 1802	Recent
[query family placement?]	
† Tyroglyphites Pampaloni, 1902	Neogene – Recent
297. <i>Tyroglyphites miocenicus</i> Pampaloni, 1902*	Ne Sicily
GAUDIELLIDAE Atyeo et al., 1974	Recent
= PARTAMONACOPTIDAE author, date?	
= PLATYGLYPHIDAE Kurosa, 1976	
no fossil record	
GLYCACARIDAE Griffiths, 1977	Recent
no fossil record	
LARDOGLYPHIDAE Oudemans, 1877	Recent
no fossil record	
SAPRACARIDAE Fain, 1988	Recent
no fossil record	
SCATOGLYPHIDAE Zachvatkin & Volgin, 1956	Recent
no fossil record	
SUIDASIIDAE Hughes, 1948	Recent
no fossil record	
TYROGLYPHIDAE Donnadieu, 1868	Quaternary – Recent
Tyroglyphidae sp. in Aoki (1974)	Qt Mizunami copal
HYPODERATOIDEA Murray, 1877	Recent
HYPODERATIDAE Murray, 1877	Recent
no fossil record	
PSOROPTIDIA Yunker, 1955 (unranked clade)	Neogene – Recent
PTEROLICHOIDEA Trouessart & Mégnin, 1884	Recent
= FREYANOIDEA Dubinin, 1953	
ASCOURACARIDAE Gaud & Atyeo, 1976	Recent
no fossil record	

- CAUDIFERIDAE** Gaud & Atyeo, 1978 Recent
no fossil record
- CHEYLABIDIDAE** Gaud, 1983 Recent
no fossil record
- CRYPTUROPTIDAE** Gaud, Atyeo & Berla, 1972 Recent
no fossil record
- EUSTATHIIDAE** Oudemans, 1905 Recent
no fossil record
- FALCULIFERIDAE** Oudemans, 1905 Recent
no fossil record
- FREYANIDAE** Dubinin, 1953 Recent
no fossil record
- GABUCINIIDAE** Gaud & Atyeo, 1975 Recent
no fossil record
- KIWILICHIDAE** Dabert, 1994 Recent
no fossil record
- KRAMERELLIDAE** Gaud & Mouchet, 1961 Recent
no fossil record
- OCHROLICHIDAE** Gaud & Atyeo, 1978 Recent
no fossil record
- OCONNORIIDAE** Gaud, Atyeo & Klompen, 1989 Recent
no fossil record
- PTEROLICHIDAE** Trouessart & Mégnin, 1884 Recent
no fossil record
- PTILOXENIDAE** Gaud, 1982 Recent
no fossil record
- RECTIJANUIDAE** Gaud, 1961 Recent
no fossil record
- SYRINGOBIIDAE** Trouessart, 1897 Recent
no fossil record

THORACOSATHESIDAE Gaud & Mouchet, 1959	Recent
no fossil record	
VEXILLARIIDAE Gaud & Mouchet, 1959	Recent
no fossil record	
ANALGOIDEA Trouessart & Mégnin, 1884	Recent
ALLOPTIDAE Gaud, 1957	Recent
no fossil record	
ANALGIDAE Trouessart & Mégnin, 1884	Recent
no fossil record	
APIONACARIDAE Gaud & Atyeo, 1977	Recent
no fossil record	
AVENZOARIIDAE Oudemans, 1905	Recent
no fossil record	
CYTODITIDAE Oudemans, 1908	Recent
no fossil record	
DERMATIONIDAE Fain, 1965	Recent
no fossil record	
DERMOGLYPHIDAE Mégnin & Trouessart, 1884	Recent
no fossil record	
EPIDERMOPTIDAE Trouessart, 1892	Recent
no fossil record	
GAUDOGLYPHIDAE Bruce & Johnston, 1976	Recent
no fossil record	
HETEROPSORIDAE Oudemans, 1908	Recent
no fossil record	
KNEMIDOKOPTIDAE Dubinin, 1953	Recent
no fossil record	
LAMINOSIOPTIDAE Vitzthum, 1931	Recent
no fossil record	

PROCTOPHYLLODIDAE Mégnin & Trouessart, 1884	Recent
no fossil record	
PSORALGIDAE Oudemans, 1908	Recent
no fossil record	
PSOROPTOIDIDAE Gaud, 1983	Recent
no fossil record	
PTERONYSSIDAE Oudemans, 1941	Recent
no fossil record	
PTYSSALGIDAE Atyeo & Gaud, 1979	Recent
no fossil record	
PYROGLYPHIDAE Cunliffe, 1958	Recent
no fossil record	
TARSOCHEYLIDAE Atyeo & Gaud, 1979	Recent
no fossil record	
THYSANOCERCIDAE Atyeo & Peterson, 1972	Recent
no fossil record	
TROUESSARTIIDAE Gaud, 1957	Recent
no fossil record	
TURBINOPTIDAE Fain, 1957	Recent
no fossil record	
XOLALGIDAE Dubinin, 1953	Recent
no fossil record	
SARCOPTOIDEA Murray, 1877	Neogene–Recent
= PSOROPTIOIDEA Canestrini, 1892	
ACAROPTIDAE Womersley, 1953	Recent
no fossil record	
ATOPOMELIDAE Gunter, 1942	Neogene–Recent
?Apotomelidae sp. [originally as Listrophoridae in Poinar 1988]	Ne Dominican amber
AUDYCOPTIDAE Lavoipierre, 1964	Recent

no fossil record

CHIRODISCIDAE Trouessart, 1892 Recent

no fossil record

CHIRORHYNCHOBIIDAE Fain, 1967 Recent

no fossil record

GALAGALIDAE Fain, 1963 Recent

no fossil record

GASTRONYSSIDAE Fain, 1956 Recent

no fossil record

LEMURNYSIIDAE Fain, 1957 Recent

no fossil record

LISTROPHORIDAE Mégnin & Trouessart, 1884 Recent

no fossil record

LOBALGIDAE Fain, 1965 Recent

no fossil record

MYCOPTIDAE Gunther, 1942 Recent

no fossil record

PSOROPTIDAE Canestrini, 1892 Recent

no fossil record

PNEUMOCOPTIDAE Fain, 1957 Recent

no fossil record

RHYNCOPTIDAE Lawrence, 1956 Recent

no fossil record

SARCOPTIDAE Murray, 1877 Recent

no fossil record

NOMINA DUBIA

1. *Acarus resinosus* Presl, 1822 Pa Baltic amber
2. *Strieremaeus cordiformatus* Sellnick, 1919 [as species inquirenda] Pa Baltic amber

NOMINA NUDA

1. *Erythraeus hirsutissimus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

2. *Gymnodamaeus kulczynskii* Petrunkevitch, 1955a Pa Baltic amber
3. *Trombidium fossile* Keferstein, 1834 Pa Aix-en-Provence?

MISIDENTIFICATIONS

1. *Limnochares antiquus* Heyden, 1862 [larval hemipteran insect] Pa Rott, Germany

NON NAMES IN ZOOLOGY

Taxa assigned to living mite genera based on the fossil responses of plant tissue (galls); see discussion in Dunlop & Braddy (2011)

1. *Eriophyes daphnogene* Ambrus & Hably, 1979 [fossil gall] Pa Hungary
2. *Eryophyes [sic] vilarrubiae* Villalta, 1957 [fossil gall] Ne Spain
3. *Phytopus antiquus* van Heyden, 1860 [fossil gall] Ne Rott, Germany

c. 36,900 Recent species according to Hallan (2004)

RICINULEI

16 currently valid species of fossil ricinuleid

RICINULEI Thorell, 1876c Carbon. – Recent

= RHINOGASTRA Cook, 1899
= PODOGONA Cook, 1899

† PALAEORICINULEI Selden, 1992 (suborder) Carboniferous – ?Cret.

NB: Wunderlich (2012e) treated the two suborders as superfamilies.

Ricinulei indet. in Wunderlich (2012e) K Myanmar amber

† CURCULOIDIDAE Cockerell, 1916 Carboniferous

† Amarixys Selden, 1992 Carboniferous

1. *Amarixys gracilis* (Petrunkevitch, 1945a) C Mazon Creek
2. *Amarixys stellaris* Selden, 1992 C Mazon Creek
3. *Amarixys sulcata* (Melander, 1903)* C Mazon Creek

† Curculioides Buckland, 1837 Carboniferous

4. *Curculioides adompha* Brauckmann, 1987 C Hagen-Vorhalle
5. *Curculioides anstictii* Buckland, 1837* C Coalbrookdale
6. *Curculioides eltringhami* Petrunkevitch, 1949 C Crawcrook
7. *Curculioides gigas* Selden, 1992 C Mazon Creek
8. *Curculioides granulatus* Petrunkevitch, 1949 C Ilkeston
9. *Curculioides mcluckiei* Selden, 1992 C Mazon Creek
10. *Curculioides pococki* Selden, 1992 C Coseley
11. *Curculioides scaber* (Scudder, 1890b) C Mazon Creek

† POLIOCHERIDAE Scudder, 1884 Carboniferous – ?Cret.

† Poliochera Scudder, 1884 Carboniferous – ?Cret.

12. ?*Poliochera cretacea* Wunderlich, 2012e K Myanmar amber
13. *Poliochera gibbsi* Selden, 1992 C Illinois
14. *Poliochera glabra* Petrunkevitch, 1913 C Mazon Creek
15. *Poliochera punctulata* Scudder, 1884* C Mazon Creek

† Terpsicroton Selden, 1992 Carboniferous

16. *Terpsicroton alticeps* Selden, 1992* C Coseley

NEORICINULEI Selden, 1992 (suborder) Recent

RICINOIDIDAE Ewing, 1929 Recent

= CRYPTOSTEMMIDAE Westwood, 1874

no fossil record

NOMINA DUBIA

1. *Poliochera / Curculioides pustulatus* Laurentiaux-Viera & Laurentiaux, 1963 C Kiaping

55 Recent species according to Harvey (2003)

ARACHNIDA and/or PANTETRAPULMONATA

incertae sedis

3 currently valid, unplaced fossil arachnid and/or tetrapulmonate species

- all three species below have been suggested as possible members of the so-called pantetrapulmonate arachnids; i.e. spiders and their closest relatives

† *Ecchosis* Selden & Shear, 1991 Devonian

1. *Ecchosis pulchribothrium* Selden & Shear in Selden et al. 1991* D Gilboa

† *Saccogulus* Dunlop, Fayers, Hass & Kerp, 2006 Devonian

2. *Saccogulus seldeni* Dunlop, Fayers, Hass & Kerp, 2006* D Rhynie chert

† *Xenarachne* Dunlop & Poschmann, 1997 Devonian

3. *Xenarachne wilwerathensis* Dunlop & Poschmann, 1997* D Willwerath

no Recent species

TRIGONOTARBIDA

68 currently valid species of fossil trigonotarbid

- † **TRIGONOTARBIDA** Petrunkevitch, 1949 Silurian – Permian
- = ANTHRACOMARTI Karsch, 1882
 - = MERIDOGASTRA Thorell & Lindström, 1885
 - = EURYMARTI Matthew, 1895
- plesion genus**
- † **Palaeotarbus** Dunlop, 1999 Silurian
- = † *Eotarbus* Dunlop, 1996 [preoccupied]
 - 1. *Palaeotarbus jerami* (Dunlop, 1996)* S Ludford Lane
- † **PALAEOCHARINIDAE** Hirst, 1923 Devonian
- † **Aculeatarbus** Shear, Selden & Rolfe, 1987 Devonian
- 2. *Aculeatarbus depressus* Shear, Selden & Rolfe, 1987* D Gilboa
- † **Gelasinotarbus** Shear, Selden & Rolfe, 1987 Devonian
- 3. *Gelasinotarbus bifidus* Shear, Selden & Rolfe, 1987 D Gilboa
 - 4. *Gelasinotarbus bonamoae* Shear, Selden & Rolfe, 1987* D Gilboa
 - 5. *Gelasinotarbus heptops* Shear, Selden & Rolfe, 1987 D Gilboa
 - 6. *Gelasinotarbus reticulatus* Shear, Selden & Rolfe, 1987 D Gilboa
- † **Gigantocharinus** Shear, 2000 Devonian
- 7. *Gigantocharinus szatmaryi* Shear, 2000* D Red Hill, USA
- † **Gilboarachne** Shear, Selden & Rolfe, 1987 Devonian
- 8. *Gilboarachne griersoni* Shear, Selden & Rolfe, 1987* D Gilboa
- † **Palaeocharinus** Hirst, 1923 Devonian
- = † *Palaeocharinoides* Hirst, 1923
 - 9. *Palaeocharinus calmani* Hirst, 1923 D Rhynie cherts
 - 10. *Palaeocharinus hornei* (Hirst, 1923) D Rhynie cherts
 - 11. *Palaeocharinus kidstoni* Hirst, 1923 D Rhynie cherts
 - 12. *Palaeocharinus rhyniensis* Hirst, 1923* D Rhynie cherts
 - 13. *Palaeocharinus scourfieldi* Hirst, 1923 D Rhynie cherts
 - 14. *Palaeocharinus tuberculatus* Fayers, Dunlop & Trewin, 2005 D Rhynie cherts
- † **Spinocharinus** Poschmann & Dunlop, 2011 Devonian
- 15. *Spinocharinus steinmeyeri* Poschman & Dunlop, 2011* D Bürdenbach
- † **ARCAEOMARTIDAE** Poschmann & Dunlop, 2010 Devonian
- † **Archaeomartus** Størmer, 1970 Devonian
- 16. *Archaeomartus levis* Størmer, 1970* D Alken an der Mosel
 - i. = *Archaeomartus tuberculatus* Størmer, 1970 D Alken an der Mosel

- † ANTHRACOMARTIDAE Haase, 1890 Carboniferous
- = † PROMYGALIDAE Frič, 1904
 - = † BRACHYPYGIDAE Pocock, 1911
 - = † CORYPHOMARTIDAE Petrunkevitch, 1945
 - = † PLEOMARTIDAE Petrunkevitch, 1945
- † *Anthracomartus* Karsch, 1882 Carboniferous
- = † *Brachylycosa* Frič, 1904
 - = † *Cleptomartus* Petrunkevitch, 1949
 - = † *Coryphomartus* Petrunkevitch, 1945a
 - = † *Cryptomartus* Petrunkevitch, 1945a
 - = † *Oomartus* Petrunkevitch, 1953
 - = † *Perneria* Frič, 1904
 - = † *Pleomartus* Petrunkevitch, 1945a
 - = † *Promygale* Frič, 1901
17. *Anthracomartus bohemica* (Frič, 1901) C Nýřany
18. *Anthracomartus carcinoides* (Frič, 1901) C Nýřany
- i. = *Promygale rotundata* Frič, 1901 C Nýřany
 - ii. = *Perneria salticoides* Frič, 1904 C ?Nýřany
19. *Anthracomartus elegans* Frič, 1901 C Nýřany
20. *Anthracomartus hindii* Pocock, 1911 C Coseley
- i. = *Cleptomartus hangardi* Guthörl, 1965 C Saar, Germany
 - ii. = *Cryptomartus meyeri* Guthörl, 1964 C Aachen
 - iii. = *Cleptomartus planus* Petrunkevitch, 1949 C Coseley
 - iv. = *Cryptomartus rebskei* Brauckmann, 1984 C Saarbrücken
21. *Anthracomartus granulatus* Frič, 1904 C Nowa Ruda
22. *Anthracomartus janae* (Opluštil, 1986) C Kladno
23. *Anthracomartus kustae* Petrunkevitch, 1953 C Rakovník
24. *Anthracomartus minor* Kušta, 1884 C Rakovník
- i. = *Anthracomartus socius* Kušta, 1888 C Rakovník
25. *Anthracomartus nyranensis* (Petrunkevitch, 1953) C Nýřany
26. *Anthracomartus palatinus* Ammon, 1901 C Brücke, Germany
27. *Anthracomartus preisti* Pocock, 1911 C Coseley
- i. = *Anthracomartus denuiti* Pruvost, 1922 C Charleroi
 - ii. = *Cleptomartus plautus* Petrunkevitch, 1949 C Coseley
28. *Anthracomartus radvanicensis* (Opluštil, 1985) C Radvanice
29. *Anthracomartus triangularis* Petrunkevitch, 1913 C Joggins
30. *Anthracomartus trilobitus* Scudder, 1884 C Fayetteville
31. *Anthracomartus voelkelianus* Karsch, 1882* C Europe
- Anthracomartus* sp. in Wright & Selden (2011) C Kansas
- † *Brachypyge* Woodward, 1878b Carboniferous
32. *Brachypyge carbonis* Woodward, 1878b* C Mons

- † *Maiocercus* Pocock, 1911 Carboniferous
33. *Maiocercus celticus* (Pocock, 1902)* C Coal Measures
 i. = *Maiocercus orbicularis* Gill, 1911 C Westhoughton
- † ANTHRACOSIRONIDAE Pocock, 1903a Devonian – Carbon.
- † *Anthracosiro* Pocock, 1903a Carboniferous
34. *Anthracosiro fritschii* Pocock, 1903b C Coseley
 i. = *Anthracosiro elongatus* Waterlot, 1934 C Marlebach, France
35. *Anthracosiro woodwardi* Pocock, 1903a* C Coal Measures
 i. = *Anthracosiro corsini* Pruvost, 1926 C Noeux, France
 ii. = *Anthracosiro latipes* Gill, 1909 C Ryton-on-Tyne, UK
- † *Arianrhoda* Dunlop & Selden, 2004 Devonian
36. *Arianrhoda bennetti* Dunlop & Selden, 2004* D Tredomen
- † *Vratislavia* Frič, 1904 Carboniferous
37. *Vratislavia silesica* (Roemer, 1878)* C Silesia
- † TRIGONOTARBIDAE Petrunkevitch, 1949 Devonian – Carbon.
- † *Trigonotarbus* Pocock, 1911 Devonian – Carbon.
38. *Trigonotarbus arnoldi* Petrunkevitch, 1955b C Decazeville
 39. *Trigonotarbus johnsoni* Pocock, 1911* C Coseley
 40. *Trigonotarbus stoermeri* Schultka, 1991 D Rheinischen Schiefer.
- Family uncertain**
- † *Namurotarbus* Poschmann & Dunlop, 2010 Carboniferous
41. *Namurotarbus roessleri* (Dunlop & Brauckmann, 2006)* C Hagen-Vorhalle
- † *Tynecotarbus* Hradská & Dunlop, 2013 Carboniferous
42. *Tynecotarbus tichaveki* Hradská & Dunlop, 2013 C Týnec
- † *Permotarbus* Dunlop & Rößler, 2013 Permian
43. *Permotarbus schuberti* Dunlop & Rößler, 2013 P Chemnitz
- † LISSOMARTIDAE Dunlop, 1995 Carboniferous
- † *Lissomartus* Petrunkevitch, 1949 Carboniferous
44. *Lissomartus carbonarius* (Petrunkevitch, 1913) C Mazon Creek
 45. *Lissomartus schucherti* (Petrunkevitch, 1913)* C Mazon Creek
- † APHANTOMARTIDAE Petrunkevitch, 1945a Devonian – Permian
- = † TRIGONOMARTIDAE Petrunkevitch, 1949
- † *Alkenia* Størmer, 1970 Devonian
46. *Alkenia mirabilis* Størmer, 1970* D Alken an der Mosel
- † *Aphantomartus* Pocock, 1911 Carbon. – Permian
- = † *Trigonomartus* Petrunkevitch, 1913
 = † *Phrynomartus* Petrunkevitch, 1945a

47. *Aphantomartus areolatus* Pocock, 1911* C-P Coal Measures
- i. = *Aphantomartus pococki* Pruvost, 1912 C Anzin, France
 - ii. = *Trigonomartus dorlodotii* Pruvost, 1930 C Rien, France
 - iii. = *Eophryrus waechteri* Guthörl, 1938 C Saar
 - iv. = ?*Trigonomartus pruvosti* van der Heide, 1951 C Limbourg
 - v. = ?*Brachylycosa manebachensis* Müller, 1957 C Rotliegenden
48. *Aphantomartus ilfeldicus* (Scharf, 1924) P Rotliegend
49. *Aphantomartus pustulatus* (Scudder, 1884) C Coal Measures
- i. = ?*Kreischeria villeti* Pruvost, 1912 C Pas de Calais
 - ii. = *Cleptomartus plötzensis* Simon, 1971 C Halleschen Mulde
- † **KREISCHERIIDAE Haase, 1890** Carboniferous
- † **Anzinia Petrunkevitch, 1953** Carboniferous
50. *Anzinia thevenini* (Pruvost, 1919)* C Anzin
- † **Gondwanarache Pinto & Hünicken, 1980** Carboniferous
51. *Gondwanarache argentinensis* Pinto & Hünicken, 1980* C Bajo de Vélez
- † **Hemikreischeria Frič, 1904** Carboniferous
52. *Hemikreischeria geinitzi* (Thevenin, 1902)* C France
- † **Kreischeria Geinitz, 1882** Carboniferous
53. *Kreischeria wiedei* Geinitz, 1882* C Zwickau
- † **Pseudokreischeria Petrunkevitch, 1953** Carboniferous
54. *Pseudokreischeria pococki* (Gill, 1924) C Crawcrook
- i. = *Eophryrus varius* Petrunkevitch, 1949 C Crawcrook
- † **EOPHRYNIDAE Karsch, 1882** Carboniferous
- = † **HEMIPHRYNIDAE Frič, 1904**
- † **Eophrynus Woodward, 1871b** Carboniferous
55. *Eophrynus prestvicii* (Buckland, 1837)* C Coalbrookdale
56. *Eophrynus udus* Brauckmann, Koch & Kemper, 1985 C Hagen-Vorhalle
- † **Nyranytarbus Harvey & Selden, 1995** Carboniferous
- = † *Hemiphrynus* Frič, 1901 [preoccupied]
57. *Nyranytarbus hofmanni* (Frič, 1901) C Nýřany
58. *Nyranytarbus longipes* (Frič, 1901)* C Nýřany
- † **Petrovicia Frič, 1904** Carboniferous
59. *Petrovicia proditoria* Frič, 1904* C Petrovice
- † **Planomartus Petrunkevitch, 1953** Carboniferous
60. *Planomartus krejci* (Kušta, 1883)* C Rakovník
- i. = *Anthracomartus affinis* Kušta, 1885 C Rakovník
- † **Pleophrynus Petrunkevitch, 1945a** Carboniferous
61. *Pleophrynus verrucosus* (Pocock, 1911) C Coal Measures
- i. = *Eophrynus warei* Dix & Pringle, 1930 C Glyncoch, UK
 - ii. = *Pleophrynus ensifer* Petrunkevitch, 1945a* C Mazon Creek

- iii. = *Eophrynus jugatus* Ambrose & Romano, 1972 C Kilmersdon, UK
62. *Pleophrynus hawsei* Dunlop, Wang, Selden & Krautz, 2014 C Kinney Brick Quarry
- † **Pocononia Petrunkevitch, 1953** Carboniferous
63. *Pocononia whitei* (Ewing, 1930)* C Pocono Shales
- † **Somaspidion Jux, 1982** Carboniferous
64. *Somaspidion hammapheron* Jux, 1982* C Dinslaken
- † **Stenotrogulus Frič, 1904** Carboniferous
- = † *Cyclotrogulus* Frič, 1904
- = † *Pseudoeophrynus* Příbyl, 1958
65. *Stenotrogulus salmii* (Stur, 1877)* C Ostrava
- i. = *Cyclotrogulus sturii* Frič, 1904 [non Hasse, 1890] C Ostrava
- ii. = *Pseudoeophrynus ostraviensis* Příbyl, 1958 C Ostrava
- TRIGONOTARBIDA *incertae sedis*
- † **Anthracophrynus André, 1913** Carboniferous
66. *Anthracophrynus tuberculatus* André, 1913* C Dudweiler
- † **Areomartus Petrunkevitch, 1913** Carboniferous
67. *Areomartus ovatus* Petrunkevitch, 1913* C West Virginia
- † ‘**Eophrynus**’
68. ‘*Eophrynus*’ *scharfi* Scharf, 1924 P Rotliegend

NOMINA DUBIA

1. *Anthracomartus buchi* (Goldenberg, 1873) C Saarbrücken
2. *Anthracomartus hageni* (Goldenberg, 1873) C Saarbrücken
3. *Elaverimartus pococki* Petrunkevitch, 1953 C Ellismuir
 - i. = *Palaeophalangium Scoticum* Peach in Murdoch, 1893 [*nomen nudum*] C Ellismuir
4. *Eurymartus latus* Matthew, 1895 C Fern Ledges
5. ?*Eurymartus spinulosus* Matthew, 1895 C Fern Ledges
6. *Trigonomartus woodruffi* (Scudder, 1893) C Rhode Island

no Recent species

URARANEIDA

2 currently valid species of uraraneid

- The uraraneids were previously interpreted as true spiders (Araneae), but are now thought to be a more basal lineage which produced silk but lacked spinnerets.

† URARANEIDA Selden & Shear *in Selden et al., 2008* Devonian – Permian

FAMILY UNCERTAIN

† Attercopus Selden & Shear *in Selden et al. (1991)* Devonian

1. *Attercopus fimbriunguis* (Shear, Selden & Rolfe, 1987)* D Gilboa, New York

† PERMARACHNIDAE Eskov & Selden, 2005 Permian

† Permarachne Eskov & Selden, 2005 Permian

2. *Permarachne novokshonovi* Eskov & Selden, 2005* P Matveyevka

ARANEAE

1,192 currently valid species of fossil spider

ARANEAE Clerck, 1757	Carbon. – Recent
‘mesotheles’	Carbon. – Recent
† ARTHROLYCOSIDAE Frič, 1904	Carboniferous
† <i>Arthrolycosa</i> Harger, 1874	Carbon. – Permian
1. <i>Arthrolycosa antiqua</i> Harger, 1874*	C Mazon Creek
2. <i>Arthrolycosa danielsi</i> Petrunkevitch, 1913	C Mazon Creek
<i>Arthrolycosa</i> sp. <i>in Eskov & Selden (2005)</i>	P Kityak river
<i>Arthrolycosa</i> sp. <i>in Selden et al. (2014)</i>	C Chunya, Russia
<i>Arthrolycosa</i> sp. <i>in Selden et al. (2014)</i>	C Donets Basin
† <i>Eocteniza</i> Pocock, 1911	Carboniferous
3. <i>Eocteniza silvicola</i> Pocock, 1911*	C Coseley
† ARTHROMYGALIDAE Petrunkevitch, 1923	Carboniferous
† <i>Arthromygale</i> Petrunkevitch, 1923	Carboniferous
4. <i>Arthromygale fortis</i> (Frič, 1904)*	C Rakovník
i. = <i>Arthrolycosa beecheri</i> Frič, 1904	C Rakovník
† <i>Eolycosa</i> Kušta, 1885	Carboniferous
5. <i>Eolycosa lorenzi</i> Kušta, 1885*	C Rakovník
† <i>Geralycosa</i> Kušta, 1888	Carboniferous
6. <i>Geralycosa fritschi</i> Kušta, 1888*	C Rakovník
† <i>Kustaria</i> Petrunkevitch, 1953	Carboniferous
= † <i>Scudderia</i> Kušta, 1888 [preoccupied]	
7. <i>Kustaria carbonaria</i> (Kušta, 1888)*	C Rakovník
† <i>Palaranea</i> Frič, 1873	Carboniferous
8. <i>Palaranea borassifoliae</i> Frič, 1873*	C Czech Republic
† <i>Protocteniza</i> Petrunkevitch, 1949	Carboniferous
9. <i>Protocteniza britannica</i> Petrunkevitch, 1949*	C Coseley
† <i>Protolycosa</i> Roemer, 1866	Carboniferous
10. <i>Protolycosa anthracophilia</i> Roemer, 1866*	C Silesia
11. <i>Protolycosa cebennensis</i> Laurentiaux-Viera & Laurentiaux, 1963	C Cévennes, France
† <i>Rakovnicia</i> Kušta, 1884a	Carboniferous
12. <i>Rakovnicia antiqua</i> Kušta, 1884a*	C Rakovník
† PYRITARANEIDAE Petrunkevitch, 1953	Carboniferous

† <i>Dinopilio</i> Frič, 1904	Carboniferous
13. <i>Dinopilio gigas</i> Frič, 1904*	C Rakovník
14. <i>Dinopilio parvus</i> Petrunkevitch, 1953	C Kent, UK
† <i>Pyritaranea</i> Frič, 1901	Carboniferous
15. <i>Pyritaranea tubifera</i> Frič, 1901*	C Nýřany
MESOTHELAE Pocock, 1892	Carbon. – Recent
plesion genus	
† <i>Palaeothele</i> Selden, 2000	Carboniferous
= † <i>Eothele</i> Selden, 1996 [preoccupied]	
16. <i>Palaeothele montceauensis</i> (Selden, 1996)*	C Montceau-les-Mines
LIPHISTIIDAE Pocock, 1892	Recent
= HEPTATHELIDAE Haupt, 1983	
no fossil record	
OPISTHOTHELAE Pocock, 1892	Triassic – Recent
Opisthothelae <i>incertae sedis</i>	
† <i>Eoatypus</i> McCook, 1888	Palaeogene
17. <i>Eoatypus woodwardii</i> McCook, 1888*	Pa Isle of Wight
MYGALOMORPHAE Pocock, 1892	Triassic – Recent
Mygalomorpha indet. 1–3 <i>in</i> Wunderlich (2008d)	K Myanmar amber
ATYPOIDEA Thorell, 1870a	Triassic – Recent
† <i>Friularachne</i> Dalla Vecchia & Selden, 2013	Triassic
18. <i>Friularachne rigoi</i> Dalla Vecchia & Selden, 2013*	Tr Friuli, Italy
ATYPIDAE Thorell, 1870a	Cretaceous – Recent
= CALOMMATOIDAE Thorell, 1887	
† <i>Ambioriphagus</i> Eskov & Zonstein, 1990	Cretaceous
19. <i>Ambioriphagus ponomarenkoi</i> Eskov & Zonstein, 1990*	K Central Mongolia
† <i>Balticatypus</i> Wunderlich, 2011 <i>h</i>	Palaeogene
20. <i>Balticatypus beigeli</i> Wunderlich, 2011 <i>h</i>	Pa Baltic amber
21. <i>Balticatypus juvenis</i> Wunderlich, 2011 <i>h</i> *	Pa Baltic amber
22. <i>Balticatypus spinosus</i> Wunderlich, 2011 <i>h</i>	Pa Baltic amber
ANTRODIAETIDAE Gertsch <i>in</i> Comstock, 1940	Cretaceous – Recent
= BRACHYBOTHRIDAE Simon, 1892	
= ACCATYMIDAE Kishida, 1930	
† <i>Cretacattyma</i> Eskov & Zonstein, 1990	Cretaceous
23. <i>Cretacattyma raveni</i> Eskov & Zonstein, 1990*	K Central Mongolia

MECICOBOTHRIIDAE Holmberg, 1882	Cretaceous – Recent
	= HEXURIDAE Simon, 1889b	
† <i>Cretohexura</i> Eskov & Zonstein, 1990	Cretaceous
24. <i>Cretohexura coylei</i> Eskov & Zonstein, 1990*	K Transbaikalia
† <i>Cretomegahexura</i> Eskov & Zonstein, 1990	Cretaceous
25. <i>Cretomegahexura platnicki</i> Eskov & Zonstein, 1990*	K Central Mongolia
HEXATHELIDAE Simon, 1892b	Triassic – Recent
† <i>Rosamygale</i> Selden & Gall, 1992	Triassic
26. <i>Rosamygale grauvogeli</i> Selden & Gall, 1992*	Tr Vosges, France
DIPLURIDAE Simon, 1889b	Cretaceous – Recent
† <i>Clostes</i> Menge, 1869	Palaeogene
27. <i>Clostes priscus</i> Menge, 1869*	Pa Baltic / Bitt. amber
† <i>Cretadiplura</i> Selden in Selden et al., 2006	Cretaceous
28. <i>Cretadiplura ceara</i> Selden in Selden et al., 2006*	K Crato Formation
† <i>Dinodiplura</i> Selden in Selden et al., 2006	Cretaceous
29. <i>Dinodiplura ambulacra</i> Selden in Selden et al., 2006*	K Crato Formation
<i>Ischnothele</i> Ausserer, 1875	?Neogene – Recent
? <i>Ischnothele</i> sp. in Wunderlich (1988)	Ne Dominican amber
<i>Masteria</i> L. Koch, 1873	Neogene – Recent
	= † <i>Microsteria</i> Wunderlich, 1988	
30. <i>Masteria sexoculata</i> (Wunderlich, 1988)	Ne Dominican amber
? <i>Masteria</i> sp. in Schawaller (1982c: as ? <i>Ischnothele</i>)	Ne Dominican amber
genus uncertain		
Dipluridae sp. 1–3 in Wunderlich (2004a)	Pa Baltic amber
Dipluridae sp. in Wunderlich (2004a)	Ne Dominican amber
Dipluridae indet. in Wunderlich (2012d)	K Myanmar amber
CYRTAUCHENIIDAE Simon, 1892b	Neogene – Recent
<i>Bolostromus</i> Ausserer, 1875	Neogene – Recent
31. <i>Bolostromus destructus</i> Wunderlich, 1988	Ne Dominican amber
CTENIZIDAE Thorell, 1887	Palaeogene – Recent
	= HALONOPROCTIDAE Pocock, 1903	
† <i>Baltocteniza</i> Eskov & Zonstein, 2000	Palaeogene
32. <i>Baltocteniza kulickae</i> Eskov & Zonstein, 2000	Pa Baltic amber
† <i>Electrocteniza</i> Eskov & Zonstein, 2000	Palaeogene
33. <i>Electrocteniza sadilenkoi</i> Eskov & Zonstein, 2000	Pa Baltic amber
<i>Ummidia</i> Thorell, 1875	Palaeogene – Recent
34. <i>Ummidia damzeni</i> Wunderlich, 2000	Pa Baltic amber

35. <i>Ummidia malinowskii</i> Wunderlich, 2000	Pa	Baltic amber
<i>Ummidia</i> sp. in Wunderlich (2004a)	Pa	Baltic amber
? <i>Ummidia</i> sp. in Wunderlich (2011h)	Pa	Baltic amber
EUCTENIZIDAE Raven, 1985		Recent
no fossil record		
IDIOPIDAE Simon, 1892b		Recent
no fossil record		
ACTINOPODIDAE Simon, 1892b		Recent
= ERIODONTIDAE C. L. Koch & Berendt, 1854		
[based on a generic synonym; listed in Bonnet as syn. of Clubionidae!]		
no fossil record		
MIGIDAE Simon, 1892b		Recent
no fossil record		
NEMESIIDAE Simon, 1892b		Cretaceous – Recent
= PYCNOTHELIDAE Chamberlin, 1917		
† <i>Cretamygale</i> Selden, 2002		Cretaceous
36. <i>Cretamygale chasei</i> Selden, 2002*	K	Isle of Wight
† <i>Eodiplurina</i> Petrunkevitch, 1922		Palaeogene
[NB: Selden (2001) questioned this familial placement based on claw structure]		
37. <i>Eodiplurina cockerelli</i> Petrunkevitch, 1922*	Pa	Florissant
MICROSTIGMATIDAE Roewer, 1942		Neogene – Recent
= MICROMYGALIDAE Wunderlich, 2004b		
† <i>Parvomygale</i> Wunderlich, 2004b		Neogene
38. <i>Parvomygale distincta</i> Wunderlich, 2004b*	Ne	Dominican amber
BARYCHELIDAE Simon, 1889b		Neogene – Recent
<i>Psalistops</i> Simon, 1889b		Neogene – Recent
39. <i>Psalistops hispaniolensis</i> Wunderlich, 1988*	Ne	Dominican amber
THERAPHOSIDAE Thorell, 1870a		Neogene – Recent
= AVICULARIIDAE Simon, 1874		
Theraphosidae gen. et sp. indet. in Dunlop et al. (2008)	Ne	Chiapas amber
<i>Hemirraghus</i> Simon, 1903		Neogene – Recent
<i>Hemirraghus</i> sp. in García-Villafuerte (2008)	Ne	Chiapas amber
† <i>Ischnocolinopsis</i> Wunderlich, 1988		Neogene
40. <i>Ischnocolinopsis acutus</i> Wunderlich, 1988*	Ne	Dominican amber

PARATROPIDIDAE Simon, 1889a	Recent
no fossil record	
ARANEOMORPHAE Smith, 1902	Triassic – Recent
ARANEOMORPHAE indet.	
† <i>Argyrarachne</i> Selden <i>in</i> Selden et al., 1999	Triassic
41. <i>Argyrarachne solitus</i> Selden <i>in</i> Selden et al., 1999*	Tr Virginia
† <i>Triassaraneus</i> Selden <i>in</i> Selden et al., 1999	Triassic
42. <i>Triassaraneus andersonorum</i> Selden <i>in</i> Selden et al., 1999*	Tr KwaZulu-Natal
HYPOCHILIDAE Marx, 1888	Recent
= ECTATOSTICTIDAE Lehtinen, 1967	
no fossil record	
AUSTROCHILOIDEA Zapfe, 1955	Recent
AUSTROCHILIDAE Zapfe, 1955	Recent
= THAIDIDAE Lehtinen, 1967	
= HICKMANIIDAE Lehtinen, 1967	
no fossil record	
GRADUNGULIDAE Forster, 1955	Recent
no fossil record	
ARANEOCLADA Platnick, 1977	Triassic – Recent
HAPLOGYNAE Simon, 1893	Jurassic – Recent
FILISTATIDAE Ausserer, 1867	Neogene – Recent
Misionella Ramírez & Grismado, 1997	Neogene – Recent
43. <i>Misionella didicostae</i> Penney, 2005a	Ne Dominican amber
SICARIIDAE Keyserling, 1880a	Neogene – Recent
= LOXOSCELIDAE Simon, 1893	
Loxosceles Heineken & Lowe, 1832	Neogene – Recent
44. <i>Loxosceles aculicaput</i> Wunderlich, 2004c	Ne Dominican amber
45. <i>Loxosceles defecta</i> Wunderlich, 1988	Ne Dominican amber
46. <i>Loxosceles deformis</i> Wunderlich, 1988	Ne Dominican amber
<i>Loxosceles</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
SCYTODIDAE Blackwall, 1864	Cretaceous – Recent
Syctodidae sp. 1–2 <i>in</i> Wunderlich (2004b)	Pa Bitterfeld amber
Scytodes Latreille, 1804a	?Cretaceous – Recent
47. ? <i>Scytodes hani</i> Wunderlich, 2012d	K Jordanian amber
48. <i>Scytodes marginalis</i> Wunderlich, 2004as	Qt Madagascan copal

49. *Scytodes piliformis* Wunderlich, 1988 Ne Dominican amber
50. *Scytodes planithorax* Wunderlich, 1988 Ne Dominican amber
51. *Scytodes stridulans* Wunderlich, 1988 Ne Dominican amber
52. *Scytodes weitschati* Wunderlich, 1993a Pa Baltic amber
- Scytodes sp. in Wunderlich (1988) Ne Dominican amber
- Scytodes sp. in Wunderlich (2011h) Pa Baltic amber
- PERIEGOPIDAE Simon, 1893** **Recent**
- no fossil record
- DRYMUSIDAE Simon, 1893** **Recent**
- no fossil record
- † **PRAETERLEPTONETIDAE** Wunderlich 2008d **Cretaceous**
- Praeterleptonetidae indet. in Wunderlich (2008d) K Myanmar amber
- † **Palaeohygropoda** Penney, 2004c **Cretaceous**
53. *Palaeohygropoda myanmarensis* Penney, 2004c* K Myanmar amber
- † **Praeterleptoneta** Wunderlich, 2008d **Cretaceous**
54. *Praeterleptoneta spinipes* Wunderlich, 2008d* K Myanmar amber
55. *Praeterleptoneta tibialis* Wunderlich, 2011i K Myanmar amber
- † **PHOLCOCHYROCERIDAE** Wunderlich, 2008d (n. stat. 2012d) **Cretaceous**
- † **Pholcochyrocer** Wunderlich, 2008d **Cretaceous**
56. ?*Pholcochyrocer baculum* Wunderlich, 2012d K Myanmar amber
57. *Pholcochyrocer guttulaequeae* Wunderlich, 2008d* K Myanmar amber
58. *Pholcochyrocer pecten* Wunderlich, 2012d K Myanmar amber
- LEPTONETIDAE** Simon, 1890 **Cretaceous – Recent**
- † **Eoleptoneta** Wunderlich, 1991 **Palaeogene**
59. *Eoleptoneta curvata* Wunderlich, 2004c Pa Bitterfeld amber
60. *Eoleptoneta duocalcar* Wunderlich, 2004c Pa Baltic amber
61. *Eoleptoneta kutscheri* Wunderlich, 1991* Pa Bitterfeld amber
62. *Eoleptoneta multispinae* Wunderlich, 2011h Pa Baltic amber
63. *Eoleptoneta pseudoarticulata* Wunderlich, 2011h Pa Baltic amber
64. *Eoleptoneta similis* Wunderlich, 2004c Pa Baltic amber
- † **Oligoleptoneta** Wunderlich 2004c **Palaeogene**
65. *Oligoleptoneta altoculus* Wunderlich 2004c* Pa Baltic amber
66. *Oligoleptoneta cymbiospina* Wunderlich, 2011h Pa Baltic amber
- † **Palaeoleptoneta** Wunderlich 2012d **Cretaceous**
67. *Paleoleptoneta calcar* Wunderlich, 2012d* K Myanmar amber
- TELEMIDAE** Fage, 1913 **Palaeogene – Recent**

Telema Simon, 1882	Palaeogene – Recent
68. ? <i>Telema moritzi</i> Wunderlich, 2004c	Pa Baltic / Bitt. amber
OCHYROCERATIDAE Fage, 1912 s. l. [incl. PSILODERCINAE]	Cretaceous – Recent
= † EOPSILODERCIDAE Wunderlich, 2008d	
[NB: Wunderlich (2012d) recognised this as a junior synonym of a family Psilodercidae; Platnick does not recognise this family]	
?Eopsilodercidae indet. 1–3 <i>in</i> Wunderlich (2008d)	K Myanmar amber
† Arachnolithulus Wunderlich, 1988	Neogene
69. <i>Arachnolithulus longipes</i> Wunderlich, 2004c	Ne Dominican amber
70. <i>Arachnolithulus pygmaeus</i> Wunderlich, 1988*	Ne Dominican amber
? <i>Arachnolithulus</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
† Eopsiloderces Wunderlich, 2008d	Cretaceous
71. <i>Eopsiloderces loxosceloides</i> Wunderlich, 2008d	K Myanmar amber
† Furcembolus Wunderlich, 2008d	Cretaceous
72. <i>Furembolus andersoni</i> Wunderlich, 2008d	K Myanmar amber
Leclercera Deeleman-Reinhold, 1995	Cretaceous – Recent
73. <i>Leclercera longissipes</i> Wunderlich, 2012d	K Myanmar amber
74. <i>Leclercera spicula</i> Wunderlich, 2012d	K Myanmar amber
Psiloderces Simon, 1892	?Cretaceous – Recent
75. ? <i>Psiloderces filiformis</i> Wunderlich, 2012d	K Myanmar amber
PHOLCIDAE C. L. Koch, 1851	Palaeogene – Recent
Pholcidae sp. 1–2 <i>in</i> Wunderlich (2004b)	Pa Baltic amber
Pholcidae sp. <i>in</i> Wunderlich (2004au)	Pa Fu Shun amber
Coryssocnemis Simon, 1893	Neogene – Recent
76. ? <i>Coryssocnemis velteni</i> Wunderlich, 2004c	Ne Dominican amber
Leptopholcus Simon, 1893	Neogene
77. <i>Leptopholcus kiskeya</i> Huber & Wunderlich, 2006	Ne Dominican amber
Modisimus Simon, 1893	Neogene – Recent
78. <i>Modisimus calcar</i> Wunderlich, 1988	Ne Dominican amber
79. <i>Modisimus calcaroides</i> Wunderlich, 1988	Ne Dominican amber
80. <i>Modisimus crassifemoralis</i> Wunderlich, 1988	Ne Dominican amber
81. <i>Modisimus oculatus</i> Wunderlich, 1988	Ne Dominican amber
82. <i>Modisimus tuberosus</i> Wunderlich, 1988	Ne Dominican amber
<i>Modisimus</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
† Paraspermophora Wunderlich, 2004c	Palaeogene
83. <i>Paraspermophora bitterfeldensis</i> Wunderlich, 2004c	Pa Bitterfeld amber
84. <i>Paraspermophora perplexa</i> Wunderlich, 2004c*	Pa Baltic amber
<i>Paraspermophora</i> sp. <i>in</i> Wunderlich (2004c, 2011h)	Pa Baltic / Bitt. amber
Pholcophora Banks, 1896	Neogene – Recent
85. <i>Pholcophora brevipes</i> Wunderlich, 1988	Ne Dominican amber

86. *Pholcophora gracilis* Wunderlich, 1988 Ne Dominican amber
87. *Pholcophora longicornis* Wunderlich, 1988 Ne Dominican amber
- Quamtana Huber, 2003** Palaeogene – Recent
88. *Quamtana huberi* Penney, 2007a Pa Le Quesnoy amber
- † **Serratochorus** Wunderlich, 1988 Neogene
89. *Serratochorus pygmaeus* Wunderlich, 1988* Ne Dominican amber
- PLECTREURIDAE Simon, 1893** Jurassic – Recent
- † **Eoplectreurus** Selden & Huang, 2010 Jurassic
90. *Eoplectreurus gertschi* Selden & Huang, 2010* J Daohugou
- † **Montsecarachne** Selden, 2014a Cretaceous
91. *Montsecarachne amicorum* Selden, 2014a* K El Montsec
- NB: erroneously cited as *amicus* in the abstract.
- † **Palaeoplectreurus** Wunderlich, 2004c Palaeogene
92. *Palaeoplectreurus baltica* Wunderlich, 2004c* Pa Baltic amber
- Plectreurus** Simon, 1893 Neogene – Recent
93. *Plectreurus pittfieldi* Penney, 2009 Ne Dominican amber
- DIGUETIDAE F. O. P.-Cambridge, 1899** Recent
- no fossil record
- CAPONIIDAE Simon, 1890** Neogene – Recent
- = COLOPHONIDAE O. P.-Cambridge, 1874 [based on a generic homonym]
- Nops** MacLeay, 1839 Neogene – Recent
94. *Nops lobatus* Wunderlich, 1988 Ne Dominican amber
- i. = *Nops segmentatus* Wunderlich, 1988 Ne Dominican amber
- Nops* sp. in Wunderlich (1988) Ne Dominican amber
- TETRABLEMMIDAE O. P.-Cambridge, 1873** Cretaceous – Recent
- = PHAEDOMOIDAE Thorell, 1890 [based on a generic homonym]
- = PACULLIDAE Simon, 1894
- Tetrablemmidae gen. indet. in Wunderlich (2012d) K Myanmar amber
- † **Balticoblemma** Wunderlich, 2004c Palaeogene
95. *Balticoblemma unicirculum* Wunderlich, 2004c* Pa Baltic amber
- † **Eogamasomorpha** Wunderlich, 2008d Cretaceous
96. *Eogamasomorpha nubila* Wunderlich, 2008d* K Myanmar amber
- † **Eoscaphiella** Wunderlich, 2011i Cretaceous
97. *Eoscaphiella ohlhoffi* Wunderlich, 2011i* K Myanmar amber
- Monoblemma** Gertsch, 1941 Neogene
98. ?*Monoblemma spinosum* Wunderlich, 1988* Ne Dominican amber
- † **Saetosoma** Wunderlich, 2012d Cretaceous
99. *Saetosoma filiembolus* Wunderlich, 2012d* K Myanmar amber

TROGLORAPTORIDAE Griswold, Audisio & Ledford, 2012	Recent
no fossil record	
DYSDEROIDEA Bristowe, 1938	Cretaceous – Recent
?Dysderoidea s. l. indet 1–2 <i>in</i> Wunderlich (2008d)	K Myanmar amber
SEGESTRIIDAE Simon, 1893	Cretaceous – Recent
?Segestriidae indet <i>in</i> Wunderlich (2008d)	K Myanmar amber
Ariadna Audouin, 1826	Cretaceous – Recent
100. ? <i>Ariadna amissiocoli</i> Wunderlich, 2008d	K Jordanian amber
101. <i>Ariadna copalis</i> Wunderlich, 2008a	Qt ?Madagascan copal
102. <i>Ariadna defuncta</i> Wunderlich 2004c	Pa Bitterfeld amber
103. <i>Ariadna hintzei</i> Wunderlich, 2004as	Qt Madagascan copal
104. <i>Ariadna ovalis</i> Wunderlich, 2008a	Pa Baltic amber
105. <i>Ariadna parva</i> Wunderlich, 2008a	Pa Baltic amber
106. <i>Ariadna paucispinosa</i> Wunderlich, 1988	Ne Dominican amber
107. <i>Ariadna resinae</i> Hickman, 1957	Ne? Australian copal
? <i>Ariadna</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
† Lebansegestria Wunderlich 2008d	Cretaceous
108. <i>Lebansegestria azari</i> Wunderlich, 2008d*	K Lebanese amber
† Microsegestria Wunderlich & Milki, 2004	Cretaceous
109. <i>Microsegestria poinari</i> Wunderlich & Milki, 2004*	K Lebanese amber
† Palaeosegestria Penney, 2004a	Cretaceous
110. <i>Palaeosegestria lutzii</i> Penney, 2004a*	K New Jersey amber
Segestria Latreille, 1804a	Cretaceous – Recent
111. <i>Segestria cristata</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
112. <i>Segestria flexio</i> Wunderlich, 2004c	Pa Baltic amber
113. <i>Segestria mortalis</i> Wunderlich 2004c	Pa Baltic amber
114. <i>Segestria plicata</i> Petrunkevitch, 1950	Pa Baltic amber
115. <i>Segestria scudderii</i> Petrunkevitch, 1922	Pa Florissant
116. <i>Segestria secessa</i> Scudder, 1890a	Pa Florissant
117. <i>Segestria succinei</i> Berland, 1939	Pa Baltic amber
118. <i>Segestria tomentosa</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
i. = <i>Segestria plicata</i> Petrunkevitch, 1950 [provisional]	Pa Baltic amber
<i>Segestria</i> sp. <i>in</i> Penney (2002)	K New Jersey amber
<i>Segestria</i> sp. <i>in</i> Wunderlich (2004c)	Pa Baltic amber
<i>Segestria</i> sp. <i>in</i> Selden (2014b)	Pa Isle of Wight
† Vetsegestria Wunderlich, 2004c	Palaeogene
119. <i>Vetsegestria quinquespinosa</i> Wunderlich, 2004c*	Pa Baltic / Bitter. amber
DYSDERIDAE C. L. Koch, 1837	Palaeogene – Recent
† Dasumiana Wunderlich, 2004c	Palaeogene

120. *Dasumiana emicans* Wunderlich, 2004c* Pa Baltic amber
121. ?*Dasumiana subita* (Petrunkevitch, 1958) Pa Baltic amber
122. *Dasumiana valga* Wunderlich, 2004c Pa Baltic amber
- Dysdera Latreille, 1804** Palaeogene – Recent
123. *Dysdera dilatata* Zhang, Sun & Zhang, 1994 Ne Shanwang
- Harpactea Bristowe, 1939** Palaeogene – Recent
124. *Harpactea communis* Wunderlich, 2004c Pa Baltic amber
125. *Harpactea extincta* Petrunkevitch, 1950 Pa Baltic amber
126. *Harpactea hombergi* (Scopoli, 1763) [Recent] Qt England
127. *Harpactea longibulbus* Wunderlich, 2011h Pa Baltic amber
128. *Harpactea tersa* (C. L. Koch & Berendt, 1854) ... [provisional transfer] Pa Baltic amber
- Harpactea* sp. in Wunderlich (2011h) Pa Bitterfeld amber
- † **Segistriites** Straus, 1967 Neogene
129. *Segistriites cromei* Straus, 1967* Ne Willershausen
- Dysderidae?**
- † **Mistura** Petrunkevitch, 1971 Neogene
130. *Mistura perplexa* Petrunkevitch, 1971* Ne Chiapas amber
- OONOPIDAE Simon, 1890** Cretaceous – Recent
- Oonopidae gen. et sp. in Penney (2002) K New Jersey amber
- † **Burmorchestina** Wunderlich, 2008a Cretaceous
131. *Burmorchestina pulcher* Wunderlich, 2008a* K Myanmar amber
- † **Canadaorchestina** Wunderlich, 2008a Cretaceous
132. *Canadaorchestina albertensis* (Penney, 2006a)* K Manitobian amber
- † **Fossilopaea** Wunderlich, 1988 Neogene
133. *Fossilopaea sulci* Wunderlich, 1988* Ne Dominican amber
- Heteroonops Dalmas, 1916** ?Neogene – Recent
- Heteroonops* sp. in Wunderlich (1988) Ne Dominican amber
- Opopaea Simon, 1891** ?Neogene – Recent
- ?*Opopaea* sp. in Wunderlich (1988) Ne Dominican amber
- Orchestina Simon, 1882** Cretaceous – Recent
134. *Orchestina (Baltorchestina) angulata* Wunderlich, 2012f [replacement name] Pa Bitterfeld amber
- i. = *Orchestina (B.) rectangulata* Wunderlich, 2011h [preoccupied]
135. *Orchestina baltica* Petrunkevitch, 1942 Pa Baltic amber
136. *Orchestina (Baltorchestina) bitterfeldensis* Wunderlich, 2008a Pa Bitterfeld amber
137. *Orchestina breviembolus* Wunderlich, 1981 Pa Baltic amber
138. *Orchestina (Baltorchestina) brevis* Wunderlich, 2008a Pa Baltic amber
139. *Orchestina crassiembolus* Wunderlich, 1981 Pa Baltic amber
140. *Orchestina (Baltorchestina) crassipatellaris* Wunderlich, 1981 Pa Baltic amber

141. *Orchestina (Baltorchestina) crassitibialis* Wunderlich, 1981 Pa Baltic amber
142. *Orchestina (Baltorchestina) colchembolus* Wunderlich, 1981 Pa Baltic amber
143. *Orchestina colombiensis* Wunderlich, 2004at Qt Colombian copal
144. *Orchestina dominicana* Wunderlich, 1981 Ne Dominican amber
145. *Orchestina forceps* Wunderlich, 1981 Pa Baltic amber
146. *Orchestina (Baltorchestina) forfex* Wunderlich, 2011h Pa Baltic amber
147. *Orchestina (Baltorchestina) furca* Wunderlich, 1981 Pa Baltic amber
148. *Orchestina fushunensis* Wunderlich, 2004au Pa Fu Shun amber
149. *Orchestina gappi* Saupe et al., 2012 K Archingeay amber
150. *Orchestina gracilitibialis* Wunderlich, 2004c Pa Baltic amber
151. *Orchestina (Baltorchestina) imperialis* Petrunkevitch, 1963 Pa Baltic/Bitter. amber
152. *Orchestina kenyana* Wunderlich, 1981 Qt East African copal
153. *Orchestina longimana* Wunderlich, 1981 Qt East African copal
154. *Orchestina madagascariensis* Wunderlich, 2004as Qt Madagascan copal
155. *Orchestina mortua* Petrunkevitch, 1971 Ne Chiapas amber
156. *Orchestina (Baltorchestina) multisetae* Wunderlich, 2008a Pa Baltic amber
157. *Orchestina (Gallorchestina) parisiensis* Penney, 2007b Pa Le Quesnoy amber
158. *Orchestina (Baltorchestina) perfecta* Wunderlich, 2008a Pa Baltic amber
159. *Orchestina pusilla* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
160. *Orchestina rabagensis* Saupe et al., 2012 K El Soplao amber
161. *Orchestina (Baltorchestina) rectangulata* Wunderlich, 2008a Pa Baltic amber
162. *Orchestina (Baltorchestina) sternalis* Wunderlich, 2008a Pa Baltic amber
163. *Orchestina tibialis* Wunderlich, 1988 Ne Dominican amber
164. *Orchestina truncata* Wunderlich, 2004at Qt Colombian copal
165. *Orchestina tuberosa* Wunderlich, 1981 Pa Baltic amber
- Orchestina* sp. in Nishikawa (1974) Qt Mizunami copal
- Orchestina* sp. in Saupe et al. (2012) K Álava amber
- Orchestina* sp. in Soriano et al. (2010) K San Just amber
- Orchestina* sp. in Wunderlich (2011h) Pa Bitterfeld amber
- Stenoonops* Simon, 1891** **Palaeogene – Recent**
166. *Stenoonops incertus* (Wunderlich, 1988) Ne Dominican amber
167. ?*Stenoonops rugosus* Wunderlich, 2004c Pa Bitterfeld amber
168. *Stenoonops seldeni* (Penney, 2000) Ne Dominican amber
- ORSOLOBIDAE Cooke, 1965** **Recent**
- no fossil record
- † **PLUMORSOLIDAE** Wunderlich, 2008d **Cretaceous**
- ?Plumorsolidae indet. in Wunderlich (2008d) K Myanmar amber
- ?Plumorsolidae indet. in Wunderlich (2011i) K Myanmar amber
- † **Plumorsolus** Wunderlich, 2008d **Cretaceous**

169. *Plumorsolus gondwanensis* Wunderlich, 2008d K Lebanese amber
- ENTELEGYNAE Simon, 1893** Triassic – Recent
- PALPIMANOIDEA Thorell, 1870a** Jurassic – Recent
- family uncertain
- † **Seppo Selden & Dunlop, 2014** Jurassic
170. *Seppo kopeneni* Selden & Dunlop, 2014* J Grimmen, Germany
- † **Sinaranea Selden, Huang & Ren, 2008** Jurassic
171. *Sinaranea metaxyostraca* Selden, Huang & Ren, 2008* J Daohugou, China
- ARCHAELIDAE C. L. Koch & Berendt, 1854** Jurassic – Recent
- Archaea C. L. Koch & Berendt, 1854** Palaeogene – Recent
172. ?*Archaea bitterfeldensis* Wunderlich, 2004d Pa Bitterfeld amber
173. *Archaea compacta* Wunderlich, 2004d Pa Baltic amber
174. *Archaea paradoxa* C. L. Koch & Berendt, 1854* Pa Baltic amber
- i. = *Archaea laevigata* C. L. Koch & Berendt, 1854 Pa Baltic amber
- ii. = *Archaea incompta* Menge in C. L. Koch & Berendt,
1854 Pa Baltic amber
175. *Archaea poungeti* Simon, 1884b Pa Baltic amber
- † **Baltarchaea Eskov, 1992** Palaeogene
176. *Baltarchaea conica* (C. L. Koch & Berendt, 1854)* Pa Baltic amber
- † **Burmesarchaea Wunderlich, 2008d** Cretaceous
177. *Burmesarchaea grimaldii* (Penney, 2003a) K Myanmar amber
- † **Eoarchaea Forster & Platnick, 1984** Palaeogene
178. *Eoarchaea hyperoptica* (Menge in C. L. Koch & Berendt, 1854)* Pa Baltic amber
179. *Eoarchaea vidua* Wunderlich, 2004d Pa Baltic amber
- † **Eomysmauchenius Wunderlich, 2008d** Cretaceous
180. *Eomysmauchenius septentrionalis* Wunderlich, 2008d* K Myanmar amber
- Eriauchenius O. P.-Cambridge, 1881** Quaternary – Recent
181. *Eriauchenius gracilicollis* (Millot, 1948) [Recent] Qt Copal
- i. = *Archaea copalensis* Lourenço, 2000b Qt Copal
- † **Filiauchenius Wunderlich, 2008d** Cretaceous
182. *Filiauchenius paudentatus* Wunderlich, 2008d* K Myanmar amber
- † **Jurarchaea Eskov, 1987** Jurassic
183. *Jurarchaea zherikhini* Eskov, 1987* J Kazakhstan
- † **Lacunauchenius Wunderlich, 2008d** Cretaceous
184. *Launauchenius speciosus* Wunderlich, 2008d* K Myanmar amber
- † **Myrmecarchaea Wunderlich, 2004d** Palaeogene
185. *Myrmecarchaea petiolus* Wunderlich, 2004d* Pa Baltic amber
186. *Myrmecarchaea pediculus* Wunderlich, 2004d Pa Baltic amber
- † **Patarchaea Selden, Huang & Ren, 2008** Jurassic

187. *Patarchaea muralis* Selden, Huang & Ren, 2008* J Daohugou, China
 † **Saxonarchaea** Wunderlich, 2004d Palaeogene
188. *Saxonarchaea dentata* Wunderlich, 2004d* Pa Bitterfeld amber
 189. *Saxonarchaea diabolica* Wunderlich, 2004d Pa Bitterfeld amber
- MECYSMAUCHENIIDAE Simon, 1895** Cretaceous – Recent
 † **Archaeomecys** Saupe & Selden, 2009 Cretaceous
 190. *Archaeomecys arcantiensis* Saupe & Selden, 2009 K Charente amber
- PARARCHAEIIDAE Forster & Platnick, 1984** Recent
 no fossil record
- HOLARCHAEIIDAE Forster & Platnick, 1984** Recent
 no fossil record
- MICROPHOLCOMMATIDAE Hickman, 1944** Palaeogene – Recent
 † **Cenotextricella** Penney in Penney et al., 2007 Palaeogene
 191. *Cenotextricella simoni* Penney in Penney et al., 2007 Pa Le Quesnoy amber
- HUTTONIIDAE Simon, 1893** Cretaceous – Recent
 unnamed genus and species in Penney & Selden (2006) K Manitoban amber
- STENOCHILIDAE Thorell, 1873** Recent
 no fossil record
- † **MICROPALPIMANIDAE** Wunderlich, 2008d Cretaceous
 † **Micropalpimanus** Wunderlich, 2008d Cretaceous
Micropalpimanus sp. indet in Wunderlich (2012d) K Myanmar amber
 192. *Micropalpimanus poinari* Wunderlich, 2008d K Myanmar amber
- PALPIMANIDAE Thorell, 1870a** Neogene – Recent
 = OTITHOPOIDAE Thorell, 1869 [younger name protected by usage]
 = CERSIDAE Canestrini & Pavesi, 1870
- Otiothops** MacLeay, 1839 Neogene – Recent
Otiothops sp. 1–2 in Wunderlich (1988) Ne Dominican amber
- † **LAGONOMEGOPIDAE** Eskov & Wunderlich, 1995 Cretaceous
 † **Archaelagonops** Wunderlich, 2012d Cretaceous
 193. *Archaelagonops salticoides* Wunderlich, 2012d* K Myanmar amber
- † **Burlagonomegops** Penney, 2005b Cretaceous
 194. *Burlagonomegops alavensis* Penney, 2006b K Álava amber
 195. *Burlagonomegops eskovi* Penney, 2005b* K Myanmar amber
- † **Lagonoburmops** Wunderlich, 2012d Cretaceous

196. *Lagonoburmops plumosus* Wunderlich, 2012d* K Myanmar amber
- † ***Lagonomegops* Eskov & Wunderlich, 1995** Cretaceous
197. *Lagonomegops americanus* Penney, 2005b K New Jersey amber
198. *Lagonomegops sukatchevae* Eskov & Wunderlich, 1995* K Taimyr amber
- † ***Myanlagonops* Wunderlich, 2012d** Cretaceous
199. *Myanlagonops gracilipes* Wunderlich, 2012d* K Myanmar amber
- † ***Zarquagonomegops* Kaddumi, 2007** Cretaceous
200. *Zarquagonomegops wunderlichi* Kaddumi, 2007* K Jordanian amber
- † **GRANDOCULIDAE Penney, 2011** Cretaceous
- NB: The validity of this family has been challenged (cf. Wunderlich 2012d).
- † ***Grandoculus* Penney, 2004b** Cretaceous
201. *Grandoculus chemahawinensis* Penney, 2004b* K Manitobian amber
- † **SPATIATORIDAE Petrunkevitch, 1942** Palaeogene
- † ***Spatiator* Petrunkevitch, 1942** Palaeogene
202. *Spatiator caulis* Wunderlich, 2008a Pa Baltic amber
203. *Spatiator martensi* Wunderlich, 2006 Pa Baltic amber
204. *Spatiator praeceps* Petrunkevitch, 1942* Pa Baltic amber
- Spatiator* sp. in Wunderlich (2011h) Pa Baltic amber
- MALKARIDAE Davies, 1980** Recent
- = STERNODIDAE Moran, 1986
- no fossil record
- MIMETIDAE Simon, 1881** Palaeogene – Recent
- = CTENOPHORIDAE Blackwall, 1870 [younger name protected by usage]
- Mimetidae gen. et sp. indet. in Penney et al. (2012a) Pa Indian amber
- Mimetini sp. 1–4 in Wunderlich (2004q) Pa Baltic amber
- Ero* C. L. Koch, 1836** Palaeogene – Recent
- = †*Palaeoero* Wunderlich, 2004q
- = †*Succinero* Wunderlich, 2004q
- [Wunderlich revalidated both as putative subgenera]
205. *Ero carboneana* Petrunkevitch, 1942 Pa Baltic amber
206. *Ero aberrans* Petrunkevitch, 1958 Pa Baltic amber
- [Treated as a *nomen dubium* by Harms & Dunlop (2009)]
207. *Ero (Succinero) clunis* Wunderlich, 2012c Pa Baltic amber
208. *Ero (Succinero) gracilitibialis* Wunderlich, 2012c Pa Baltic amber
209. *Ero (Paleoero) longitarsus* (Wunderlich, 2004q) Pa Baltic amber
210. *Ero permunda* Petrunkevitch, 1942 Pa Baltic amber
211. *Ero (Succinero) rovnoensis* (Wunderlich, 2004ar) Pa Rovno amber
212. *Ero (Succinero) veta* Wunderlich, 2012c Pa Baltic amber

<i>Mimetus</i> Hentz, 1832	Palaeogene – Recent
? <i>Mimetus</i> sp. in Wunderlich (1988)	Ne Dominican amber
213. <i>Mimetus bituberculatus</i> Wunderlich, 1988	Ne Dominican amber
214. <i>Mimetus brevipes</i> Wunderlich, 2004q	Pa Baltic amber [synonymised by Harms & Dunlop (2009), but resurrected by Wunderlich (2012c)]
215. ? <i>Mimetus longipes</i> Wunderlich, 2004q	Pa Baltic amber
† <i>Protomimetus</i> Wunderlich, 2011	Palaeogene
216. ? <i>Protomimetus breviclypeus</i> Wunderlich, 2011h	Pa Baltic amber
217. <i>Protomimetus longiclypeus</i> Wunderlich, 2011h*	Pa Baltic amber
ERESOIDEA C. L. Koch, 1851	Cretaceous – Recent
ERESIDAE C. L. Koch, 1851	?Miocene – Recent
no body fossil record, but a web attributed to the extant genus <i>Seothyra</i> was described by Pickford (2000) from Miocene aeolianites in the Namib Desert of Namibia	
'OECOBIOIDEA'	
Oecobioidea fam. indet. in Wunderlich (2008d)	K Myanmar amber
OECOBIIDAE Blackwall, 1862	Cretaceous – Recent
= UROCTEIDAE Thorell, 1869	
† <i>Lebanoecobius</i> Wunderlich, 2004e	Cretaceous
218. <i>Lebanoecobius schleei</i> Wunderlich, 2004e*	K Lebanese amber
† <i>Mizalia</i> C. L. Koch & Berendt, 1854	Palaeogene
= † <i>Paruroctea</i> Petrunkevitch, 1942	
219. <i>Mizalia blauvelti</i> (Petrunkevitch, 1942)	Pa Baltic amber
220. <i>Mizalia gemini</i> Wunderlich, 2004e	Pa Baltic amber
221. <i>Mizalia rostrata</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
i. = <i>Mizalia pilosula</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
222. <i>Mizalia spirembolus</i> Wunderlich, 2004e	Pa Baltic amber
<i>Mizalia</i> sp. in Wunderlich (2011h)	Pa Baltic/Blter. amber
<i>Oecobius</i> Lucas, 1846	?Cretaceous – Recent
223. <i>Oecobius piliformis</i> Wunderlich, 1988	Ne Dominican amber
? <i>Oecobius</i> sp. indet in Penney (2002)	K New Jersey amber
<i>Uroctea</i> Dufour, 1820	Palaeogene – Recent
224. <i>Uroctea galloprovincialis</i> Gourret, 1887	Pa Aix-en-Provence
† <i>Zamilia</i> Wunderlich, 2008d	Cretaceous
225. <i>Zamilia antecessor</i> Wunderlich, 2008d	K Myanmar amber
HERSILIIDAE Thorell, 1870a	Cretaceous – Recent
= CHALINUROIDAE Thorell, 1873	
Hersiliidae sp. 1–3 in Wunderlich (2004d)	Pa Baltic amber
Hersiliidae sp. in Wunderlich (2011f)	Qt Madagascar copal

† <i>Burmesiola</i> Wunderlich, 2011 <i>i</i>	Cretaceous
226. <i>Burmesiola cretacea</i> Wunderlich, 2011 <i>*</i>	K Myanmar amber
† "Fictotama" Petrunkevitch, 1963 (<i>nomen dubium</i>)	Neogene
[Wunderlich 2011f placed a new species in this genus, which was previously considered a <i>nomen dubium</i> . He did not formally revalidate the genus]	
227. "Fictotama" <i>maculosa</i> Wunderlich, 2011 <i>g</i>	Ne Dominican amber
† <i>Gerdia</i> Menge, 1869	Palaeogene
228. <i>Gerdia myura</i> Menge, 1869*	Pa Baltic amber
† <i>Gerdiosis</i> Wunderlich, 2004 <i>e</i>	Palaeogene
229. <i>Gerdiosis infringens</i> Wunderlich, 2004 <i>e*</i>	Pa Baltic amber
† <i>Gerdiorum</i> Wunderlich 2004 <i>e</i>	Palaeogene
230. <i>Gerdiorum inflexum</i> Wunderlich 2004 <i>e*</i>	Pa Baltic amber
<i>Hersilia</i> Audouin, 1826	Palaeogene – Recent
= † <i>Hersiliopsis</i> Wunderlich, 2004 <i>e</i>	
231. <i>Hersilia aquisextana</i> Gourret, 1887	Pa Aix-en-Provence
232. <i>Hersilia longipes</i> Giebel, 1856	Pa Baltic amber
233. <i>Hersilia madagascarensis</i> (Wunderlich, 2004 <i>e</i>)	Qt–R Madagas. copal
234. ? <i>Hersilia miranda</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† <i>Hersiliiana</i> Wunderlich, 2004 <i>e</i>	Quaternary – Recent
235. <i>Hersiliiana brevipes</i> Wunderlich, 2004 <i>e*</i>	Qt Madagascan copal
† <i>Prototama</i> Petrunkevitch, 1971	Neogene
= † <i>Priscotama</i> Petrunkevitch, 1971	
236. <i>Prototama antiqua</i> (Petrunkevitch, 1971)	Ne Chiapas amber
237. <i>Prototama maior</i> (Wunderlich, 1988)	Ne Dominican amber
238. <i>Prototama media</i> (Wunderlich, 1988)	Ne Dominican amber
239. <i>Prototama minor</i> (Wunderlich, 1987)	Ne Dominican amber
240. <i>Prototama succinea</i> Petrunkevitch, 1971*	Ne Chiapas amber
<i>Prototama</i> sp. in Wunderlich (1988)	Ne Dominican amber
Superfamily uncertain	
† BURMASCUTIDAE Wunderlich, 2008 <i>d</i>	Cretaceous
† <i>Burmascutum</i> Wunderlich, 2008 <i>d</i>	Cretaceous
241. <i>Burmascutum aenigma</i> Wunderlich, 2008 <i>d*</i>	K Myanmar amber
† SALTICOIDIDAE Wunderlich, 2008 <i>d</i>	Cretaceous
† <i>Salticoidus</i> Wunderlich, 2008 <i>d</i>	Cretaceous
242. <i>Salticoidus kaddumiorum</i> Wunderlich, 2008 <i>d*</i>	K Jordanian amber
'CANOE TAPETUM' CLADE	
ORBICULARIAE Walckenaer, 1802	Triassic – Recent
DEINOPOIDEA C. L. Koch, 1851	?Jurassic – Recent
DEINOPIDAE C. L. Koch, 1851	Cretaceous – Recent

<i>Deinopis</i> MacLeay, 1839	Quaternary – Recent
243. <i>Deinopis ?madagascariensis</i> Lenz, 1886 [Recent]	Qt Madagascar copal
<i>Menneus</i> Simon, 1876b	Palaeogene – Recent
244. ? <i>Menneus pietrzeniukae</i> Wunderlich, 2004g	Pa Baltic amber
? <i>Menneus</i> sp. 1–3 in Wunderlich (2004g)	Pa Baltic amber
† <i>Palaeomicromennus</i> Penney, 2003b	Cretaceous
245. <i>Palaeomicromenneus lebanensis</i> Penney, 2003b*	K Lebanese amber
 ULOBORIDAE Thorell, 1869	?Jurassic – Recent
Uloboridae indet. in Wunderlich (2011f)	Qt Madagascar copal
† <i>Talbragaraneus</i> Selden & Beattie, 2013 [tentative assignment]	Jurassic
246. <i>Talbragaraneus jurassicus</i> Selden & Beattie, 2013*	J Talbragar, Australia
† <i>Burmuloborus</i> Wunderlich, 2008d	Cretaceous
247. <i>Burmuloborus parvus</i> Wunderlich, 2008d*	K Myanmar amber
† <i>Eomiagrammopes</i> Wunderlich, 2004f	Palaeogene
248. <i>Eomiagrammopes maior</i> Wunderlich, 2004f	Pa Baltic amber
249. <i>Eomiagrammopes minor</i> Wunderlich, 2004f	Pa Baltic amber
250. <i>Eomiagrammopes semiapertus</i> Wunderlich, 2011h	Pa Baltic amber
251. <i>Eomiagrammopes singularis</i> Wunderlich, 2004f*	Pa Baltic amber
252. <i>Eomiagrammopes spinipes</i> Wunderlich, 2004f	Pa Baltic amber
<i>Eomiagrammopes</i> sp. 1–2 in Wunderlich (2004f)	Pa Baltic amber
? <i>Eomiagrammopes</i> sp. in Wunderlich (2004f)	Pa Baltic amber
† <i>Hyptiomopes</i> Wunderlich, 2004f	Palaeogene
253. <i>Hyptiomopes bitterfeldensis</i> Wunderlich 2004f*	Pa Bitterfeld amber
? <i>Hyptiomopes</i> sp. in Wunderlich (2004f)	Pa Bitterfeld amber
<i>Hyptiotes</i> Walckenaer, 1837	Palaeogene – Recent
= † <i>Androgeus</i> C. L. Koch & Berendt, 1854	
254. <i>Hyptiotes convexus</i> Wunderlich, 2004f	Pa Baltic amber
255. <i>Hyptiotes glaber</i> Wunderlich, 2004f	Pa Baltic amber
256. <i>Hyptiotes saetosus</i> Wunderlich, 2004f	Pa Baltic amber
257. <i>Hyptiotes stellatus</i> Wunderlich, 2004f	Pa Baltic amber
258. <i>Hyptiotes triqueter</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
† <i>Jerseyuloborus</i> Wunderlich, 2011i	Cretaceous
259. <i>Jerseyuloborus longisoma</i> Wunderlich, 2011i*	K New Jersey amber
<i>Miagrammopes</i> O. P.-Cambridge, 1870	Neogene – Recent
260. <i>Miagrammopes dominicanus</i> Wunderlich, 2004e	Ne Dominican amber
<i>Miagrammopes</i> sp. in Penney (2001)	Ne Dominican amber
<i>Miagrammopes</i> sp. in Wunderlich (2011f)	Qt Madagascar copal
† <i>Ocululoborus</i> Wunderlich, 2012d	Cretaceous
261. <i>Ocululoborus curvatus</i> Wunderlich, 2012d*	K Myanmar amber
† <i>Opellianus</i> Wunderlich, 2004f	Palaeogene

262. *Opellianus excellens* Wunderlich, 2004^{f*} Pa Baltic amber
263. *Opellianus kazimierasi* Wunderlich 2004^f Pa Baltic amber
264. *Opellianus ludwigi* Wunderlich 2004^f Pa Baltic amber
- † ***Palaeomiagrammopes* Wunderlich, 2008d** Cretaceous
265. *Palaeomiagrammopes vesica* Wunderlich, 2008d^{*} K Myanmar amber
- † ***Palaeouloborus* Selden, 1990** Cretaceous
266. *Palaeouloborus lacasae* Selden, 1990^{*} K Sierra de Montsech
- † ***Paramiagrammopes* Wunderlich, 2008d** Cretaceous
267. *Paramiagrammopes cretaceus* Wunderlich, 2008d^{*} K Myanmar amber
- Paramiagrammopes* sp. in Wunderlich (2008d) K Myanmar amber
- † ***Ulobomopes* Wunderlich, 2004f** Palaeogene
268. *Ulobomopes unicus* Wunderlich, 2004^{f*} Pa Baltic amber
- ARANEOIDEA Latreille, 1806** Jurassic – Recent
- Araneoidea fam indet. in Wunderlich (2008d) K Myanmar amber
- † ***Mesarania* Hong, 1984** Jurassic
269. *Mesarania hebeiensis* Hong, 1984^{*} J Hebei, China
- CYATHOLIPIDAE Simon, 1894** Palaeogene – Recent
- = TEEMENAARIDAE Davies, 1978
- † ***Balticolipus* Wunderlich, 2004m** Palaeogene
270. *Balticolipus kruemmeri* Wunderlich, 2004m^{*} Pa Baltic / Bitt. amber
- † ***Cyathosuccinus* Wunderlich, 2004m** Palaeogene
271. *Cyathosuccinus elongatus* Wunderlich, 2004m^{*} Pa Baltic amber
- † ***Erigolipus* Wunderlich, 2004m** Palaeogene
272. *Erigolipus griswoldi* Wunderlich, 2004m^{*} Pa Baltic amber
- † ***Spinilipus* Wunderlich, 1993b** Palaeogene
273. *Spinilipus bispinosus* Wunderlich, 2004m Pa Bitterfeld amber
274. *Spinilipus curvatus* Wunderlich, 2004m Pa Bitterfeld amber
275. *Spinilipus glinki* Wunderlich, 2004m Pa Baltic amber
276. *Spinilipus kerneggeri* Wunderlich, 1993b^{*} Pa Baltic amber
277. *Spinilipus longembolus* Wunderlich, 2004m Pa Baltic amber
- † ***Succinilipus* Wunderlich, 1993b** Palaeogene
278. *Succinilipus abditus* Wunderlich, 2004m Pa Baltic / Bitt. amber
279. *Succinilipus aspinosus* Wunderlich, 2004m Pa Bitterfeld amber
280. *Succinilipus saxonensis* Wunderlich, 1993b Pa Bitterfeld amber
281. *Succinilipus similis* Wunderlich, 2004m Pa Bitterfeld amber
282. *Succinilipus teuberi* Wunderlich, 1993b^{*} Pa Baltic amber
- Succinilipus* sp. in Wunderlich (2004m) Pa Baltic / Bitt. amber
- SYNOTAXIDAE Simon, 1894** Palaeogene – Recent
- † ***Acrometa* Petrunkevitch, 1942** Palaeogene

- = † *Eogonatum* Petrunkevitch, 1942
 = † *Liticens* Petrunkevitch, 1942
 = † *Theridiometa* Petrunkevitch, 1942
 = † *Viocurus* Petrunkevitch, 1958
283. *Acrometa clava* Wunderlich, 2004n Pa Baltic amber
284. *Acrometa cristata* Petrunkevitch, 1942* Pa NE Europe ambers
- i. = *Theridiometa edwardsi* Petrunkevitch, 1942 Pa Baltic amber
 ii. = *Viocurus fossilis* Petrunkevitch, 1958 Pa Baltic amber
285. *Acrometa eichmanni* Wunderlich, 2004n Pa Baltic amber
286. *Acrometa incidunt* Wunderlich, 2004n Pa Baltic amber
287. *Acrometa minutum* (Petrunkevitch, 1942) Pa Baltic amber
288. *Acrometa pala* Wunderlich, 2004n Pa Baltic amber
289. *Acrometa robusta* (Petrunkevitch, 1942) Pa Baltic amber
290. *Acrometa pseudorobusta* Dunlop & Jekel, 2009 Pa Baltic amber
- i. = *Acrometa robusta* (Petrunkevitch, 1946) [preoccupied]
291. *Acrometa samlandica* (Petrunkevitch, 1942) Pa Baltic amber
292. *Acrometa setosus* (Petrunkevitch, 1942) Pa Baltic amber
293. *Acrometa succini* Petrunkevitch, 1942 Pa Baltic amber
- † **Anandrus** Menge, 1856 Palaeogene
- = † *Elucus* Petrunkevitch, 1942
294. *Anandrus inermis* (Petrunkevitch, 1942) Pa Baltic amber
295. *Anandrus infelix* (Petrunkevitch, 1950)* Pa Baltic amber
296. *Anandrus quaesitus* (Petrunkevitch, 1958) Pa Baltic amber
297. *Anandrus redemptus* (Petrunkevitch, 1958) Pa Baltic amber
- † **Chelicerinus** Wunderlich, 2008a Palaeogene
298. *Chelicerinus abnormis* Wunderlich, 2008a Pa Bitterfeld amber
- † **Cornuanandrus** Wunderlich, 1986 Palaeogene
299. *Cornuanandrus bifurcatus* Wunderlich, 2004n Pa Bitterfeld amber
300. *Cornuanandrus bitterfeldensis* Wunderlich, 2004n Pa Bitterfeld amber
301. *Cornuanandrus corniculans* Wunderlich, 2004n Pa Baltic amber
302. *Cornuanandrus maior* Wunderlich, 1986* Pa Baltic amber
303. *Cornuanandrus minor* Wunderlich, 2004n Pa Baltic amber
- † **Dubiosynotaxus** Wunderlich, 2004n Palaeogene
304. *Dubiosynotaxus perfectus* Wunderlich, 2004n* Pa Baltic amber
- † **Eosynotaxus** Wunderlich, 2004n Palaeogene
305. *Eosynotaxus bispinosus* Wunderlich, 2004n Pa Baltic amber
306. *Eosynotaxus bitterfeldensis* Wunderlich, 2004n Pa Bitterfeld amber
307. *Eosynotaxus custodens* Wunderlich, 2004n Pa Baltic amber
308. *Eosynotaxus fastigatus* Wunderlich, 2004n Pa Baltic amber
309. *Eosynotaxus paucispina* Wunderlich, 2004n Pa Baltic amber
310. *Eosynotaxus spinipes* Wunderlich, 2004n Pa Baltic amber
311. *Eosynotaxus wegneri* Wunderlich, 2004n* Pa Baltic amber

† <i>Gibbersynotaxus</i> Wunderlich, 2004n	Palaeogene
312. <i>Gibbersynotaxus parvus</i> Wunderlich, 2004n*	Pa Baltic amber
† <i>Protophysoglenes</i> Wunderlich, 2004n	Palaeogene
313. <i>Protophysoglenes impressum</i> Wunderlich, 2004n*	Pa Baltic amber
† <i>Pseudoacrometa</i> Wunderlich, 1986	Palaeogene
314. <i>Pseudoacrometa gracilipes</i> Wunderlich, 1986*	Pa Baltic amber
315. <i>Pseudoacrometa wittmanni</i> Wunderlich, 2004n	Pa Baltic amber
† <i>Succinitaxus</i> Wunderlich, 2004n	Palaeogene
316. <i>Succinitaxus brevis</i> Wunderlich, 2004n*	Pa Baltic, Bitterfeld & Rovno amber
317. ? <i>Succinitaxus minutus</i> Wunderlich, 2004n	Pa Baltic amber
† <i>Sulcosynotaxus</i> Wunderlich, 2004n	Palaeogene
318. <i>Sulcosynotaxus cavatus</i> Wunderlich, 2004n*	Pa Baltic amber
 NESTICIDAE Simon, 1894	Palaeogene – Recent
† <i>Balticonesticus</i> Wunderlich, 1986	Palaeogene
319. <i>Balticonesticus flexuosus</i> Wunderlich, 1986*	Pa Baltic amber
<i>Eidmanella</i> Roewer, 1935	Quaternary
320. <i>Eidmanella pallida</i> (Emerton, 1875) [Recent]	Qt Madagascar copal
† <i>Eopopino</i> Petrunkevitch, 1942	Palaeogene
321. <i>Eopopino budrysi</i> Eskov & Marusik, 1992	Pa Baltic amber
322. <i>Eopopino inopinatus affinis</i> Wunderlich, 1986	Pa Baltic amber
323. <i>Eopopino inopinatus inopinatus</i> Wunderlich, 1986	Pa Baltic amber
324. <i>Eopopino longipes</i> Petrunkevitch, 1942*	Pa Baltic amber
325. <i>Eopopino palanga</i> Eskov & Marusik, 1992	Pa Baltic amber
326. <i>Eopopino rarus rarus</i> Wunderlich, 1986	Pa Baltic amber
327. <i>Eopopino rarus solitarius</i> Wunderlich, 1986	Pa Baltic amber
328. <i>Eopopino rudloffii</i> Wunderlich, 2004o	Pa Bitterfeld amber
<i>Eopopino</i> sp. in Wunderlich (1986)	Pa Bitterfeld amber
† <i>Heteronesticus</i> Wunderlich, 1986	Palaeogene
329. <i>Heteronesticus magnoparacymbialis</i> Wunderlich, 1986*	Pa Baltic amber
† <i>Hispanonesticus</i> Wunderlich, 1986	Neogene
330. <i>Hispanonesticus latopalpus</i> Wunderlich, 1986*	Ne Dominican amber
 THERIDIIDAE Sundevall, 1833	?Cretaceous – Recent
= PHYCOIDAE Thorell, 1873	
= EPISINIDAE O. P.-Cambridge, 1879a	
= HADROTARSIDAE Thorell, 1881	
?Theridiidae gen. et sp. indet in McAlpine & Martin (1969)	K Canadian amber
Theridiidae gen. et sp. in Nishikawa (1974)	Qt Mizunami copal
<i>Achaearanea</i> Strand, 1929	Neogene – Recent
331. <i>Achaearanea extincta</i> Wunderlich, 1988	Ne Dominican amber

<i>Achaeareana</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
Argyrodes Simon, 1864	Neogene – Recent
332. <i>Argyrodes (Ariamnes) copalis</i> Wunderlich, 2008b	Qt Colombian copal
333. <i>Argyrodes (Ariamnes) resina</i> Wunderlich, 2011f.....	Qt Madagascar copal
334. <i>Argyrodes (Rhomphaea) gibbifera</i> Wunderlich, 2004as	Qt Madagascar copal
335. <i>Argyrodes parvipatellaris</i> Wunderlich, 1988	Ne Dominican amber
<i>Argyrodes</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
† Balticoridion Wunderlich, 2008b	Palaeogene
336. <i>Balticoridion dubium</i> Wunderlich, 2008b*	Pa Baltic / Bitt. amber
† Balticpholcomma Wunderlich, 2008b	Palaeogene
337. <i>Balticpholcomma scutatum</i> Wunderlich, 2008b*	Pa Baltic amber
† Caudasinus Wunderlich, 2008b	Palaeogene
338. <i>Caudasinus bispinosus</i> Wunderlich, 2008b	Pa Baltic amber
339. <i>Caudasinus caudatus</i> Wunderlich, 2008b*	Pa Baltic amber
340. <i>Caudasinus regeneratus</i> Wunderlich, 2008b	Pa Baltic amber
<i>Caudasinus</i> sp. <i>in</i> Wunderlich (2008b)	Pa Baltic amber
Chrosiothes Simon, 1894	Neogene – Recent
341. <i>Chrosiothes biconigerus</i> Wunderlich, 1988	Ne Dominican amber
342. <i>Chrosiothes curvispinosus</i> Wunderlich, 1988	Ne Dominican amber
343. <i>Chrosiothes emulgatus</i> Wunderlich, 1988	Ne Dominican amber
344. <i>Chrosiothes longispinosus</i> Wunderlich, 1988	Ne Dominican amber
345. <i>Chrosiothes monoceros</i> Wunderlich, 1988	Ne Dominican amber
346. <i>Chrosiothes tumulus</i> Wunderlich, 1988	Ne Dominican amber
347. <i>Chrosiothes unicornis</i> Wunderlich, 1988	Ne Dominican amber
Chrysso O. P.-Cambridge, 1882a	Neogene – Recent
348. <i>Chrysso conspicua</i> Wunderlich, 1988	Ne Dominican amber
349. <i>Chrysso dubia</i> Wunderlich, 1988	Ne Dominican amber
† Clavibertus Wunderlich, 2008b	Palaeogene
350. <i>Clavibertus parvus</i> Wunderlich, 2008b	Pa Baltic amber
351. <i>Clavibertus prominens</i> Wunderlich, 2008b*	Pa Baltic amber
† Clya C. L. Koch & Berendt, 1854	Palaeogene
352. <i>Clya abdita</i> Wunderlich, 2008b	Pa Baltic amber
353. <i>Clya lugubris</i> C. L. Koch & Berendt, 1854*	Pa Baltic / Rovno amber
354. <i>Clya calefacta</i> Wunderlich, 2008b	Pa Baltic amber
355. <i>Clya gracilis</i> (Petrunkevitch, 1958)	Pa Baltic amber
356. <i>Clya granulata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
357. <i>Clya obscura</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
358. <i>Clya rotata</i> Wunderlich, 2008b	Pa Baltic amber
359. <i>Clya supercalefacta</i> Wunderlich, 2008b	Pa Baltic amber
360. <i>Clya superspiralis</i> Wunderlich, 2008b	Pa Baltic amber
361. <i>Clya tricurvata</i> Wunderlich, 2008b	Pa Baltic amber

- † *Cornutidion* Wunderlich, 1988 Neogene
 362. *Cornutidion elongatum* Wunderlich, 1988* Ne Dominican amber
- Craspedisia* Simon, 1894 Neogene – Recent
 363. *Craspedisia yapchoontecki* Penney & Marusik *in* Penney et al.
 (2012b) Ne Dominican amber
- † *Cymbiopholcomma* Wunderlich, 2008b Palaeogene
 364. *Cymbiopholcomma dudum* Wunderlich, 2008b* Pa Baltic amber
 365. *Cymbiopholcomma spiculum* Wunderlich, 2008b Pa Baltic amber
- † *Dipoenata* Wunderlich, 1988 Neogene
 366. *Dipoenata altioculata* Wunderlich, 1988 Ne Dominican amber
 367. *Dipoenata cala* Wunderlich, 1988 Ne Dominican amber
 368. *Dipoenata clypeata* Wunderlich, 1988 Ne Dominican amber
 369. *Dipoenata globulus* Wunderlich, 1988 Ne Dominican amber
 370. *Dipoenata praedominicana* (Wunderlich, 1986) Qt Dominican copal
 371. *Dipoenata stipes* Wunderlich, 1988* Ne Dominican amber
 372. *Dipoenata yolanda* Wunderlich, 1988 Ne Dominican amber
Dipoenata sp. *in* Wunderlich (1988) Ne Dominican amber
- † *Eoasagena* Wunderlich, 2008b Palaeogene
 373. *Eoasagena scutata* Wunderlich, 2008b* Pa Baltic amber
- † *Eolyrifer* Wunderlich, 2008b Palaeogene
 374. *Eolyrifer longitibialis* Wunderlich, 2008b* Pa Baltic amber
- † *Eomysmena* Petrunkevitch, 1942 Palaeogene – Neogene
 = † *Antopia* Menge, 1854 [tentative synonymy]
 = † *Astodipoena* Petrunkevitch, 1958
 = † *Eodipoena* Petrunkevitch, 1942
 375. *Eomysmena asta* Petrunkevitch, 1971 Ne Chiapas amber
 376. *Eomysmena aviceps* Wunderlich, 2008b Pa Baltic amber
 377. *Eomysmena calefacta* Wunderlich, 2008b Pa Baltic amber
 378. *Eomysmena crassa* (Petrunkevitch, 1958) Pa Baltic amber
 379. *Eomysmena baltica* Petrunkevitch, 1946 Pa Baltic amber
 380. 'Eomysmena' *bassleri* (Petrunkevitch, 1942) Pa Baltic amber
 381. ?*Eomysmena kaestneri* (Petrunkevitch, 1958) Pa Baltic amber
 382. *Eomysmena militaris* (C. L. Koch & Berendt, 1854) Pa Baltic amber
 383. *Eomysmena moritura* Petrunkevitch, 1942* Pa Baltic amber
 i. = *Eomysmena consulta* (Petrunkevitch, 1958)
 [tentative synonymy] Pa Baltic amber
 384. *Eomysmena nielseni* (Petrunkevitch, 1958) Pa Baltic amber
 385. *Eomysmena oculata* (Petrunkevitch, 1942) Pa Baltic amber
 386. *Eomysmena punctulata* (C. L. Koch & Berendt, 1854) Pa Baltic amber
 387. *Eomysmena recta* Wunderlich, 2008b Pa Baltic amber
 388. *Eomysmena tenera* (Menge *in* C. L. Koch & Berendt, 1854) Pa Baltic amber

<i>Eomysmena</i> spp. in Wunderlich 2008b	Pa	Baltic / Bitt. Amber
† <i>Eoteutana</i> Wunderlich, 2008b		Palaeogene
389. <i>Eoteutana hirsuta</i> Wunderlich, 2008b*	Pa	Baltic amber
<i>Episinus</i> Latreille, 1809		Palaeogene – Recent
= † <i>Flegia</i> C. L. Koch & Berendt, 1854		
= † <i>Impulsor</i> Petrunkevitch, 1942		
= † <i>Malleator</i> Petrunkevitch, 1942		
= † <i>Mictodipoena</i> Petrunkevitch, 1958		
= † <i>Municeps</i> Petrunkevitch, 1942 [tentative synonymy]		
390. <i>Episinus anapidaeque</i> Wunderlich, 2008b	Pa	Baltic amber
391. <i>Episinus antecognatus</i> Wunderlich, 1986	Qt	Dominican copal
392. <i>Episinus appendix</i> Wunderlich, 2008b	Pa	Baltic amber
393. <i>Episinus arrodens</i> Wunderlich, 2008b	Pa	Baltic amber
394. <i>Episinus balticus</i> Marusik & Penney, 2004	Pa	Baltic / Bitt. amber
395. <i>Episinus brevipalpus</i> Wunderlich, 1988	Ne	Dominican amber
396. <i>Episinus bulla</i> Wunderlich, 2008b	Pa	Baltic amber
397. <i>Episinus chiapasanus</i> (Petrunkevitch, 1971)	Ne	Chiapas amber
398. <i>Episinus clunis</i> Wunderlich, 2008b	Pa	Baltic amber
399. <i>Episinus cochlear</i> Wunderlich, 2008b	Pa	Baltic amber
400. <i>Episinus cornutus</i> Wunderlich, 1988	Ne	Dominican amber
401. <i>Episinus cymbialis</i> Wunderlich, 2008b	Pa	Baltic amber
402. <i>Episinus dimidiatus</i> Wunderlich, 2008b	Pa	Baltic amber
403. <i>Episinus eskovi</i> Marusik & Penney, 2004	Pa	Baltic amber
404. <i>Episinus isopteraque</i> Wunderlich, 2008b	Pa	Baltic amber
405. <i>Episinus latus</i> Wunderlich, 2008b	Pa	Baltic amber
406. <i>Episinus longimanus</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
i. = <i>Malleator niger</i> Petrunkevitch, 1942	Pa	Baltic amber
407. <i>Episinus longisoma</i> Wunderlich, 2008b	Pa	Baltic amber
408. <i>Episinus minutus</i> (Petrunkevitch, 1958)	Pa	Baltic amber
409. <i>Episinus mordellidaeque</i> Wunderlich, 2008b	Pa	Baltic amber
410. <i>Episinus musculus</i> Wunderlich, 2008b	Pa	Baltic amber
411. <i>Episinus mutilus</i> (Petrunkevitch, 1958)	Pa	Baltic amber
412. <i>Episinus nausticymbium</i> Wunderlich, 2008b	Pa	Baltic amber
413. <i>Episinus neglectus</i> (Petrunkevitch, 1942)	Pa	Baltic amber
414. <i>Episinus penneyi</i> Garcia-Villafuerte, 2006a	Ne	Chiapas amber
415. <i>Episinus praecognatus</i> Wunderlich, 1982	Ne	Dominican amber
416. <i>Episinus pulcher</i> (Petrunkevitch, 1942)	Pa	Baltic amber
417. <i>Episinus regalis</i> (Petrunkevitch, 1958)	Pa	Baltic amber
418. <i>Episinus stridulus</i> (Petrunkevitch, 1958)	Pa	Baltic amber
419. <i>Episinus tibiaseta</i> Wunderlich, 2011g	Ne	Dominican amber
420. <i>Episinus transversus</i> Wunderlich, 2008b	Pa	Baltic amber
421. <i>Episinus tuberosus</i> Wunderlich, 1988	Ne	Dominican amber

<i>Episinus</i> spp. in Wunderlich (2008b)	Pa	Baltic amber
<i>Euryopis</i> Menge, 1868		Palaeogene – Recent
422. ? <i>Euryopis araneoides</i> Wunderlich, 2008b	Pa	Baltic amber
423. <i>Euryopis bitterfeldensis</i> Wunderlich, 2008b	Pa	Baltic / Bitt. amber
424. <i>Euryopis nexus</i> Wunderlich, 2008b	Pa	Baltic amber
425. <i>Euryopis streyi</i> Wunderlich, 2008b	Pa	Baltic / Bitt. Amber
<i>Euryopis/Emertonella</i> complex in Penney et al. (2012c)	Qt	Colombian copal
† <i>Euryopus</i> Menge in C. L. Koch & Berendt, 1854		Palaeogene
426. <i>Euryopus gracilipes</i> Menge in C. L. Koch & Berendt, 1854*	Pa	Baltic amber
<i>Faiditus</i> Keyserling, 1884		Neogene – Recent
427. <i>Faiditus crassipatellaris</i> (Wunderlich, 1988)	Ne	Dominican amber
† <i>Femurrapator</i> Wunderlich, 2011g		Neogene
428. <i>Femurrapator dominicanus</i> Wunderlich, 2011g*	Ne	Dominican amber
† <i>Globulidion</i> Wunderlich, 2008b		Palaeogene
429. <i>Globulidion cochlea</i> Wunderlich, 2008b*	Pa	Baltic amber
† <i>Hirsutipalpus</i> Wunderlich, 2008b		Palaeogene
430. <i>Hirsutipalpus varipes</i> Wunderlich, 2008b*	Pa	Baltic / Bitt. Amber
† <i>Kochiuridion</i> Wunderlich, 2008b		Palaeogene
431. <i>Kochiuridion scutatum</i> Wunderlich, 2008b*	Pa	Baltic / Bitt. amber
<i>Lasaeola</i> Simon, 1881		Palaeogene – Recent
= † <i>Nactodipoena</i> Petrunkevitch, 1942 [a subgenus in Wunderlich (2008b)]		
432. <i>Lasaeola acumen</i> Wunderlich, 2008b	Pa	Baltic amber
433. <i>Lasaeola baltica</i> (Marusik & Penney, 2004)	Pa	Baltic amber
434. <i>Lasaeola bitterfeldensis</i> Wunderlich, 2008b	Pa	Bitterfeld amber
435. <i>Lasaeola communis</i> Wunderlich, 2008b	Pa	Baltic amber
436. <i>Lasaeola (Nactodipoena) dunbari</i> (Petrunkevitch, 1942)	Pa	Baltic amber
437. ? <i>Lasaeola furca</i> Wunderlich, 2008b	Pa	Baltic amber
438. <i>Lasaeola germanica</i> (Petrunkevitch, 1958)	Pa	Baltic amber
439. <i>Lasaeola (Phycosoma) inclinata</i> Wunderlich, 2012a	Qt	Madagascan copal
440. <i>Lasaeola infulata</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic / Bitt. Amber
441. <i>Lasaeola larvaque</i> Wunderlich, 2008b	Pa	Baltic amber
442. <i>Lasaeola latisulci</i> Wunderlich, 2008b	Pa	Baltic amber
443. <i>Lasaeola pristina</i> (Wunderlich, 1986)	Ne	Dominican amber
444. <i>Lasaeola puta</i> Wunderlich, 1988	Ne	Dominican amber
445. <i>Lasaeola sexsaetosa</i> Wunderlich, 2008b	Pa	Baltic amber
446. ? <i>Lasaeola sigillata</i> Wunderlich, 2008b	Pa	Bitterfeld amber
447. <i>Lasaeola vicina</i> (Wunderlich, 1982)	Ne	Dominican amber
448. <i>Lasaeola vicinoides</i> Wunderlich, 1988	Ne	Dominican amber
<i>Lasaeola</i> sp. in Wunderlich (1988)	Ne	Dominican amber
<i>Lasaeola</i> spp. in Wunderlich (2008b)	Pa	Baltic / Bitt. amber
† <i>Medela</i> Petrunkevitch, 1942 [?Theridiidae, cf. Wunderlich (2008b)].....		Palaeogene

449. *Medela baltica* Petrunkevitch, 1942* Pa Baltic amber
- † ***Mimetidion*** Wunderlich, 2008b Palaeogene
450. *Mimetidion furca* Wunderlich, 2008b* Pa Baltic amber
- † ***Nanomysmena*** Petrunkevitch, 1958 Palaeogene
451. *Nanomysmena aculeata* Petrunkevitch, 1958 Pa Baltic amber
452. *Nanomysmena munita* Petrunkevitch, 1958 Pa Baltic amber
453. *Nanomysmena palanga* Marusik & Penney, 2004 Pa Baltic amber
454. *Nanomysmena petrunkevitchi* Marusik & Penney, 2004 Pa Baltic amber
455. *Nanomysmena pseudogracilis* Marusik & Penney, 2004 Pa Baltic amber
- † ***Nanosteatoda*** Wunderlich, 2008b Palaeogene
456. *Nanosteatoda breviscutum* Wunderlich, 2008b Pa Baltic amber
457. *Nanosteatoda trisetae* Wunderlich, 2008b Pa Baltic amber
- † ***Obscuropholcomma*** Wunderlich, 2008b Palaeogene
458. *Obscuropholcomma* sp. in Wunderlich (2012b) Pa Rovno amber
459. *Obscuropholcomma tegens* Wunderlich, 2008b* Pa Baltic amber
- Phoroncidia*** Westwood, 1835 Quaternary – Recent
460. *Phoroncidia ?aculeata* Westwood, 1835 [Recent] Qt Madagascan copal
- Platnickina*** Koçak & Kemal, 2008 Quaternary – Recent
461. *Platnickina duosetae* Wunderlich, 2012a Qt Madagascan copal
- † ***Praetereuryopis*** Wunderlich, 2008b Palaeogene
462. *Praetereuryopis phoroncidoides* Wunderlich, 2008b* Pa Baltic amber
- † ***Pronepos*** Petrunkevitch, 1963 Neogene
463. *Pronepos exilis* Petrunkevitch, 1963* Ne Chiapas amber
464. *Pronepos fossilis* Petrunkevitch, 1963 Ne Chiapas amber
- † ***Protosteatoda*** Wunderlich, 2008b Palaeogene
465. *Protosteatoda gutta* Wunderlich, 2008b Pa Baltic amber
- † ***Pseudoteutana*** Wunderlich, 2008b Palaeogene
466. *Pseudoteutana stigmatosa* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- i. = *Eomysmena stridens* Petrunkevitch, 1958 Pa Baltic amber
- ii. = *Flegia succini* Petrunkevitch, 1942 Pa Baltic amber
- † ***Rugopholcomma*** Wunderlich, 2008b Palaeogene
467. *Rugopholcomma patellaris* Wunderlich, 2008b* Pa Baltic amber
- † ***Spinisinus*** Wunderlich, 2008b Palaeogene
468. *Spinisinus parvioculi* Wunderlich, 2008b Pa Baltic amber
469. *Spinisinus splendidus* Wunderlich, 2008b* Pa Baltic amber
- † ***Spinitharinus*** Wunderlich, 2008b Palaeogene
470. *Spinitharinus bulbosus* Wunderlich, 2008b* Pa Baltic / Bitt. amber
471. *Spinitharinus cheliceratus* Wunderlich, 2008b Pa Baltic / Bitt. amber
472. *Spinitharinus coniectens* Wunderlich, 2008b Pa Baltic amber
473. *Spinitharinus curvatus* Wunderlich, 2008b Pa Baltic amber
474. *Spinitharinus cymbioseta* Wunderlich, 2008b Pa Baltic amber

<i>Spinitharinus</i> spp. in Wunderlich (2008b)	Pa	Baltic amber
<i>Spintharus</i> Hentz, 1850		Neogene – Recent
475. <i>Spintharus longisoma</i> Wunderlich, 1988	Ne	Dominican amber
<i>Steatoda</i> Sundevall, 1833		?Palaeogene – Recent
476. 'Steatoda' <i>anticus</i> (Berland, 1939)	Pa	Baltic amber
<i>Stemmops</i> O. P.-Cambridge, 1894		Neogene – Recent
477. <i>Stemmops incertus</i> Wunderlich, 1988	Ne	Dominican amber
478. <i>Stemmops prominens</i> Wunderlich, 1988	Ne	Dominican amber
<i>Styposis</i> Simon, 1894		Neogene – Recent
479. <i>Styposis pholcoides</i> Wunderlich, 1988	Ne	Dominican amber
† <i>Succinobertus</i> Wunderlich, 2008b		Palaeogene
480. <i>Succinobertus adjacens</i> Wunderlich, 2008b*	Pa	Baltic / Bitt. Amber
† <i>Succinura</i> Wunderlich, 2008b		Palaeogene
481. <i>Succinura aciesaeta</i> Wunderlich, 2008b	Pa	Baltic amber
482. <i>Succinura bellavista</i> Wunderlich, 2008b*	Pa	Baltic amber
483. <i>Succinura circuta</i> Wunderlich, 2008b	Pa	Baltic amber
484. <i>Succinura dubia</i> Wunderlich, 2008b	Pa	Baltic amber
485. <i>Succinura fuscoruber</i> Wunderlich, 2008b	Pa	Baltic amber
486. <i>Succinura ovalis</i> Wunderlich, 2008b	Pa	Baltic amber
<i>Succinura</i> sp. in Wunderlich (2008b)	Pa	Baltic amber
<i>Theridion</i> Walckenaer, 1805		?Cretaceous – Recent
487. 'Theridion' <i>alutaceum</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
488. <i>Theridion annulipes</i> Heer, 1865	Ne	Öhningen
489. <i>Theridion atalus</i> Chang, 2004 [both generic and familial assignment unreliable!]	K	Jehol Biota
490. 'Theridion' <i>berendti</i> Marusik & Penney, 2004	Pa	Baltic amber
i. = <i>Theridion globosa</i> C. L. Koch & Berendt, 1854 [preoccupied]		
491. <i>Theridion bucklandi</i> Thorell, 1870a	Pa	Aix-en-Provence
492. <i>Theridion contrarium</i> Wunderlich, 1988	Ne	Dominican amber
493. <i>Theridion crassipalpum</i> Berland, 1939	Pa	Aix-en-Provence
494. 'Theridion' <i>detersum</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
495. <i>Theridion erectoides</i> Wunderlich, 1988	Ne	Dominican amber
496. <i>Theridion erectum</i> Wunderlich, 1988	Ne	Dominican amber
497. 'Theridion' <i>globosus</i> (Presl, 1822)	Pa	Baltic amber
498. <i>Theridion globulus</i> Heer, 1865	Ne	Öhningen
499. 'Theridion' <i>hirtum</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
500. <i>Theridion inversum</i> Wunderlich, 1988	Ne	Dominican amber
501. <i>Theridion maculipes</i> Heer, 1865	Ne	Öhningen
502. 'Theridion' <i>oblongum</i> (Presl, 1822)	Pa	Baltic amber
503. 'Theridion' <i>ovale</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
504. 'Theridion' <i>ovatum</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber

505. '*Theridion*' *simplex* C. L. Koch & Berendt, 1854 Pa Baltic amber
506. *Theridion variosoma* Wunderlich, 1988 Ne Dominican amber
507. *Theridion wunderlichi* Penney, 2001 Ne Dominican amber
i. = *Theridion ovale* Wunderlich, 1988 [preoccupied]
- † ***Thyelia* C. L. Koch & Berendt, 1854** **Palaeogene**
508. *Thyelia anomala* C. L. Koch & Berendt, 1854 Pa Baltic amber
509. *Thyelia convexa* C. L. Koch & Berendt, 1854 Pa Baltic amber
510. *Thyelia fossula* C. L. Koch & Berendt, 1854 Pa Baltic amber
511. *Thyelia marginata* C. L. Koch & Berendt, 1854 Pa Baltic amber
512. *Thyelia pallida* C. L. Koch & Berendt, 1854 Pa Baltic amber
513. *Thyelia scotina* C. L. Koch & Berendt, 1854 Pa Baltic amber
514. *Thyelia tristis* C. L. Koch & Berendt, 1854* Pa Baltic amber
515. *Thyelia villosa* C. L. Koch & Berendt, 1854 Pa Baltic amber
- Ulesanis* L. Koch, 1872** **Palaeogene – Recent**
516. *Ulesanis antecessor* Wunderlich, 2008b Pa Baltic Amber
517. *Ulesanis frontprocera* Wunderlich, 2008b Pa Baltic Amber
518. *Ulesanis longicymbium* Wunderlich, 2008b Pa Baltic Amber
519. *Ulesanis ovalis* Wunderlich, 2008b Pa Baltic / Bitt. amber
520. *Ulesanis parva* Wunderlich, 2008b Pa Baltic / Bitt. amber
- † ***Unispinatoda* Wunderlich, 2008b** **Palaeogene**
521. *Unispinatoda aculeata* Wunderlich, 2008b* Pa Baltic / Bitt. Amber
- † ***Vicipholcomma* Wunderlich, 2008b** **Palaeogene**
522. *Vicipholcomma spiralis* Wunderlich, 2008b* Pa Baltic Amber
- Theridiidae incertae sedis**
523. '*Eomysmena*' *succini* (Petrunkevitch, 1942) Pa Baltic amber
524. '*Anelosimus*' *clypeatus* Wunderlich, 1988 Ne Dominican amber
- THERIDIOSOMATIDAE Simon, 1881** **Cretaceous – Recent**
- Theridiosomatidae gen. et sp. indet *in* Wunderlich (2004*j*) Pa Baltic amber
- Theridiosomatidae gen. et sp. indet *in* Wunderlich (2011*f*) Qt Madagascar copal
- Baalzebub* Coddington, 1986** **?Cretaceous – Recent**
525. ?*Baalzebub mesozoicum* Penney, 2014 K Vendée amber
- † ***Eocoddingtonia* Selden, 2010** **Cretaceous**
526. *Eocoddingtonia eskovi* Selden, 2010* K Baissa, Transbaikalia
- † ***Eoepeirotypus* Wunderlich, 2004*j*** **Palaeogene**
527. *Eoepeirotypus retrobulbus* Wunderlich, 2004*j** Pa Baltic amber
- Eoepeirotypus* sp. *in* Wunderlich (2004) Pa Bitterfeld amber
- † ***Eotheridiosoma* Wunderlich, 2004*j*** **Palaeogene**
528. ?*Eotheridiosoma hamatum* Wunderlich, 2011*e* Pa Baltic amber
529. *Eotheridiosoma tuber* Wunderlich, 2004*j** Pa Bitterfeld amber
530. *Eotheridiosoma volutum* Wunderlich, 2004*j* Pa Bitterfeld amber

- † *Hypotheridiosoma* Wunderlich, 2012d Cretaceous
 531. *Hypotheridiosoma paracymbium* Wunderlich, 2012d* K Myanmar amber
- † *Leviunguis* Wunderlich, 2012d Cretaceous
 532. *Leviunguis bruckschi* Wunderlich, 2012d* K Myanmar amber
- † *Palaeoepirotypus* Wunderlich, 1988 Neogene
 533. *Palaeoepirotypus iuvenis* Wunderlich, 1988* Ne Dominican amber
 534. *Palaeoepirotypus iuvenoides* Wunderlich, 1988 Ne Dominican amber
- † *Spinitheridiosoma* Wunderlich, 2004j Palaeogene
 NB: type species designated from the wrong genus!
 535. *Spinitheridiosoma balticum* Wunderlich, 2004j Pa Baltic amber
 536. *Spinitheridiosoma bispinosum* Wunderlich, 2004j Pa Bitterfeld amber
 537. *Spinitheridiosoma rima* Wunderlich, 2004j Pa Baltic amber
- Theridiosoma* O. P.-Cambridge, 1879b Neogene – Recent
 538. *Theridiosoma incompletum* Wunderlich, 1988 Ne Dominican amber
- † *Umerosoma* Wunderlich, 2004j Palaeogene
 539. *Umerosoma multispina* Wunderlich, 2004j* Pa Baltic amber
- SYMPHYTOGNATHIDAE Hickman, 1931** Recent
 no fossil record
- ANAPIDAE Simon, 1895** Palaeogene – Recent
 = TEXTRICELLIDAE Hickman, 1945
- † *Balticonopsis* Wunderlich, 2004k Palaeogene
 540. *Balticonopsis bispina* Wunderlich, 2004k Pa Baltic amber
 541. *Balticonopsis bitterfeldensis* Wunderlich, 2004k Pa Bitterfeld amber
 542. *Balticonopsis bulbosa* Wunderlich, 2004k Pa Baltic amber
 543. *Balticonopsis ceranowiczae* Wunderlich, 2004k Pa Baltic amber
 544. *Balticonopsis holti* Wunderlich, 2004k* Pa Baltic amber
 545. *Balticonopsis perkovskyi* Wunderlich, 2004ar Pa Rovno amber
 546. *Balticonopsis thomasi* Wunderlich, 2004k Pa Baltic amber
Balticonopsis sp. in Wunderlich (2004k) Pa Baltic amber
- † *Dubianapis* Wunderlich, 2004k Palaeogene
 547. *Dubianapis obscura* Wunderlich, 2004k* Pa Baltic amber
- † *Flagellanapis* Wunderlich, 2004k Palaeogene
 548. *Flagellanapis voigti* Wunderlich, 2004k* Pa Baltic/Bitt. Amber
- † *Fossilanapis* Wunderlich, 2004k Palaeogene
 549. *Fossilanapis anderseri* Wunderlich, 2004k Pa Baltic amber
 550. *Fossilanapis baetcheri* Wunderlich, 2004k* Pa Baltic amber
 551. *Fossilanapis eichmanni* Wunderlich, 2004k Pa Baltic amber
 552. *Fossilanapis flexiotarsus* Wunderlich, 2004k Pa Baltic amber
 553. *Fossilanapis multispinae* Wunderlich, 2011h Pa Baltic amber
 554. *Fossilanapis saltans* Wunderlich, 2004k Pa Baltic amber

555. *Fossilanapis unispinum* Wunderlich, 2004k Pa Baltic amber
Fossilanapis sp. in Wunderlich (2004k) Pa Bitterfeld amber
Fossilanapis sp. in Wunderlich (2011h) Pa Baltic amber
- † ***Palaeoanapis* Wunderlich, 1988** Neogene
556. *Palaeoanapis nana* Wunderlich, 1988* Ne Dominican amber
- † ***Ruganapis* Wunderlich, 2004k** Palaeogene
557. *Ruganapis scutata* Wunderlich, 2004k* Pa Baltic amber
- † ***Saxonanapis* Wunderlich, 2004k** Palaeogene
558. *Saxonanapis grabenhorsti* Wunderlich, 2004k* Pa Baltic/Bitt. Amber
- † ***Tuberanapis* Wunderlich, 2004k** Palaeogene
559. *Tuberanapis parvibulbus* Wunderlich, 2004k* Pa Baltic amber
- COMAROMIDAE Wunderlich, 2004** [stat. nov. 2011] Palaeogene – Recent
- † ***Balticorma* Wunderlich, 2004k** Palaeogene
= † *Balticorma* [sic] Weitschat & Wichard, 2002 [nomen nudum]
560. *Balticorma damzeni* Wunderlich, 2011h Pa Baltic amber
561. *Balticorma ernstorom* Wunderlich, 2004k Pa Baltic/Bitt. amber
562. *Balticorma gracilipes* Wunderlich 2004k Pa Baltic/Bitt. amber
563. *Balticorma reschi* Wunderlich, 2004k* Pa Baltic amber
564. *Balticorma serafinorum* Wunderlich, 2004k Pa Baltic/Bitt. amber
565. *Balticorma tibialis* Wunderlich, 2004k Pa Baltic amber
566. *Balticorma wheateri* Penney & Marusik in Penney et al. (2011) Pa Baltic amber
- MYSMENIDAE Petrunkevitch, 1928** Palaeogene – Recent
- Mysmeninae sp. in Wunderlich (2004ar) Pa Rovno amber
- † ***Dominicanopsis* Wunderlich, 2004k** Neogene
567. *Dominicanopsis grimaldii* Wunderlich, 2004k* Ne Dominican amber
- † ***Eomysmenopsis* Wunderlich, 2004k** Palaeogene
568. *Eomysmenopsis spinipes* Wunderlich, 2004k* Pa Baltic / Bitt. Amber
- Mysmena* Simon, 1894** Palaeogene – Recent
- Mysmena* (s. l.) sp. indet in Wunderlich (2012a) Qt Madagascan copal
569. *Mysmena* (s.l.) *copalis* Wunderlich, 2011f Qt Madagascan copal
570. *Mysmena curvata* Wunderlich, 2011h Pa Baltic amber
571. *Mysmena dominicana* Wunderlich, 1998 Qt Madagascan copal
572. *Mysmena fossilis* Petrunkevitch, 1971 Ne Chiapas amber
573. *Mysmena groehni* Wunderlich, 2004k Pa Baltic / Bitt. amber
574. *Mysmena grotae* Wunderlich, 2004k Pa Baltic amber
- Mysmenopsis* Simon, 1897b** Neogene – Recent
575. *Mysmenopsis lissycoleyae* Penney, 2000 Ne Dominican amber
- † ***Palaeomysmena* Wunderlich, 2004k** Palaeogene
576. *Palaeomysmena hoffeinsorum* Wunderlich, 2004k* Pa Baltic amber

† BALTSUCCINIDAE Wunderlich, 2004/	Palaeogene
† Baltsuccinus Wunderlich, 2004/	Palaeogene
577. <i>Baltsuccinus flagellaceus</i> Wunderlich, 2004/ [*]	Pa Baltic amber
578. <i>Baltsuccinus similis</i> Wunderlich, 2004/	Pa Baltic amber
† PROTHERIDIIDAE Wunderlich, 2004/	Cretaceous – Palaeo.
† Protheridion Wunderlich, 2004/	Palaeogene
579. <i>Protheridion bitterfeldensis</i> Wunderlich, 2004/	Pa Bitterfeld amber
580. <i>Protheridion detritus</i> Wunderlich, 2004/	Pa Baltic amber
581. <i>Protheridion obscurum</i> Wunderlich, 2004/	Pa Baltic amber
582. <i>Protheridion punctatum</i> Wunderlich, 2004/	Pa Baltic amber
583. <i>Protheridion tibialis</i> Wunderlich, 2004/ [*]	Pa Baltic amber
† Zarqaraneus Wunderlich, 2008d	Cretaceous
584. <i>Zarqaraneus hudei</i> Wunderlich, 2008d [*]	K Jordanian amber
† PRAETHERIDIIDAE Wunderlich, 2004/ (n. stat. 2012)	Palaeogene
† Praetheridion Wunderlich, 2004/	Palaeogene
585. <i>Praetheridion fleissneri</i> Wunderlich, 2004/ [*]	Pa Baltic amber
SYNAPHRIDAE Wunderlich, 1986	Palaeogene – Recent
† Iardinidis Wunderlich 2004k	Palaeogene
586. <i>Iardinidis brevipes</i> Wunderlich, 2004k [*]	Pa Baltic amber
PIMOIDAE Wunderlich, 1986	Palaeogene – Recent
Pimoa Chamberlin & Ivie, 1943	Palaeogene – Recent
587. <i>Pimoa expandens</i> Wunderlich, 2004r	Pa Baltic amber
588. <i>Pimoa (Eopimoa) hormigai</i> Wunderlich, 2004r	Pa Baltic amber
589. <i>Pimoa inopinata</i> Wunderlich, 2004r	Pa Baltic amber
590. <i>Pimoa liedtkei</i> Wunderlich, 2004r	Pa Baltic amber
591. <i>Pimoa lingua</i> Wunderlich, 2004r	Pa Baltic amber
592. <i>Pimoa (Eopimoa) longiscapus</i> Wunderlich, 2008a	Pa Baltic amber
593. <i>Pimoa multicuspuli</i> Wunderlich, 2004r	Pa Baltic amber
594. <i>Pimoa (Eopimoa) obruens</i> Wunderlich, 2008a	Pa Baltic amber
<i>Pimoa</i> sp. in Wunderlich (2004r)	Pa Baltic amber
<i>Pimoa (Eopimoa)</i> sp. in Wunderlich (2008a)	Pa Baltic amber
PUMILIOPIMOIDAE Wunderlich, 2008a	Palaeogene – Recent
† Pumiliopimoa Wunderlich, 2008a	Palaeogene
595. <i>Pumiliopimoa parma</i> Wunderlich, 2008a [*]	Pa Baltic amber
SINOPIMOIDAE Li & Wunderlich, 2008	Recent
no fossil record	

LINYPHIIDAE Blackwall, 1859	Cretaceous – Recent
= MICRYPHANTIDAE Bertkau, 1878a	
= ERIGONIDAE Simon, 1884c	
?Linyphiidae gen. et sp. indet <i>in</i> McAlpine & Martin (1969)	K Canadian amber
Linyphiidae gen. et sp. indet <i>in</i> Penney (2002)	K New Jersey amber
Linyphiidae gen. et sp. indet <i>in</i> Schmidt <i>et al.</i> (2010)	K Ethiopian amber
Linyphiinae gen. et sp. indet <i>in</i> Penney & Selden (2002)	K Lebanese amber
[NB: Wunderlich (2012d) questioned the veracity of these Cretaceous linyphiids.]	
† <i>Agynetiphantes</i> Wunderlich, 2004s	Palaeogene
596. <i>Agynetiphantes gibbiferus</i> Wunderlich, 2004s*	Pa Baltic amber
<i>Ceratinopsis</i> Emerton, 1882	Quaternary – Recent
597. <i>Ceratinopsis deformans</i> (Wunderlich, 1998)	Qt Madagascan copal
<i>Cnephalcotes</i> Simon, 1884c	Quaternary – Recent
598. <i>Cnephalcotes obscurus</i> (Blackwall, 1834b) [Recent]	Qt England
† <i>Custodela</i> Petrunkevitch, 1942	Palaeogene
= † <i>Obnisis</i> Petrunkevitch, 1942 [tentative synonymy]	
599. <i>Custodela acuta</i> Wunderlich, 2004s	Pa Baltic amber
600. <i>Custodela acutula</i> Wunderlich, 2004s	Pa Bitterfeld amber
601. <i>Custodela bispina</i> Wunderlich, 2004s	Pa Bitterfeld amber
602. <i>Custodela bispinosa</i> Wunderlich, 2004s	Pa Bitterfeld amber
603. <i>Custodela cheiracantha</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
604. <i>Custodela clava</i> Wunderlich, 2004s	Pa Baltic amber
605. <i>Custodela curva</i> Wunderlich, 2004s	Pa Baltic amber
606. <i>Custodela curvata</i> Wunderlich, 2004s	Pa Bitterfeld amber
607. <i>Custodela divergens</i> Wunderlich, 2004s	Pa Baltic amber
608. <i>Custodela expandens</i> Wunderlich, 2004s	Pa Baltic amber
609. <i>Custodela falcata</i> Wunderlich, 2004s	Pa Baltic amber
610. <i>Custodela femurspinosa</i> Wunderlich, 2004s	Pa Bitterfeld amber
611. <i>Custodela henningseni</i> Wunderlich, 2004s	Pa Baltic amber
612. <i>Custodela kochi</i> Wunderlich, 2004s	Pa Baltic amber
613. <i>Custodela lamellata</i> (Wunderlich, 1988)	Pa Baltic amber
614. <i>Custodela lanx</i> Wunderlich, 2004s	Pa Baltic amber
615. <i>Custodela oblonga</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
616. <i>Custodela obtusa</i> Wunderlich, 2004s	Pa Baltic amber
617. ? <i>Custodela parva</i> Wunderlich, 2004s	Pa Bitterfeld amber
618. <i>Custodela pseudokochi</i> Wunderlich, 2004s	Pa Baltic amber
619. <i>Custodela stridulans</i> Wunderlich, 2004s	Pa Bitterfeld amber
620. <i>Custodela tenuipes</i> (Petrunkevitch, 1942)	Pa Baltic amber
621. <i>Custodela tibialis</i> Wunderlich, 2004s	Pa Baltic amber
<i>Custodela</i> sp. <i>in</i> Wunderlich (2004s)	Pa Bitterfeld amber
† <i>Custodela</i> Wunderlich, 2004s	Palaeogene

622.	<i>Custodelela hamata</i> Wunderlich, 2004s*	Pa	Bitterfeld amber
†	<i>Eolabulla</i> Wunderlich, 2004s		Palaeogene
623.	<i>Eolabulla falcata</i> Wunderlich, 2004s	Pa	Baltic amber
624.	<i>Eolabulla gladiiformis</i> Wunderlich, 2004s	Pa	Baltic amber
625.	<i>Eolabulla laminata</i> Wunderlich, 2004s*	Pa	Baltic amber
626.	<i>Eolabulla perforata</i> Wunderlich, 2004s	Pa	Baltic amber
627.	<i>Eolabulla sagitta</i> Wunderlich, 2004s	Pa	Baltic amber
628.	<i>Eolabulla similis</i> Wunderlich, 2004s	Pa	Baltic amber
	<i>Eolabulla</i> sp. 1–2 in Wunderlich (2004s)	Pa	Baltic amber
†	<i>Eophantes</i> Wunderlich, 2004s		Palaeogene
629.	<i>Eophantes complicatus</i> Wunderlich, 2004s*	Pa	Baltic amber
630.	? <i>Eophantes seorsum</i> Wunderlich, 2012c	Pa	Baltic amber
<i>Erigone</i> Audouin, 1826			Neogene – Recent
	<i>Erigone</i> sp. in Hopkins et al. (1976)	Qt	Alaska
631.	<i>Erigone atra</i> Blackwall, 1833 [Recent]	Qt	England
632.	? <i>Erigone dechenii</i> Bertkau, 1878b	Ne	Rott, Germany
<i>Floricomus</i> Crosby & Bishop, 1925			Neogene – Recent
633.	<i>Floricomus fossilis</i> Penney, 2005c	Ne	Dominican amber
<i>Gonatium</i> Menge, 1868			Quaternary – Recent
634.	<i>Gonatium rubens</i> (Blackwall, 1833) [Recent]	Qt	England
<i>Hypselistes</i> Simon, 1894			Quaternary – Recent
635.	<i>Hypselistes jacksoni</i> (O. P.-Cambridge, 1902) [Recent]	Qt	England
<i>Linyphia</i> Latreille, 1804a			Palaeogene – Recent
636.	<i>Linyphia andraei</i> Bertkau, 1878b	Ne	Rott, Germany
637.	<i>Linyphia byrami</i> Cockerell, 1925	Pa	Green River
638.	<i>Linyphia florissanti</i> Petrunkevitch, 1922	Pa	Florissant
639.	<i>Linyphia pachygnathoides</i> Petrunkevitch, 1922	Pa	Florissant
640.	<i>Linyphia quievreuxi</i> Berland, 1939	Pa	Aix-en-Provence
641.	<i>Linyphia retensa</i> Scudder, 1890a	Pa	Florissant
642.	<i>Linyphia rottensis</i> Bertkau, 1878b	Ne	Rott, Germany
643.	<i>Linyphia seclusa</i> (Scudder, 1890a)	Pa	Florissant
† <i>Madagascarphantes</i> Wunderlich, 2012a			Quaternary
644.	<i>Madagascarphantes vomerans</i> Wunderlich, 2012a*	Qt	Madagascan copal
† <i>Malepellis</i> Petrunkevitch, 1971			Neogene
645.	<i>Malepellis extincta</i> Petrunkevitch, 1971*	Ne	Chiapas amber
<i>Meioneta</i> Hull, 1920			Neogene – Recent
646.	<i>Meioneta bigibber</i> (Wunderlich, 1988)	Ne	Dominican amber
647.	<i>Meioneta fastigata</i> (Wunderlich, 1988)	Ne	Dominican amber
648.	<i>Meioneta separata</i> (Wunderlich, 1988)	Ne	Dominican amber
	<i>Meioneta</i> sp. in Wunderlich (1988)	Ne	Dominican amber
<i>Micryphantes</i> C. L. Koch, 1833			Palaeogene

649. *Micryphantes molybdinus* C. L. Koch & Berendt, 1854 Pa Baltic amber
650. *Micryphantes regularis* C. L. Koch & Berendt, 1854 Pa Baltic amber
- † ***Mystagogus* Petrunkevitch, 1942** ...[Wunderlich suggests possibly in Cyatholipidae] **Palaeogene**
651. *Mystagogus dubius* Petrunkevitch, 1958 Pa Baltic amber
652. *Mystagogus glaber* Petrunkevitch, 1942* Pa Baltic amber
- † ***Paralabulla* Wunderlich, 2004s** **Palaeogene**
653. *Paralabulla bitterfeldensis* Wunderlich, 2004s* Pa Bitterfeld amber
654. ?*Paralabulla dubia* Wunderlich, 2004s Pa Baltic amber
655. *Paralabulla succinifera* Wunderlich, 2004s Pa Baltic amber
- Paralabulla* sp. in Wunderlich (2004s, 2012c) Pa Bitterfeld amber
- Pocadicnemis* Simon, 1884c** **Quaternary – Recent**
656. *Pocadicnemis pumila* (Blackwall, 1841) **[Recent]** Qt England
- Savignia* Blackwall, 1833** **Quaternary – Recent**
657. *Savignia frontata* Blackwall, 1833 **[Recent]** Qt England
- Selenyphantes* Gertsch & Davis, 1946** **Neogene – Recent**
- = † *Palaeolinypbia* Wunderlich, 1986
658. *Selenyphantes flagellifera* (Wunderlich, 1986) Ne Dominican amber
- † ***Succineta* Wunderlich, 2004s** **Palaeogene**
659. *Succineta brevispina* Wunderlich, 2004s Pa Baltic amber
660. *Succineta discoidalis* Wunderlich, 2004s* Pa Baltic amber
- Succineta* sp. in Wunderlich (2004s) Pa Baltic amber
- † ***Succiphantes* Wunderlich, 2004s** **Palaeogene**
661. *Succiphantes tanasevitchi* Wunderlich, 2004s Pa Baltic amber
662. *Succiphantes velteni* Wunderlich, 2004s* Pa Baltic amber
- Toschia* Caporiacco, 1949** **Quaternary – Recent**
663. ?*Toschia fossilis* Wunderlich, 2004as Qt Madagascan copal
- TETRAGNATHIDAE Menge, 1866** **Cretaceous – Recent**
- = PACHYGNATHIDAE Menge, 1866
- = METIDAE Simon, 1894
- = NANOMETIDAE Forster & Forster, 1999
- † ***Anameta* Wunderlich, 2004h** **Palaeogene**
664. *Anameta distenda* Wunderlich, 2004h* Pa Bitterfeld amber
665. *Anameta kuntneri* Wunderlich, 2008a Pa Baltic amber
- Azilia* Keyserling, 1882** **Neogene – Recent**
666. *Azilia hispaniolensis* Wunderlich, 1988 Ne Dominican amber
- i. = *Azilia muellenmeisteri* Wunderlich, 1988 Ne Dominican amber
- Azilia* sp. in Wunderlich (1988) Ne Dominican amber
- † ***Balticgnatha* Wunderlich, 2011h** **Palaeogene**
667. *Balticgnatha projectens* Wunderlich 2011h* Pa Baltic amber
- † ***Battleucauge* Wunderlich, 2008a** **Palaeogene**
668. *Battleucauge gillespieae* Wunderlich 2008a* Pa Baltic amber

669. *Baltleuauge propinqua* Wunderlich, 2012c Pa Baltic amber
- † ***Corneometa*** Wunderlich, 2004h Palaeogene
670. *Corneometa baltica* Wunderlich 2004h* Pa Baltic amber
671. *Corneometa pilosipes* Wunderlich 2004h Pa Baltic amber
- Cyrtognatha*** Keyserling, 1882 Neogene – Recent
672. *Cyrtognatha weitschati* Wunderlich, 1988 Ne Dominican amber
- † ***Eometa*** Petrunkevitch, 1958 Palaeogene
673. *Eometa calefacta* Wunderlich, 2004h Pa Baltic amber
674. *Eometa longipes* Petrunkevitch, 1958 Pa Baltic amber
675. *Eometa occulta* Wunderlich, 2004h Pa Baltic amber
676. *Eometa perfecta* Wunderlich, 2004h Pa Baltic amber
677. *Eometa samlandica* Petrunkevitch, 1958* Pa Baltic amber
- Eometa* sp. 1–2 in Wunderlich (2004h) Pa Baltic amber
- Homalometra*** Simon, 1897b Neogene – Recent
678. *Homalometra fossilis* Wunderlich, 1988 Ne Dominican amber
- † ***Huergina*** Selden & Penney, 2003 Cretaceous
679. *Huergina diazromeralei* Selden & Penney, 2003* K Las Hoyas, Spain
- † ***Macryphantes*** Selden, 1990 Cretaceous
680. *Macryphantes cowdeni* Selden, 1990* K Sierra de Montsech
- Meta*** C. L. Koch, 1836 Palaeogene – Recent
681. *Meta (Praetermeta) maculosa* Wunderlich, 2008a Pa Baltic amber
682. *Meta (Praetermeta) velans* (Wunderlich, 2004h) Pa Baltic amber
- † ***Palaeometa*** Petrunkevitch, 1922 Palaeogene
683. *Palaeometa opertanea* (Scudder, 1890a)* Pa Florissant
- † ***Palaeopachygnatha*** Petrunkevitch, 1922 Palaeogene
684. *Palaeopachygnatha cockerelli* Petrunkevitch, 1922 Pa Florissant
685. *Palaeopachygnatha scudderri* Petrunkevitch, 1922* Pa Florissant
- † ***Priscometa*** Petrunkevitch, 1958 Palaeogene
686. *Priscometa capta* Wunderlich, 2004h Pa Baltic amber
687. *Priscometa minor* Wunderlich, 2004h Pa Baltic amber
688. *Priscometa tenuipes* Petrunkevitch, 1958* Pa Baltic amber
- † ***Samlandicmeta*** Wunderlich, 2012c Palaeogene
689. *Samlandicmeta mutila* Wunderlich, 2012c Pa Baltic amber
- Tetragnatha*** Latreille, 1804a Palaeogene – Recent
690. *Tetragnatha parva* (Hong, 1985) Ne Shanwang
691. *Tetragnatha pristina* Schawaller, 1982c Ne Dominican amber
692. *Tetragnatha tertaria* Scudder, 1885 Pa Florissant
- NEPHILIDAE** Simon, 1894 Jurassic – Recent
- Nephilidae indet. in Wunderlich (2012c) Pa Baltic amber
- † ***Cretaraneus*** Selden, 1990 Cretaceous

693. *Cretaraneus liaoningensis* Cheng, Meng & Wang *in Cheng et al.*,
2008 K Jehol biota
694. *Cretaraneus martensnetoi* Mesquita, 1996 K Crato Formation
695. *Cretaraneus vilaltae* Selden, 1990* K Sierra de Montsech
- † *Eonephila* Wunderlich, 2004*i* Palaeogene
696. *Eonephila bitterfeldensis* Wunderlich, 2004*i* Pa Bitterfeld amber
697. *Eonephila excellens* Wunderlich, 2004**i* Pa Baltic amber
698. *Eonephila longembolus* Wunderlich, 2004*i* Pa Baltic amber
- † *Geratonephila* Poinar *in* Poinar & Buckley, 2012 Cretaceous
699. *Geratonephila burmanica* Poinar *in* Poinar & Buckley, 2012* K Myanmar amber
- † *Luxurioneephila* Wunderlich, 2004*i* Palaeogene
700. *Luxurioneephila spinifera* Wunderlich, 2004*i* Pa Baltic amber
- † *Minutunguis* Wunderlich, 2011*f* Quaternary
701. *Minutunguis silvestris* Wunderlich, 2011*f** Qt Madagascar copal
- Nephila* Leach, 1815 Cretaceous – Recent
702. *Nephila breviembolus* Wunderlich, 1986 Ne Dominican amber
703. *Nephila dommeli* Wunderlich, 1982 Ne Dominican amber
704. *Nephila furca* Wunderlich, 1986 Ne Dominican amber
705. *Nephila longembolus* Wunderlich, 1986 Ne Dominican amber
706. *Nephila pennatipes* Scudder, 1885 Pa Florissant
707. *Nephila tenuis* Wunderlich, 1986 Ne Dominican amber
- Nephila* sp. *in* Dunlop & Penney (2012) K Crato Formation
- † *Palaeonephila* Wunderlich, 2004*i* Palaeogene
708. *Palaeonephila brevis* Wunderlich, 2004*i* Pa Baltic amber
709. *Palaeonephila curvata* Wunderlich, 2004**i* Pa Baltic amber
710. *Palaeonephila dilitans* Wunderlich, 2004*i* Pa Baltic amber
711. *Palaeonephila fibula* Wunderlich, 2004*i* Pa Baltic amber
712. *Palaeonephila longipes* Wunderlich, 2004*i* Pa Baltic amber
- † **MONGOLARACHNIDAE** Selden, Shi & Ren, 2013 Jurassic
- † *Mongolarachne* Selden, Shi & Ren, 2013 Jurassic
713. *Mongolarachne jurassica* (Selden, Shih & Ren, 2011)* J Daohugou
- † **JURARANEIDAE** Eskov, 1984 Jurassic
- † *Juraraneus* Eskov, 1984 Jurassic
714. *Juraraneus rASNITSYNI* Eskov, 1984 J Transbaikalia
- ARANEIDAE** Simon, 1895 Cretaceous – Recent
- = EPEIRIDAE Sundevall, 1833 [based on a generic synonym]
- = EUETRIIDAE Thorell, 1887 [based on a generic synonym]
- = ARGIOPIDAE Simon, 1890
- = ZYGIELLIDAE Simon, 1929

- ?Araneinae sp. *in* Wunderlich (2004*h*) Pa Baltic amber
- Araneidae gen. et sp. indet. *in* Ribera (2003) Qt Girona, Spain
- ?Mangorini indet. *in* Wunderlich (2011*a*) Pa Baltic amber
- Araneidae incertae sedis *in* Selden (2014*b*) Pa Isle of Wight
- † **Anepeira** Wunderlich, 2004*i* **Palaeogene**
715. *Anepeira complicata* Wunderlich, 2004*†* Pa Baltic amber
- † **Araneometa** Wunderlich, 1988 **Neogene**
716. *Araneometa excelsa* Wunderlich, 1988 Ne Dominican amber
717. *Araneometa herringi* Wunderlich, 1988* Ne Dominican amber
718. *Araneometa spirembolus* Wunderlich, 1988 Ne Dominican amber
- Araneometa* sp. *in* Wunderlich (1988) Ne Dominican amber
- Araneus** Clerck, 1757 **?Cretaceous – Recent**
719. ?*Araneus* sp. *in* Wunderlich (2012*c*) Pa Baltic amber
720. *Araneus absconditus* (Scudder, 1890*a*) Pa Florissant
721. *Araneus aethus* Chang, 2004 [generic assignment unreliable!] K Jehol biota
722. *Araneus beipiaoensis* Chang, 2004 [generic assignment unreliable!] K Jehol biota
723. *Araneus carbonaceous* Zhang, Sun & Zhang, 1994 Ne Shanwang
724. *Araneus cinefactus* (Scudder, 1890*a*) Pa Florissant
725. *Araneus defunctus* Petrunkevitch, 1958 Pa Baltic amber
726. *Araneus delitus* (Scudder, 1890*a*) Pa Florissant
727. *Araneus emertoni* (Scudder, 1890*a*) Pa Florissant
728. *Araneus exustus* Petrunkevitch, 1963 Ne Chiapas amber
729. *Araneus kinchloeae* Dunlop & Jekel, 2009 Pa Florissant
- i. = *Araneus indistinctus* (Petrunkevitch, 1922) [preoccupied]
730. *Araneus inelegans* Zhang, Sun & Zhang, 1994 Ne Shanwang
731. *Araneus leptopodus* Zhang, Sun & Zhang, 1994 Ne Shanwang
732. *Araneus liaoxiensis* Chang, 2004 [generic assignment unreliable!] K Jehol biota
733. *Araneus longimanus* (Petrunkevitch, 1922) Pa Florissant
734. *Araneus (Calinurus) longipes* Dalman, 1826 Qt Copal
735. *Araneus luanus* Zhang, Sun & Zhang, 1994 Ne Shanwang
736. *Araneus meeki* (Scudder, 1890*a*) Pa Florissant
737. *Araneus molassicus* (Heer, 1865) Ne Öhningen
738. *Araneus nanus* Wunderlich, 1988 Ne Dominican amber
739. *Araneus piceus* Lin, Zhang & Wang, 1989 Ne Shanwang
740. *Araneus reheensis* Chang, 2004 [generic assignment unreliable!] K Jehol biota
741. *Araneus ruidipedalis* Zhang, Sun & Zhang, 1994 Ne Shanwang
742. *Araneus troschelii* (Bertkau, 1878*b*) Ne Rott, Germany
743. *Araneus vulcanalis* (Scudder, 1890*a*) Pa Florissant
- Argiope** Audouin, 1826 **Neogene – Recent**
- = † *Magnaranea* Hong, 1985
744. *Argiope furva* (Hong, 1985) Ne Shanwang

† <i>Bararaneus</i> Wunderlich, 2004i	Palaeogene
745. ? <i>Bararaneus annulatus</i> Wunderlich, 2004i	Pa Baltic amber
746. <i>Bararaneus evolvens</i> Wunderlich, 2004*	Pa Baltic amber
† <i>Chrysometata</i> Wunderlich, 2004h	Palaeogene
747. <i>Chrysometata palaearctica</i> Wunderlich, 2004h*	Pa Baltic amber
† <i>Cyclososoma</i> Petrunkevitch, 1958	Palaeogene
748. <i>Cyclososoma succini</i> Petrunkevitch, 1958*	Pa Baltic amber
<i>Enacrosoma</i> Mello-Leitão, 1932	Neogene – Recent
749. <i>Enacrosoma verrucosa</i> (Wunderlich, 1988)	Ne Dominican amber
† <i>Eoaraneus</i> Wunderlich, 2004i	Palaeogene
750. <i>Eoaraneus complexus</i> Wunderlich, 2004*	Pa Baltic amber
† <i>Eochorizopes</i> Wunderlich, 2008a	Palaeogene
751. <i>Eochorizopes szeklinskiae</i> Wunderlich, 2008a*	Pa Baltic amber
† <i>Eozygiella</i> Wunderlich, 2004h	Palaeogene
752. <i>Eozygiella compacta</i> Wunderlich, 2004h*	Pa Baltic amber
† <i>Fossilaraneus</i> Wunderlich, 1988	Neogene
753. <i>Fossilaraneus incertus</i> Wunderlich, 1988*	Ne Dominican amber
<i>Gea</i> C. L. Koch, 1843a	Palaeogene – Recent
754. <i>Gea krantzi</i> von Heyden, 1859	Ne Rott, Germany
† <i>Graea</i> Thorell, 1869	Palaeogene
= † <i>Eustaloides</i> Petrunkevitch, 1942	
755. ? <i>Graea aberrans</i> Wunderlich, 2004h	Pa Baltic amber
756. <i>Graea bitterfeldensis</i> Wunderlich, 2004h	Pa Bitterfeld amber
757. <i>Graea breviembolus</i> Wunderlich, 2004h	Pa Baltic amber
758. <i>Graea brevis</i> Wunderlich, 2004h	Pa Baltic amber
759. <i>Graea calceatus</i> (Petrunkevitch, 1950)	Pa Baltic amber
760. <i>Graea epeiroidea</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
761. <i>Graea impudica</i> Wunderlich, 2004h	Pa Baltic amber
762. <i>Graea lingula</i> Wunderlich, 2004h	Pa Baltic amber
763. <i>Graea magnocoli</i> Wunderlich, 2012c	Pa Baltic amber
764. <i>Graea minor</i> (Petrunkevitch, 1950)	Pa Baltic amber
765. <i>Graea setosa</i> (Petrunkevitch, 1942)	Pa Baltic amber
766. <i>Graea succini</i> Petrunkevitch, 1942	Pa Baltic amber
<i>Hypognatha</i> Guérin, 1839	Quaternary – Recent
767. <i>Hypognatha testudinaria</i> (Taczanowski, 1879) [Recent]	Qt Colombian copal
† <i>Meditrina</i> Petrunkevitch, 1942	Palaeogene
768. <i>Meditrina circumvallata</i> Petrunkevitch, 1942*	Pa Baltic amber
† <i>Mesozygiella</i> Penney & Ortuño, 2006	Cretaceous
769. <i>Mesozygiella dunlopi</i> Penney & Ortuño, 2006*	K Álava amber
† <i>Miraraneus</i> Wunderlich, 2004i	Palaeogene
770. <i>Miraraneus peregrinus</i> Wunderlich, 2004*	Pa Baltic amber

† <i>Mirometa</i> Petrunkevitch, 1963	Neogene
771. <i>Mirometa valdespinosa</i> Petrunkevitch, 1963	Ne Chiapas amber
<i>Molinaranea</i> Mello-Leitão, 1940	Neogene – Recent
772. <i>Molinaranea mitnickii</i> Saupe, Selden & Penney, 2010	Ne Dominican amber
† <i>Pycnosinga</i> Wunderlich, 1988	Neogene
773. <i>Pycnosinga fossilis</i> Wunderlich, 1988*	Ne Dominican amber
† <i>Testudinaroides</i> Dunlop & Jekel, 2008	Neogene
= † <i>Testudinaria</i> Zhang, Sun & Zhang, 1994 [preoccupied]	
774. <i>Testudinaroides papposa</i> (Zhang, Sun & Zhang, 1994)	Ne Shanwang
† <i>Tethneus</i> Scudder, 1885	Palaeogene
= † <i>Melanites</i> Hong, 1985	
775. <i>Tethneus guyoti</i> Scudder, 1890a	Pa Florissant
776. <i>Tethneus hentzi</i> Scudder, 1885*	Pa Florissant
777. <i>Tethneus obduratus</i> Scudder, 1890a	Pa Florissant
778. <i>Tethneus orbiculatus</i> (Hong, 1985)	Ne Shanwang
779. <i>Tethneus provectus</i> Scudder, 1890a	Pa Florissant
780. <i>Tethneus robustus</i> Petrunkevitch, 1922	Pa Florissant
781. <i>Tethneus twenhofeli</i> Petrunkevitch, 1922	Pa Florissant
<i>Zilla</i> C. L. Koch, 1834	Palaeogene – Recent
782. <i>Zilla gracilis</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
783. <i>Zilla porrecta</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
784. <i>Zilla veterana</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
RETROLATERAL TIBIAL APOPHYSIS CLADE	Cretaceous – Recent
?RTA-clade <i>in</i> Wunderlich (2008d)	K Myanmar amber
LYCOSOIDEA Sundevall, 1833	Cretaceous – Recent
† <i>Korearachne</i> Selden, Nam, Kim & Kim, 2012	Cretaceous
785. <i>Korearachne jinju</i> Selden, Nam, Kim & Kim, 2012*	K Sacheon, S. Korea
[Tentative assignment to Lycosoidea; disputed by Wunderlich (2012d) who suggested it could be a haplogyne spider in Pholcoidea or Leptonetoidea]	
LYCOSIDAE Sundevall, 1833	?Cretaceous – Recent
Lycosidae gen. et sp. <i>in</i> Bottali (1975)	Qt Italy
Lycosidae gen. et sp. <i>in</i> Schawaller (1982d)	Ne Willershausen
Lycosidae gen. et sp. <i>in</i> Penney (2001)	Ne Dominican amber
Lycosidae gen. et sp. <i>in</i> Kim & Nam (2012) [unreliable record]	K Lioyuan, China
<i>Alopecosa</i> Simon, 1885b	Quaternary – Recent
786. <i>Alopecosa ?pulverulenta</i> (Clerck, 1757) [Recent]	Qt England
† <i>Dryadia</i> Zhang, Sun & Zhang, 1994	Palaeogene
787. <i>Dryadia acanthopoda</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
<i>Lycosa</i> Latreille, 1804a	Palaeogene – Recent
788. <i>Lycosa florissanti</i> Petrunkevitch, 1922	Pa Florissant

789.	<i>Lycosa lithographica</i> Schawaller & Ono, 1979	Ne Randecker Maar
790.	<i>Lycosa malleata</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
791.	<i>Lycosa miocaena</i> Schawaller & Ono, 1979	Ne Randecker Maar
792.	<i>Lycosa subterranea</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
Pardosa C. L. Koch, 1847		Quaternary – Recent
793.	<i>Pardosa pullata</i> (Clerck, 1757) [Recent]	Qt England
	<i>Pardosa</i> sp. in Scott (2003)	Qt England
Pirata Sundevall, 1833		Quaternary – Recent
794.	<i>Pirata ?piraticus</i> (Clerck, 1757) [Recent]	Qt England
Trochosa C. L. Koch, 1847		Quaternary – Recent
795.	<i>Trochosa terricola</i> Thorell, 1856 [Recent]	Qt England
† PARATTIDAE Petrunkevitch, 1922		Palaeogene
† Parattus Petrunkevitch, 1922		Palaeogene
796.	<i>Parattus evocatus</i> (Scudder, 1890a)	Pa Florissant
797.	<i>Parattus latitatus</i> (Scudder, 1890a)	Pa Florissant
798.	<i>Parattus oculatus</i> Petrunkevitch, 1922	Pa Florissant
799.	<i>Parattus resurrectus</i> (Scudder, 1890a)*	Pa Florissant
TRECHALEIDAE Simon, 1890		Palaeogene – Recent
	= TRICLARIDAE O. P.-Cambridge, 1877 [<i>nomen oblitum</i>]	
	= PERISSOBLEMMATIDAE O. P.-Cambridge, 1882b [based on a synonym]	
Trechaleidae sp. in Wunderlich (2004aa)		Pa Baltic amber
† Eotrechalea Wunderlich, 2004aa		Palaeogene
800.	<i>Eotrechalea annulata</i> Wunderlich, 2004aa*	Pa Baltic amber
† Esuritor Petrunkevitch, 1942		Palaeogene
801.	<i>Esuritor aculeatus</i> Petrunkevitch, 1958	Pa Baltic amber
802.	<i>Esuritor spinipes</i> Petrunkevitch, 1942*	Pa Baltic amber
† Linoptes Menge, 1854		Palaeogene
803.	?'Linoptes' <i>oculeus</i> Menge in C. L. Koch & Berendt, 1854*	Pa Baltic amber
NB: <i>Linoptes</i> mentioned as a <i>nomen nudum</i> by Wunderlich (2004z); this species listed by Wunderlich (2004aa) under Trechaleidae and another species under Pisauridae (see below)		
PISAURIDAE Simon, 1890		Palaeogene – Recent
	= BRADYSTICHIDAE Simon, 1884	
	= DOLOMEDIDAE Simon, 1898a	
	= HALIDAE Jocqué, 1994	
Pisauridae sp. in Wunderlich (1988)		Pa Dominican amber
Pisauridae sp. in Wunderlich (2004z)		Pa Baltic amber
Dolomedes Latreille, 1804a		Quaternary – Recent
804.	<i>Dolomedes fimbriatus</i> (Clerck, 1757) [Recent]	Qt England
† 'Linoptes' Menge, 1854		Palaeogene

= † *Eopisaurella* Petrunkevitch, 1958

NB: See notes on *Linoptes* under Trechaleidae above!

805. ?'Linoptes' *valdespinosa* (Petrunkevitch, 1958)* Pa Baltic amber

?'Linoptes' sp. 1–8 in Wunderlich (2004z) Pa Baltic amber

† *Palaeoperenethis* Selden & Penney, 2009 Palaeogene

806. *Palaeoperenethis thaleri* Selden & Penney, 2009* Pa British Columbia

OXYOPIDAE Thorell, 1870a Palaeogene – Recent

= SPHASIDAE O. P.-Cambridge, 1871

= HAMATALIVIDAE Marx, 1890b

Oxyopidae sp. in Wunderlich 2004ab Pa Bitterfeld amber

Oxyopes Latreille, 1804a Palaeogene – Recent

807. *Oxyopes defectus* Wunderlich, 1988 Ne Dominican amber

808. 'Oxyopes' *succini* Petrunkevitch, 1958 Pa Baltic amber

Oxyopes sp. in Wunderlich (1988, 2004ab) Ne Dominican amber

† *Planoxyopes* Petrunkevitch, 1963 Neogene

809. *Planoxyopes eximius* Petrunkevitch, 1963* Ne Chiapas amber

i. = *Planoxyopes fossilis* Wunderlich, 1988 [lapsus] Ne Chiapas amber

SENOCULIDAE Simon, 1890 Recent

= NEOTHEREUTOIDAE Holmberg, 1883 [based on a generic synonym]

no fossil record

STIPHIDIIDAE Dalmas, 1917 Recent

no fossil record

ZOROCRATIDAE Dahl, 1913 Recent

no fossil record

PSECHRIDAE Simon, 1890 Recent

no fossil record

ZOROPSIDAE Bertkau, 1882 Palaeogene – Recent

Zoropsidae sp. in Wunderlich (2004x) Pa Baltic / Bitt. amber

† *Eomatachia* Petrunkevitch, 1942 Palaeogene

810. *Eomatachia barbarus* Wunderlich, 2004x Pa Baltic amber

811. *Eomatachia bipartita* Wunderlich, 2004x Pa Baltic amber

812. *Eomatachia divergens* Wunderlich, 2004x Pa Baltic amber

813. *Eomatachia duplex* Wunderlich, 2004x Pa Baltic amber

814. *Eomatachia latifrons* Petrunkevitch, 1942* Pa Baltic amber

815. *Eomatachia recedens* Wunderlich, 2004x Pa Baltic amber

816. *Eomatachia succini* (Petrunkevitch, 1942) Pa Baltic amber

817. *Eomatachia wegneri* Wunderlich, 2004x Pa Baltic amber

818. *Eomatachia xanthippe* Wunderlich, 2004x Pa Baltic amber
- † ***Eoprychia* Petrunkevitch, 1958** Palaeogene
819. *Eoprychia succini* Petrunkevitch, 1958* Pa Baltic amber
820. *Eoprychia succinopsis* Wunderlich, 2004x Pa Baltic amber
821. *Eoprychia vicina* Wunderlich, 2004x Pa Baltic amber
- Eoprychia* sp. in Wunderlich (2004x) ?Pa not specified
- † ***Succiniopsis* Wunderlich, 2004x** Palaeogene
822. *Succiniopsis kutscheri* Wunderlich, 2004x* Pa Baltic / Bitt. Amber
823. *Succiniopsis runcinata* Wunderlich, 2012c Pa Baltic amber
824. *Succiniopsis samlandica* Wunderlich, 2004x Pa Baltic amber
- † **INSECUTORIDAE Petrunkevitch, 1942** Palaeogene
- † ***Insecutor* Petrunkevitch, 1942** Palaeogene
825. *Insecutor aculeatus* Petrunkevitch, 1942* Pa Baltic amber
826. *Insecutor mandibulatus* Petrunkevitch, 1942 Pa Baltic amber
827. ?*Insecutor pecten* Wunderlich, 2004y Pa Baltic amber
828. *Insecutor rufus* Petrunkevitch, 1942 Pa Baltic amber
829. ?*Insecutor spinifer* Wunderlich, 2004y Pa Baltic amber
- Insecutor* sp. in Wunderlich (2004y) Pa Baltic amber
- † **SUCCINOMIDAE Wunderlich, 2012c** Palaeogene
- † ***Eohalinobius* Wunderlich, 2008c** Palaeogene
830. *Eohalinobius calefactus* Wunderlich, 2012c Pa Baltic amber
831. *Eohalinobius hiddenseeensis* Wunderlich, 2012c Pa Baltic amber
832. *Eohalinobius patina* Wunderlich, 2012c Pa Baltic amber
833. *Eohalinobius scutatus* Wunderlich, 2008c Pa Baltic amber
- † ***Succinomus* Wunderlich, 2008c** Palaeogene
834. *Succinomus duomammillae* Wunderlich, 2008c Pa Baltic amber
835. ?*Succinomus gibbosus* Wunderlich, 2012c Pa Baltic amber
- CTENIDAE Keyserling, 1877** Neogene – Recent
- = ACANTHOCTENIDAE Simon, 1892b
- † ***Nanoctenus* Wunderlich, 1988** Neogene
836. *Nanoctenus longipes* Wunderlich, 1988* Ne Dominican amber
- AGELENIDAE C. L. Koch, 1837** Palaeogene – Recent
- = TEGENARIDAE Prach, 1860
- = † INCEPTORIDAE Petrunkevitch, 1942
- Agelena* Walckenaer, 1805** Palaeogene – Recent
837. *Agelena tabida* C. L. Koch & Berendt, 1854 Pa Baltic amber
- Histopona* Thorell, 1869** Palaeogene – Recent
838. ?*Histopona anthracina* Bertkau, 1878b Ne Rott, Germany

† <i>Inceptor</i> Petrunkevitch, 1942	Palaeogene
839. <i>Inceptor aculeatus</i> Petrunkevitch, 1942*	Pa Baltic amber
840. <i>Inceptor dubius</i> Petrunkevitch, 1946	Pa Baltic amber
<i>Tegenaria</i> Latreille, 1804a	Palaeogene – Recent
841. ? <i>Tegenaria fragmentum</i> Wunderlich, 2004w	Pa Baltic amber
842. <i>Tegenaria lacazei</i> Gourret, 1887	Pa Aix-en-Provence
843. ? <i>Tegenaria obtusa</i> Wunderlich, 2004w	Pa Baltic amber
844. <i>Tegenaria virilis</i> Menge in C. L. Koch & Berendt, 1854	Pa Baltic amber
DICTYNOIDEA O. P.-Cambridge, 1871	Palaeogene – Recent
Dictynoidea incertae sedis	
† <i>Sinodictyna</i> Hong, 1982	Palaeogene
845. <i>Sinodictyna fushunensis</i> Hong, 1982*	Pa Fu Shun amber
CYBAEIDAE Simon, 1898a	Palaeogene – Recent
= ARGYRONETIDAE Thorell, 1870a [both family names protected by usage]	
Argyroneta Latreille, 1804a	?Neogene – Recent
846. <i>Argyroneta aquatica</i> (Clerck, 1757) [Recent]	Qt England
847. ? <i>Argyroneta longipes</i> Heer, 1865	Ne Öhningen
† <i>Vectoraneus</i> Selden, 2001	Palaeogene
848. <i>Vectoraneus yulei</i> Selden, 2001*	Pa Bembridge Marls
DESIDAE Pocock, 1895	Palaeogene – Recent
Myro O. P.-Cambridge, 1876	Palaeogene – Recent
849. <i>Myro extinctus</i> Petrunkevitch, 1958 ...[possibly belongs in Dictynidae].....	Pa Baltic amber
850. <i>Myro hirsutus</i> Petrunkevitch, 1942	Pa Baltic amber
AMPHINECTIDAE Forster & Wilton, 1973	Recent
= NEOLANIDAE Forster & Wilton, 1973	
no fossil record	
CYCLOCTENIDAE Simon, 1898a	Recent
no fossil record	
HAHNIIDAE Bertkau, 1878a	Palaeogene – Recent
† <i>Cymbiohahnia</i> Wunderlich, 2004v	Palaeogene
851. <i>Cymbiohahnia parens</i> Wunderlich, 2004v	Pa Baltic, Bitterfeld & Rovno amber
† <i>Eohahnia</i> Petrunkevitch, 1958	Palaeogene
852. <i>Eohahnia succini</i> Petrunkevitch, 1958*	Pa Baltic amber
† <i>Protohahnia</i> Wunderlich, 2004v	Palaeogene
853. <i>Protohahnia antiqua</i> Wunderlich, 2004v*	Pa Baltic amber

854. *Protohahnia tripartita* Wunderlich, 2004v Pa Baltic amber
genus uncertain
855. 'Tegenaria' *obscura* C. L. Koch & Berendt, 1854 Pa Baltic amber
- DICTYNIDAE O. P.-Cambridge, 1871** **Cretaceous – Recent**
- = RHOIDAE Thorell, 1873
- = † ARTHRODICTYNIDAE Petrunkevitch, 1942
- Dictynidae gen. et sp. indet *in* Penney (2002) K New Jersey amber
- Dictynidae sp. 1–2 *in* Wunderlich (2004v) Pa Baltic amber
- Dictynidae sp. 1–5 *in* Wunderlich (2008d) K Myanmar amber
- Dictyninae indet *in* Wunderlich (2012b) Pa Rovno amber
- Argenna Thorell, 1870a** **Neogene – Recent**
856. *Argenna fossilis* Petrunkevitch *in* Palmer, 1957 Ne Mojave Desert
- † **Arthrodictyna Petrunkevitch, 1942** **Palaeogene**
857. *Arthrodictyna segmentata* Petrunkevitch, 1942* Pa Baltic amber
- † **Balticocryphoeca Wunderlich, 2004v** **Palaeogene**
858. *Balticocryphoeca curvitarsis* Wunderlich, 2004v* Pa Baltic / Bitt. amber
- † **Brommellina Wunderlich, 2004v** **Palaeogene**
859. *Brommellina longungulae* Wunderlich, 2004v* Pa Baltic amber
- † **Burmadictyna Wunderlich, 2008d** **Cretaceous**
860. *Burmadictyna pecten* Wunderlich, 2008d* K Myanmar amber
- † **Chelicirrum Wunderlich, 2004v** **Palaeogene**
861. *Chelicirrum stridulans* Wunderlich, 2004v* Pa Baltic amber
- † **Cryphoezaga Wunderlich, 2004v** **Palaeogene**
862. *Cryphoezaga dubia* Wunderlich, 2004v* Pa Baltic amber
- Dictyna Sundevall, 1833** **Quaternary – Recent**
863. *Dictyna rufa* Wunderlich, 2012a Qt Madagascan copal
- † **Eobrommella Wunderlich, 2004v** **Palaeogene**
864. *Eobrommella scutata* Wunderlich, 2004v* Pa Baltic amber
- † **Eocryphoeca Petrunkevitch, 1946** **Palaeogene**
865. *Eocryphoeca bitterfeldensis* Wunderlich, 2004v Pa Bitterfeld amber
866. *Eocryphoeca electrina* Wunderlich, 2004v Pa Baltic amber
867. *Eocryphoeca falcata* Wunderlich, 2004v Pa Baltic amber
868. *Eocryphoeca gibbifera* Wunderlich, 2004v Pa Baltic amber
869. *Eocryphoeca gracilipes* (C. L. Koch & Berendt, 1854)* Pa Baltic amber
870. *Eocryphoeca ligula* Wunderlich, 2004v Pa Baltic amber
871. *Eocryphoeca mammilla* Wunderlich, 2004v Pa Baltic amber
872. *Eocryphoeca splendens* Wunderlich, 2004v Pa Baltic amber
- Eocryphoeca* sp. *in* Wunderlich (2004v) Pa Baltic amber
- † **Eocryphoecara Wunderlich, 2004v** **Palaeogene**
873. *Eocryphoecara abicera* Wunderlich, 2004v* Pa Baltic amber
- † **Eodictyna Wunderlich, 2004v** **Palaeogene**

874. *Eodictyna communis* Wunderlich, 2004v* Pa Baltic amber
- † ***Eolathys* Petrunkevitch, 1950** Palaeogene
875. *Eolathys debilis* Petrunkevitch, 1950 Pa Baltic amber
876. *Eolathys succini* Petrunkevitch, 1950* Pa Baltic amber
- † ***Flagelldictyna* Wunderlich, 2012a** Quaternary
877. *Flagelldictyna copalis* Wunderlich, 2012a* Qt Madagascar copal
- † ***Gibbermastigusa* Wunderlich, 2004v** Palaeogene
878. *Gibbermastigusa lateralis* Wunderlich, 2004v* Pa Baltic amber
- † ***Hispaniolyna* Wunderlich, 1988** Neogene
879. *Hispaniolyna hirsuta* Wunderlich, 1988 Ne Dominican amber
880. *Hispaniolyna magna* Wunderlich, 1988* Ne Dominican amber
- † ***Mastigusa* Menge in C. L. Koch & Berendt, 1854** Palaeogene
- = † *Eotetralius* Wunderlich, 1982 [nomen nudum]
881. *Mastigusa acuminata* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
882. *Mastigusa arcuata* Wunderlich, 2004v Pa Baltic amber
883. *Mastigusa bitterfeldensis* Wunderlich, 2004v Pa Bitterfeld amber
884. *Mastigusa laticymbium* Wunderlich, 2004v Pa Baltic amber
885. *Mastigusa magnibulbus* Wunderlich, 2004v Pa Bitterfeld amber
886. *Mastigusa media* Wunderlich, 1986 Pa Baltic amber
887. *Mastigusa modesta* Wunderlich, 1986 Pa Baltic amber
888. *Mastigusa scutata* Wunderlich, 2004v Pa Baltic amber
- Mastigusa* sp. in Wunderlich (2004v) Pa Baltic amber
- † ***Mizagalla* Wunderlich, 2004v** Palaeogene
889. *Mizagalla quattuor* Wunderlich, 2004v* Pa Baltic amber
890. *Mizagalla tuberculata* Wunderlich, 2004v Pa Baltic amber
- † ***Palaeodictyna* Wunderlich, 1988** Neogene
891. *Palaeodictyna intermedia* Wunderlich, 1988 Ne Dominican amber
892. *Palaeodictyna longispina* Wunderlich, 1988 Ne Dominican amber
893. *Palaeodictyna singularis* Wunderlich, 1988 Ne Dominican amber
894. *Palaeodictyna spiculum* Wunderlich, 1988 Ne Dominican amber
895. *Palaeodictyna termitophila* Wunderlich, 1988* Ne Dominican amber
896. *Palaeodictyna unispina* Wunderlich, 1988 Ne Dominican amber
- † ***Palaeolathys* Wunderlich, 1986** Neogene
897. *Palaeolathys circumductus* Wunderlich, 1988 Ne Dominican amber
898. *Palaeolathys copalis* Wunderlich, 1986 Qt Dominican copal
899. *Palaeolathys quadruplex* Wunderlich, 1988 Ne Dominican amber
900. *Palaeolathys similis* Wunderlich, 1988 Ne Dominican amber
901. *Palaeolathys spinosa* Wunderlich, 1986* Ne Dominican amber
- Palaeolathys* sp. in Wunderlich (1988) Ne Dominican amber
- † ***Protomastigusa* Wunderlich, 2004v** Palaeogene
902. *Protomastigusa composita* Wunderlich, 2004v Pa Baltic amber

† <i>Scopulyna</i> Wunderlich, 2004v	Palaeogene
903. <i>Scopulyna cursor</i> Wunderlich, 2004v	Pa Baltic amber
† <i>Succinya</i> Wunderlich, 1988	Neogene
904. <i>Succinya longembolus</i> Wunderlich, 1988	Ne Dominican amber
905. <i>Succinya pulcher</i> Wunderlich, 1988*	Ne Dominican amber
906. <i>Succinya spinipalpus</i> Wunderlich, 1988	Ne Dominican amber
<i>Thallumetus</i> Simon, 1892b	Subrecent – Recent
907. <i>Thallumetus copalis</i> Wunderlich, 2004at	Qt Colombian copal
AMAUROBIIDAE Thorell, 1870a	Palaeogene – Recent
= CINIFLONIDAE Blackwall, 1841	
[partly also Dictynidae; based on a generic synonym]	
Amaurobiinae sp. <i>in</i> Wunderlich (2004u)	Pa Baltic amber
PHYXELIDIDAE Lehtinen, 1967	Recent
no fossil record	
TITANOECIDAE Lehtinen, 1967	Quaternary – Recent
† <i>Copaldictyna</i> Wunderlich, 2004v	Quaternary
Tentative transfer by Wunderlich (2012a)	
908. <i>Copaldictyna madagascariensis</i> Wunderlich, 2004v*	Qt Madagascan copal
NICODAMIDAE Simon, 1898	Recent
= MEGADICTYNIDAE Lehtinen, 1967	
no fossil record	
TENGELLIDAE Dahl, 1908	Recent
no fossil record	
EUTICHURIDAE Lehtinen, 1967	Recent
= CHEIRACANTHIDAE Wagner, 1887	
Strotarchus Simon, 1888	Neogene – Recent
= † <i>Mimeutychurus</i> Petrunkevitch, 1963 [tentative synonymy]	
909. <i>Strotarchus heidti</i> Wunderlich, 1988	Ne Dominican amber
910. <i>Strotarchus paradoxus</i> (Petrunkevitch, 1963)	Ne Chiapas amber
MITURGIDAE Simon, 1885a	Palaeogene – Recent
= ZORIDAE F.O.P.-Cambridge, 1893	
† <i>Zorapostenus</i> Wunderlich, 2008c	Palaeogene
911. <i>Zorapostenus raveni</i> Wunderlich, 2008c	Pa Baltic amber
ANYPHAENIDAE Bertkau, 1878a	Palaeogene – Recent

= AMAUROBIOIDIDAE Hickman, 1949

<i>Anyphaena</i> Sundevall, 1833	Palaeogene – Recent
912. ‘ <i>Anyphaena</i> ’ <i>fuscata</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
<i>Anyphaenoides</i> Berland, 1913	Neogene – Recent
913. <i>Anyphaenoides bulla</i> (Wunderlich, 1988)	Ne Dominican amber
<i>Lupettiana</i> Brescovit, 1997	Neogene – Recent
914. <i>Lupettiana ligula</i> (Wunderlich, 1988)	Ne Dominican amber
<i>Wulfila</i> O. P.-Cambridge, 1895	Neogene – Recent
915. <i>Wulfila spinipes</i> Wunderlich, 1988	Ne Dominican amber
 LIOCRANIDAE Simon, 1897a	 Palaeogene – Recent
?Liocranidae <i>in</i> Wunderlich (1988)	Ne Dominican amber
Apostenus Westring, 1851	Palaeogene – Recent
916. <i>Apostenus arnoldorum</i> Wunderlich, 2004ag	Pa Baltic amber
917. <i>Apostenus bigibber</i> Wunderlich, 2004ag	Pa Baltic / Bitt. amber
918. <i>Apostenus spinimanus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
Donaea Strand, 1932	Quaternary – Recent
919. <i>Donaea collustrata</i> Bosselaers & Dierick, 2010 [Recent]	Qt – R Madagascar
† Palaeospinisoma Wunderlich, 2004ag	Palaeogene
920. <i>Palaeospinisoma femoralis</i> Wunderlich, 2004ag*	Pa Baltic amber

CLUBIONOIDEA *incertae sedis*

Wunderlich (2011d) proposed removing almost all the amber fossils from the clubionids *sensu stricto*. We follow this in part for the two genera below, but would prefer a more formal treatment before accepting all these transfers. In general the delimitation of even modern clubionids, and related forms, is problematic.

† Concursator Petrunkevitch, 1958	Palaeogene
921. <i>Concursator nudipes</i> Petrunkevitch, 1958*	Pa Baltic amber
† Systariella Wunderlich, 2004af	Palaeogene
922. <i>Systariella magnioculi</i> Wunderlich, 2004af*	Pa Baltic amber

 CLUBIONIDAE Simon, 1895	 Palaeogene – Recent
Clubionidae gen. et sp. <i>in</i> Nishikawa (1974)	Qt Mizunami copal
Clubiona Latreille, 1804a	Palaeogene – Recent
923. <i>Clubiona arcana</i> Scudder, 1890a	Pa Florissant
924. <i>Clubiona attenuata</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
925. <i>Clubiona curvispinosa</i> Petrunkevitch, 1922	Pa Florissant
926. <i>Clubiona florissanti</i> Petrunkevitch, 1922	Pa Florissant
927. <i>Clubiona lanata</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
928. <i>Clubiona microphthalma</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
929. <i>Clubiona pubescens</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
930. <i>Clubiona sericea</i> C. L. Koch & Berendt, 1854	Pa Baltic amber

931.	<i>Clubiona tomentosa</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
†	<i>Desultor</i> Petrunkevitch, 1942		Palaeogene
932.	<i>Desultor depressus</i> Petrunkevitch, 1942	Pa	Baltic amber
	<i>Elaver</i> O. P.-Cambridge, 1898		Neogene – Recent
933.	<i>Elaver nutua</i> (Wunderlich, 1988)	Ne	Dominican amber
†	<i>Eobumbatrix</i> Petrunkevitch, 1922		Palaeogene
934.	<i>Eobumbatrix latebrosa</i> (Scudder, 1890a)*	Pa	Florissant
†	<i>Eodoter</i> Petrunkevitch, 1958		Palaeogene
935.	<i>Eodoter eopala</i> Wunderlich, 2004af	Pa	Baltic amber
936.	<i>Eodoter ionimammillae</i> Wunderlich, 2012c	Pa	Baltic amber
937.	<i>Eodoter magnificus</i> Petrunkevitch, 1958*	Pa	Baltic amber
938.	<i>Eodoter scutatus</i> Wunderlich, 2011d	Pa	Baltic amber
939.	? <i>Eodoter tibialis</i> Wunderlich, 2011d	Pa	Baltic amber
†	<i>Eostentatrix</i> Petrunkevitch, 1922		Palaeogene
940.	<i>Eostentatrix cockerelli</i> Petrunkevitch, 1922	Pa	Florissant
941.	<i>Eostentatrix ostentata</i> (Scudder, 1890a)*	Pa	Florissant
†	<i>Eoversatrix</i> Petrunkevitch, 1922		Palaeogene
942.	<i>Eoversatrix eversa</i> (Scudder, 1890a)*	Pa	Florissant
†	<i>Machilla</i> Petrunkevitch, 1958 [family uncertain]		Palaeogene
943.	<i>Machilla setosa</i> Petrunkevitch, 1958*	Pa	Baltic amber
†	<i>Massula</i> Petrunkevitch, 1942 [family uncertain]		Palaeogene
944.	<i>Massula klebsi</i> Petrunkevitch, 1942*	Pa	Baltic amber
†	<i>Prosocer</i> Petrunkevitch, 1963		Neogene
945.	<i>Prosocer mollis</i> Petrunkevitch, 1963*	Ne	Chiapas amber

Clubionidae incertae sedis

†	<i>Chiapasona</i> Petrunkevitch, 1963		Neogene
946.	<i>Chiapasona defuncta</i> Petrunkevitch, 1963*	Ne	Chiapas amber

CORINNIDAE Karsch, 1880a

= MYRMECIIDAE C. L. Koch, 1851 [name already used for ants]

NB: Extinct genera were not considered in the otherwise comprehensive revision of Ramírez (2014), some fossil corinnids may now belong in other families.

†	<i>Ablator</i> Petrunkevitch, 1942		Palaeogene
	= † <i>Abiguritor</i> Petrunkevitch, 1942		
947.	<i>Ablator biguttatus</i> Wunderlich, 2004ah	Pa	Baltic amber
948.	<i>Ablator curvatus</i> Wunderlich, 2004ah	Pa	Baltic amber
949.	<i>Ablator deminuens</i> Wunderlich, 2004ah	Pa	Baltic amber
950.	<i>Ablator depressus</i> Wunderlich, 2004ah	Pa	Baltic amber
951.	<i>Ablator duomammillae</i> Wunderlich, 2004ah	Pa	Baltic amber
952.	<i>Ablator felix</i> (Petrunkevitch, 1958)	Pa	Baltic amber
953.	<i>Ablator inevolvens</i> Wunderlich, 2004ah	Pa	Baltic amber

954. *Ablator longus* Wunderlich, 2004ah Pa Baltic amber
955. *Ablator nonguttatus* Wunderlich, 2004ah Pa Baltic amber
956. *Ablator parvus* Wunderlich, 2004ah Pa Baltic amber
957. *Ablator plumosus* (Petrunkewitch, 1950) Pa Baltic amber
958. *Ablator robustus* Wunderlich, 2004ah Pa Baltic amber
959. *Ablator scutatus* Wunderlich, 2004ah Pa Baltic amber
960. *Ablator splendens* Wunderlich, 2004ah Pa Baltic amber
961. *Ablator triguttatus* (C. L. Koch & Berendt, 1854)* Pa Baltic amber
- i. = *Philodromus microcephalus* C. L. Koch & Berendt,
1854 Pa Baltic amber
- ii. = *Philodromus squamiger* C. L. Koch & Berendt, 1854 ..Pa Baltic amber
- iii. = *Abiligurator niger* Petrunkewitch, 1942 Pa Baltic amber
- † ***Alterphrurolithus* Wunderlich, 2004ah** **Palaeogene**
962. *Alterphrurolithus longipes* Wunderlich, 2004ah Pa Baltic amber
- Castianeira* Keyserling, 1880b** **Neogene – Recent**
963. *Castianeira tenebricosa* Wunderlich, 1988 Ne Dominican amber
- † ***Chemmisomma* Wunderlich, 1988** **Neogene**
964. *Chemmisomma dubia* Wunderlich, 1988* Ne Dominican amber
- Corinna* C. L. Koch, 1842a** **Neogene – Recent**
965. *Corinna flagelliformis* Wunderlich, 1988 Ne Dominican amber
- † ***Cornucymbium* Wunderlich, 2004ah** **Palaeogene**
966. *Cornucymbium insolens* Wunderlich, 2004ah* Pa Baltic amber
- † ***Cryptoplanus* Petrunkewitch, 1958** **Palaeogene**
967. *Cryptoplanus bulbosus* Wunderlich, 2004ah Pa Baltic amber
968. *Cryptoplanus complicatus* Wunderlich, 2004ah Pa Baltic amber
969. *Cryptoplanus incidens* Wunderlich, 2004ah Pa Baltic amber
970. *Cryptoplanus lanatus* (Petrunkewitch, 1958) Pa Baltic amber
971. *Cryptoplanus paradoxus* Petrunkewitch, 1958* Pa Baltic amber
972. *Cryptoplanus sericatus* (C. L. Koch & Berendt, 1854) Pa Baltic amber
973. *Cryptoplanus sinuosus* Wunderlich, 2004ah Pa Baltic amber
- Cryptoplanus* sp. in Wunderlich (2004ah) Pa Baltic amber
- † ***Eomazax* Petrunkewitch, 1958** **Palaeogene**
974. *Eomazax pulcher* Petrunkewitch, 1958* Pa Baltic amber
- Megalostrata* Karsch, 1880a** **Neogene – Recent**
975. *Megalostrata grandis* Wunderlich, 1988 Ne Dominican amber
- † ***Myrmecorinna* Wunderlich, 2004ah** **Palaeogene**
976. *Myrmecorinna gracilis* Wunderlich, 2004ah* Pa Baltic amber
- † ***Palpiraptor* Wunderlich, 2011f** **Quaternary**
977. *Palpiraptor myrmarachnoides* Wunderlich, 2011f* Qt Madagascar copal
- † ***Protoorthobula* Wunderlich, 2004ah** **Palaeogene**
978. *Protoorthobula bifida* Wunderlich, 2004ah* Pa Baltic amber

979. *Protoorthobula deelemani* Wunderlich, 2004ah Pa Baltic / Bitt. Amber
- TRACHELIDAE Simon, 1897** Neogene – Recent
- Trachelas* L. Koch, 1872 Neogene
980. *Trachelas poinari* Penney, 2001 Ne Dominican amber
- PHRUROLITHIDAE Banks, 1892** Palaeogene – Recent
- Phrurolithus* C. L. Koch, 1839b Palaeogene – Recent
981. *Phrurolithus extinctus* Petrunkevitch, 1958 Pa Baltic amber
982. *Phrurolithus fossilis* Petrunkevitch, 1958 Pa Baltic amber
983. *Phrurolithus ipseni* Petrunkevitch, 1958 Pa Baltic amber
- ZODARIIDAE Thorell, 1881** Palaeogene – Recent
- = CRYPTOTHELIDAE L. Koch, 1872 [younger name protected by usage]
- = † ADJUTORIDAE Petrunkevitch, 1942
- Zodariidae gen. et sp. indet 1–4 in Wunderlich (2004ae) Pa Baltic amber
- † *Adjutor* Petrunkevitch, 1942 Palaeogene
984. *Adjutor deformis* Petrunkevitch, 1958 Pa Baltic amber
985. *Adjutor mirabilis* Petrunkevitch, 1942* Pa Baltic amber
- † *Admissor* Petrunkevitch, 1942 Palaeogene
986. *Admissor aculeatus* Petrunkevitch, 1942* Pa Baltic amber
- † *Adorator* Petrunkevitch, 1942 Palaeogene
987. *Adorator hispidus* (C. L. Koch & Berendt, 1854) Pa Baltic / Rovno amber
- i. = *Segestria cylindrica* C. L. Koch & Berendt, 1854 Pa Baltic amber
- ii. = *Eresus curtipes* C. L. Koch & Berendt, 1854 Pa Baltic amber
- iii. = *Eresus monachus* C. L. Koch & Berendt, 1854 Pa Baltic amber
- iv. = *Adorator brevipes* Petrunkevitch, 1942* Pa Baltic amber
988. *Adorator samlandicus* Petrunkevitch, 1942 Pa Baltic amber
- † *Angusdarion* Wunderlich, 2004ae Palaeogene
989. *Angusdarion humilis* Wunderlich, 2004ae* Pa Baltic amber
- † *Anniculus* Petrunkevitch, 1942 Palaeogene
990. *Anniculus balticus* Petrunkevitch, 1942* Pa Baltic amber
- † *Eocydrole* Petrunkevitch, 1958 Palaeogene
991. *Eocydrole mortua* Petrunkevitch, 1958* Pa Baltic amber
- † *Propago* Petrunkevitch, 1963 Neogene
992. *Propago debilis* Petrunkevitch, 1963* Ne Chiapas amber
- † *Spinizodarion* Wunderlich, 2004ae Palaeogene
993. *Spinizodarion ananulum* Wunderlich, 2004ae* Pa Baltic amber
- † *Zodariodamus* Wunderlich 2004ae Palaeogene
994. *Zodariodamus recurvatus* Wunderlich 2004ae* Pa Baltic amber
- PENESTOMIDAE Simon, 1903** Recent

no fossil record

† EPHALMATORIDAE Petrunkevitch, 1950	Palaeogene
† <i>Ephalmator</i> Petrunkevitch, 1950	Palaeogene
995. <i>Ephalmator bitterfeldensis</i> Wunderlich, 2004ad	Pa Bitterfeld amber
996. <i>Ephalmator calidus</i> Wunderlich, 2004ad	Pa Baltic amber
997. <i>Ephalmator debilis</i> Wunderlich, 2004ad	Pa Baltic amber
998. <i>Ephalmator distinctus</i> Wunderlich, 2004ad	Pa Baltic amber
999. <i>Ephalmator ellwangeri</i> Wunderlich, 2004ad	Pa Baltic amber
1000. ? <i>Ephalmator eximus</i> Petrunkevitch, 1958	Pa Baltic amber
1001. <i>Ephalmator fossilis</i> Petrunkevitch, 1950*	Pa Baltic amber
1002. <i>Ephalmator kerneggeri</i> Wunderlich, 2004ad	Pa Baltic amber
1003. <i>Ephalmator petrunkevitchi</i> Wunderlich, 2004ad	Pa Baltic amber
1004. <i>Ephalmator ruthildae</i> Wunderlich, 2004ad	Pa Baltic amber
1005. <i>Ephalmator tredecim</i> Wunderlich, 2012c	Pa Baltic amber
1006. <i>Ephalmator trudis</i> Wunderlich, 2004ad	Pa Baltic amber
1007. <i>Ephalmator turpiculus</i> Wunderlich, 2004ad	Pa Baltic amber
<i>Ephalmator</i> sp. in Wunderlich (2004ad)	Pa Baltic amber

CHUMMIDAE Jocqué, 2001

Recent

no fossil record

HOMALONYCHIDAE Simon, 1893

Recent

no fossil record

GNAPHOSOIDEA Simon, 1893

Palaeogene – Recent

AMMOXENIDAE Simon, 1893

Recent

no fossil record

CITHAERONIDAE Simon, 1893

Recent

no fossil record

GALLIENIELLIDAE Millot, 1947

Recent

no fossil record

TROCHANTERIIDAE Karsch, 1879

Palaeogene – Recent

= PLATORIDAE Simon, 1890

† *Eotrochanteria* Wunderlich, 2004am

Palaeogene

 1008. *Eotrochanteria kruegeri* Wunderlich, 2004am*

Pa Baltic amber

† *Sosybius* C. L. Koch & Berendt, 1854

Palaeogene

= † *Adamator* Petrunkevitch, 1942

= † *Adjuncctor* Petrunkevitch, 1942

= † *Adulatrix* Petrunkevitch, 1942

1009. *Sosybius berendti* Wunderlich, 2004am Pa Baltic amber
1010. *Sosybius decumana* (C. L. Koch & Berendt, 1854) Pa Baltic amber
1011. *Sosybius falcatus* Wunderlich, 2004am Pa Baltic amber
1012. *Sosybius fusca* (Petrunkevitch, 1942) Pa Baltic amber
1013. *Sosybius kochi* Wunderlich, 2004am Pa Baltic amber
1014. *Sosybius lateralis* Wunderlich, 2004am Pa Baltic amber
1015. *Sosybius longipes* Wunderlich, 2004am Pa Baltic amber
1016. *Sosybius major* C. L. Koch & Berendt, 1854 Pa Baltic amber
1017. *Sosybius minor* C. L. Koch & Berendt, 1854* Pa Baltic amber
1018. *Sosybius mizgirisi* Wunderlich, 2004am Pa Baltic amber
1019. *Sosybius parva* (Petrunkevitch, 1942) Pa Baltic amber
1020. *Sosybius perniciosus* Wunderlich, 2004am Pa Baltic amber
1021. *Sosybius rufa* (Petrunkevitch, 1942) Pa Baltic amber
1022. *Sosybius similis* Petrunkevitch, 1942 Pa Baltic amber
1023. *Sosybius succineus* (Petrunkevitch, 1942) Pa Baltic amber
1024. *Sosybius tibialis* Wunderlich, 2004am Pa Baltic amber
1025. *Sosybius unispinosus* Wunderlich, 2004am Pa Baltic amber
- Sosybius* sp. in Wunderlich (2004am, ar) Pa Baltic / Rovno amber
- † *Thereola* Petrunkevitch, 1955 **Palaeogene**
- = † *Thereola* Koch & Berendt, 1854 [preoccupied]
1026. *Thereola petiolata* (C. L. Koch & Berendt, 1854)* [♀ = ?*Dasuminia* sp.
according to Wunderlich 2004b] Pa Baltic amber
1027. *Thereola pubescens* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
- † *Trochanteridromulus* Wunderlich, 2004am **Palaeogene**
1028. *Trochanteridromulus glabripes* Wunderlich, 2004am* Pa Baltic amber
- † *Trochanteridromus* Wunderlich, 2004am **Palaeogene**
1029. *Trochanteridromus scutatus* Wunderlich, 2004am* Pa Baltic amber
- † *Veterator* Petrunkevitch, 1963 **Neogene**
1030. *Veterator angustus* Wunderlich, 1988 Ne Dominican amber
1031. *Veterator ascutum* Wunderlich, 1988 Ne Dominican amber
1032. *Veterator extinctus* Petrunkevitch, 1963* Ne Chiapas amber
1033. *Veterator incompletus* Wunderlich, 1982 Ne Dominican amber
1034. *Veterator longipes* Wunderlich, 1988 Ne Dominican amber
1035. *Veterator loricatus* Wunderlich, 1988 Ne Dominican amber
1036. *Veterator porrectus* Wunderlich, 1988 Ne Dominican amber
1037. *Veterator viduus* Wunderlich, 1988 Ne Dominican amber
- Veterator* sp. 1–2 in Wunderlich (1988) Ne Dominican amber
- LAMPONIDAE Simon, 1893** **Recent**
- no fossil record

PRODIDOMIDAE Simon, 1884a	Quaternary – Recent
= MILTIIDAE Thorell, 1873 [based on a generic synonym]	
Prodidomus Hentz, 1847	Quaternary – Recent
1038. <i>Prodidomus madagascariensis</i> Wunderlich, 2011c	Qt Madagascar copal
GNAPHOSIDAE Pocock, 1898	?Cretaceous – Recent
= DRASSIDAE Sundevall, 1833 [based on a generic synonym]	
† Captrix Petrunkevitch, 1942	Palaeogene
1039. <i>Captrix lineata</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
Drassodes Westring, 1851	Palaeogene – Recent
1040. <i>Drassodes cupreus</i> (Blackwall, 1834a) [Recent]	Qt England
1041. ? <i>Drassodes femurus</i> Lin, Zhang & Wang, 1989	Ne Shanwang
1042. ? <i>Drassodes sextii</i> Berland, 1939	Pa Aix-en-Provence
† Drassyllinus Wunderlich, 1988	Neogene
1043. <i>Drassyllinus aliter</i> Wunderlich, 1988*	Ne Dominican amber
† Eognaphosops Wunderlich, 2011b	Palaeogene
1044. <i>Eognaphosops cryptoplanoides</i> Wunderlich 2011b*	Pa Baltic amber
† Eomactator Petrunkevitch, 1958	Palaeogene
1045. <i>Eomactator hamatus</i> Wunderlich, 2011b	Pa Baltic amber
1046. <i>Eomactator hirsutipes</i> Wunderlich, 2011b	Pa Baltic amber
1047. <i>Eomactator mactatus</i> Petrunkevitch, 1958*	Pa Baltic amber
1048. <i>Eomactator obscurior</i> Wunderlich, 2011b	Pa Baltic amber
Gnaphosa Latreille, 1804a	?Cretaceous – Recent
1049. <i>Gnaphosa affinis</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
i. = <i>Philodromus dubius</i> C. L. Koch & Berendt, 1854	
1050. <i>Gnaphosa ambigua</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1051. <i>Gnaphosa liaoningensis</i> Chang, 2004	
[generic assignment unreliable!]	K Jehol biota
Micaria Westring, 1851	Palaeogene – Recent
1052. <i>Micaria procera</i> C. L. Koch & Berendt, 1954	Pa Baltic amber
1053. <i>Micaria tenella</i> Heer, 1865	Ne Öhningen
† Palaeodrassus Petrunkevitch, 1922	Palaeogene
1054. <i>Palaeodrassus cockerelli</i> Petrunkevitch, 1922	Pa Florissant
1055. <i>Palaeodrassus florissanti</i> Petrunkevitch, 1922	Pa Florissant
1056. <i>Palaeodrassus hesternus</i> (Scudder, 1890a)	Pa Florissant
1057. <i>Palaeodrassus ingenuus</i> (Scudder, 1890a)*	Pa Florissant
1058. <i>Palaeodrassus interitus</i> (Scudder, 1890a)	Pa Florissant
Scopoides Platnick, 1989	Palaeogene – Recent
1059. <i>Scopoides dominicanus</i> Wunderlich, 2011g	Ne Dominican amber
Zelotes Gistel, 1848	Palaeogene
1060. <i>Zelotes concinna</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1061. <i>Zelotes mundula</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber

- i. = *Melanophora nobilis* C. L. Koch & Berendt, 1854 Pa Baltic amber
1062. *Zelotes regalis* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- † ***Zelotetis* Wunderlich, 2011b** **Palaeogene**
1063. *Zelotetis calefacta* Wunderlich, 2011b Pa Baltic amber
- SELENOPIDAE Simon, 1897a** **Palaeogene – Recent**
- † ***Garcorops* Corronca, 2003** **Quaternary – Recent**
1064. *Garcorops jadis* Bosselaers, 2004 Qt Madagascar copal
- i. = ?*Anyplops cortex* Wunderlich, 2004as Qt Madagascar copal
- Selenops* Latreille, 1819** **Palaeogene – Recent**
1065. *Selenops benoiti* Wunderlich, 2004as Qt Madagascar copal
1066. *Selenops beynai* Schawaller, 1984 Ne Dominican amber
1067. *Selenops dominicanus* Wunderlich, 2004an Ne Dominican amber
- Selenops* sp. in Wunderlich (1988) Ne Dominican amber
- Selenops* sp. in García-Villafuerte (2006b) Ne Chiapas amber
- Selenops* sp. in Penney (2007) Pa Le Quesnoy amber
- SPARASSIDAE Bertkau, 1872** **Palaeogene – Recent**
- = HETEROPODIDAE Thorell, 1873
- = MICROMMATIDAE Bertkau, 1878a
- = EUSPARASSIDAE Järvi, 1912
- Sparassidae* sp. 1–2 in (Wunderlich 2008c) Pa Baltic amber
- † ***Caduceator* Petrunkevitch, 1942** **Palaeogene**
1068. *Caduceator minutus* Petrunkevitch, 1942* Pa Baltic amber
1069. *Caduceator quadrimaculatus* Petrunkevitch, 1950 Pa Baltic amber
- † ***Collacteus* Petrunkevitch, 1942** **Palaeogene**
1070. *Collacteus captivus* Petrunkevitch, 1942* Pa Baltic amber
- † ***Eostaianus* Petrunkevitch, 1950** **Palaeogene**
1071. *Eostaianus succini* Petrunkevitch, 1950* Pa Baltic amber
- † ***Eostasina* Petrunkevitch, 1942** **Palaeogene**
1072. *Eostasina aculeata* Petrunkevitch, 1942* Pa Baltic amber
- Eusparassus* Simon 1903** **Palaeogene – Recent**
1073. *Eusparassus crassipes* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- Heteropoda* Latreille, 1804a** **Palaeogene – Recent**
- = † *Retina* Hong, 1985
1074. *Heteropoda rpbusta* [sic] (Hong, 1985) Ne Shanwang
- [NB: as '*H. robusta*' this would be a junior homonym of a living species.]
- Pseudosparianthis* Simon, 1887** **Neogene – Recent**
1075. *Pseudosparianthis pfeifferi* (Wunderlich, 1988) Ne Dominican amber
- Zachria* L. Koch, 1875** **Palaeogene – Recent**
- [NB: An Australian genus; Wunderlich (2012c) regarded at least *Z. desiderabilis* as gen. indet.]
1076. *Zachria desiderabilis* Petrunkevitch, 1950 Pa Baltic amber

1077. *Zachria peculiata* Petrunkevitch, 1946 Pa Baltic amber
1078. *Zachria restincta* Petrunkevitch, 1958 Pa Baltic amber
- PHILODROMIDAE Thorell, 1870a** **Cretaceous – Recent**
- Philodromidae sp. *in* Wunderlich (1988) Ne Dominican amber
- Philodromidae sp. *in* Wunderlich (2004ae) Ne Baltic amber
- † Cretadromus Cheng, Shen & Gao, 2009** **Cretaceous**
1079. *Cretadromus liaoningensis* Cheng, Shen & Gao, 2009 K Liaoning Province
[NB: Wunderlich (2012d) suggested this could be a Theridosomatidae]
- † Eothanatus Petrunkevitch, 1950** **Palaeogene – Recent**
1080. *Eothanatus diritatis* Petrunkevitch, 1950* Pa Baltic amber
- THOMISIDAE Sundevall, 1833** **Palaeogene – Recent**
- = APANTHOCHILIDAE Thorell, 1873
- = MISUMENIDAE Thorell, 1887
- = STIPHROPODIDAE Simon, 1895
- = XYSTICIDAE Dahl, 1912
- = BORBOROPACTIDAE Wunderlich, 2004ao
- Thomisidae gen. et sp. *in* Nishikawa (1974) Qt Mizunami copal
- Thomisidae gen. et sp. *in* Bottali (1975) Qt Italy
- Thomisidae gen. et sp. *in* Schawaller (1982d) Ne Willershausen
- Thomisidae gen. et sp. *in* Wunderlich (1988) Ne Dominican amber
- Thomisidae gen. et sp. 1–2 *in* Wunderlich (2004ap) Pa Baltic amber
- Thomisidae gen. et sp. *in* Garcíá-Villafuerte (2006b) Ne Chiapas amber
- Coriarachne Thorell, 1870b** **Quaternary – Recent**
- Coriarachne sp. *in* Cutler (1970) Qt Wyoming
- † Ecotona Lin, Zhang & Wang, 1989 [ex Araneidae]** **Neogene**
1081. *Ecotona brunnea* Zhang, Sun & Zhang, 1994 Ne Shanwang
1082. *Ecotona pilulifera* Zhang, Sun & Zhang, 1994 Ne Shanwang
1083. *Ecotona transipeda* Lin, Zhang & Wang, 1989* Ne Shanwang
- † Facundia Petrunkevitch, 1942** **Palaeogene**
1084. *Facundia clara* Petrunkevitch, 1942* Pa Baltic amber
- † Fiducia Petrunkevitch, 1950** **Palaeogene**
1085. *Fiducia tenuipes* Petrunkevitch, 1950* Pa Baltic amber
- † Filiolella Petrunkevitch, 1955a** **Palaeogene**
- = † *Filiola* Petrunkevitch, 1942 [preoccupied]
1086. *Filiolella argentata* (Petrunkevitch, 1942)* Pa Baltic amber
- † Heterotmarus Wunderlich, 1988** **Neogene**
1087. *Heterotmarus altus* Wunderlich, 1988* Ne Dominican amber
- † Komisumena Ono, 1981** **Neogene**
1088. *Komisumena rosae* Ono, 1981* Ne Dominican amber

† <i>Miothomisus</i> Zhang, Sun & Zhang, 1994	Neogene
1089. <i>Miothomisus subnudus</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
1090. <i>Miothomisus sylvaticus</i> Zhang, Sun & Zhang, 1994*	Ne Shanwang
<i>Misumena</i> Latreille, 1804a	Palaeogene – Recent
1091. <i>Misumena samlandica</i> Petrunkevitch, 1942	Pa Baltic amber
† <i>Palaeoxysticus</i> Wunderlich, 1985	Neogene
1092. <i>Palaeoxysticus extinctus</i> Wunderlich, 1985	Ne Randecker Maar
† <i>Parvulus</i> Zhang, Sun & Zhang, 1994	Neogene
1093. <i>Parvulus latissimus</i> Zhang, Sun & Zhang, 1994*	Ne Shanwang
† <i>Succinaenigma</i> Wunderlich, 2004ap	Palaeogene
1094. <i>Succinaenigma raptor</i> Wunderlich, 2004ap*	Pa Baltic amber
† <i>Succiniraptor</i> Wunderlich, 2004ao	Palaeogene
1095. <i>Succiniraptor radiatus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
i. = <i>Succiniraptor paradoxus</i> Wunderlich, 2004ao*	Pa Baltic amber
<i>Synema</i> Simon, 1864	Palaeogene – Recent
1096. <i>Synema enigmaticum</i> Berland, 1939	Pa Aix-en-Provence
† <i>Syphax</i> C. L. Koch & Berendt, 1854	Palaeogene
1097. <i>Syphax asper</i> Petrunkevitch, 1950	Pa Baltic amber
1098. <i>Syphax crassipes</i> Petrunkevitch, 1942	Pa Baltic amber
1099. <i>Syphax fuliginosus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1100. <i>Syphax gracilis</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1101. <i>Syphax megacephalus</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
1102. <i>Syphax thoracicus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† <i>Thomisidites</i> Straus, 1967	Neogene
1103. <i>Thomisidites hercynicus</i> , Straus, 1967*	Ne Willershausen
† <i>Thomisiraptor</i> Wunderlich, 2004ap	Palaeogene
1104. <i>Thomisiraptor liedtkei</i> Wunderlich, 2004ap*	Pa Baltic amber
<i>Thomisus</i> Walckenaer, 1805	Palaeogene – Recent
1105. <i>Thomisus defossus</i> Scudder, 1890a	Pa Florissant
1106. <i>Thomisus disjunctus</i> Scudder, 1890a	Pa Florissant
1107. <i>Thomisus lividus</i> Heer, 1865	Ne Öhningen
1108. <i>Thomisus resutus</i> Scudder, 1890a	Pa Florissant
1109. <i>Thomisus sulzeri</i> Heer, 1865	Ne Öhningen
<i>Xysticus</i> C. L. Koch, 1835	Palaeogene – Recent
1110. ? <i>Xysticus annulipes</i> Bertkau, 1878b	Ne Rott, Germany
1111. <i>Xysticus archaeopalpus</i> Leech & Matthews, 1971	Ne Alaska
1112. <i>Xysticus oeningensis</i> (Heer, 1865)	Ne Öhningen
<i>Xysticus</i> sp. in Protescu (1937)	Pa Romanian amber
SALTICIDAE Blackwall, 1841	Palaeogene – Recent
= ATTIDAE Sundevall, 1833 [based on a generic synonym]	

= LYSSOMANIDAE Peckham & Wheeler, 1889		
Salticidae gen. et sp. <i>in</i> Schawaller (1982d)	Ne Willershausen
Salticidae incertae sedis <i>in</i> Selden (2014b)	Pa Isle of Wight
† Almolinus Petrunkevitch, 1958	Palaeogene
1113. <i>Almolinus bitterfeldensis</i> Wunderlich, 2004aq	Pa Bitterfeld amber
1114. <i>Almolinus clarus</i> Petrunkevitch, 1958*	Pa Baltic amber
1115. <i>Almolinus ligula</i> Wunderlich, 2004aq	Pa Baltic amber
? <i>Almolinus</i> sp. <i>in</i> Wunderlich (2004aq)	Pa Baltic amber
† Attoides Brongniart, 1877	Palaeogene
1116. <i>Attoides eresiformis</i> Brongniart, 1877	Pa Aix-en-Provence
† Calilinus Wunderlich, 2004aq	Palaeogene
1117. <i>Calilinus fleissneri</i> Wunderlich, 2004aq*	Pa Baltic amber
† Cenattus Petrunkevitch, 1942	Palaeogene
1118. <i>Cenattus exophthalmicus</i> Petrunkevitch, 1942*	Pa Baltic amber
Corythalia C. L. Koch, 1851	Neogene – Recent
1119. <i>Corythalia ocululiter</i> Wunderlich, 1988	Ne Dominican amber
1120. <i>Corythalia pilosa</i> Wunderlich, 1982	Ne Dominican amber
1121. <i>Corythalia scissa</i> Wunderlich, 1988	Ne Dominican amber
† Descangeles Wunderlich, 1988	Neogene
1122. <i>Descangeles pygmaeus</i> Wunderlich, 1988*	Ne Dominican amber
<i>Descangeles</i> sp. 1–2 <i>in</i> Wunderlich (1988)	Ne Dominican amber
Descanso Peckham & Peckham, 1892	Neogene – Recent
<i>Descanso</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
† Distanilinus Wunderlich, 2004aq	Palaeogene
1123. <i>Distanilinus filum</i> Wunderlich, 2004aq	Pa Baltic amber
1124. <i>Distanilinus nutus</i> Wunderlich, 2004aq*	Pa Baltic amber
1125. <i>Distanilinus paranutus</i> Wunderlich, 2004aq	Pa Baltic amber
1126. <i>Distanilinus pernatus</i> Wunderlich, 2004aq	Pa Baltic amber
† Eoattopsis Gourret, 1887	Palaeogene
1127. <i>Eoattopsis hirsutus</i> Gourret, 1887*	Pa Aix-en-Provence
† Eolinus Petrunkevitch, 1942	Palaeogene
1128. <i>Eolinus balticus</i> Žabka, 1988	Pa Baltic amber
1129. <i>Eolinus fungus</i> Wunderlich, 2004aq	Pa Baltic amber
1130. <i>Eolinus insuriens</i> Wunderlich, 2004aq	Pa Baltic amber
1131. <i>Eolinus prominens</i> Wunderlich, 2004aq	Pa Baltic amber
1132. <i>Eolinus samlandica</i> Wunderlich, 2004aq	Pa Baltic amber
1133. <i>Eolinus succineus</i> Petrunkevitch, 1942*	Pa Baltic amber
1134. <i>Eolinus theryi</i> Petrunkevitch, 1942	Pa Baltic amber
1135. <i>Eolinus theryoides</i> Wunderlich, 2004aq	Pa Baltic amber
1136. <i>Eolinus tyschenkoi</i> Proszynski & Žabka, 1980	Pa Baltic amber
1137. <i>Eolinus vates</i> Wunderlich, 2004aq	Pa Baltic amber

<i>Eolinus</i> sp. in Wunderlich (2004aq)	Pa	Baltic amber
<i>Euophrys</i> C. L. Koch, 1834		Palaeogene – Recent
1138. <i>Euophrys gibberula</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
1139. <i>Euophrys randeckensis</i> Schawaller & Ono, 1979	Ne	Randecker Maar
† <i>Evagoratus</i> Zhang, Sun & Zhang, 1994		Neogene
1140. <i>Evagoratus longicruris</i> Zhang, Sun & Zhang, 1994	Ne	Shanwang
† <i>Gorgopsidis</i> Wunderlich, 2004aq		Palaeogene
1141. <i>Gorgopsidis bechlyi</i> Wunderlich, 2004aq*	Pa	Baltic amber
† <i>Gorgopsina</i> Petrunkevitch, 1955a		Palaeogene
1142. <i>Gorgopsina amabilis</i> Wunderlich, 2004aq	Pa	Baltic amber
1143. <i>Gorgopsina constricta</i> Wunderlich, 2004aq	Pa	Baltic amber
1144. <i>Gorgopsina expandens</i> Wunderlich, 2004aq	Pa	Baltic amber
1145. ‘ <i>Gorgopsina</i> ’ <i>fasciata</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
1146. <i>Gorgopsina flexuosa</i> Wunderlich, 2004aq	Pa	Baltic amber
1147. <i>Gorgopsina formosa</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
1148. <i>Gorgopsina fractura</i> Wunderlich, 2004ar	Pa	Rovno amber
1149. <i>Gorgopsina frenata</i> (C. L. Koch & Berendt, 1854)*	Pa	Baltic amber
1150. <i>Gorgopsina inclusa</i> Wunderlich, 2004aq	Pa	Baltic amber
1151. <i>Gorgopsina jucunda</i> (Petrunkevitch, 1942)	Pa	Baltic amber
1152. <i>Gorgopsina marginata</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
1153. <i>Gorgopsina melanocephala</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
1154. <i>Gorgopsina naumanni</i> Giebel, 1856	Pa	Baltic amber
1155. <i>Gorgopsina paulula</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
1156. <i>Gorgopsina rectangularis</i> Wunderlich, 2011h	Pa	Baltic amber
1157. <i>Gorgopsina speciosa</i> Wunderlich, 2004aq	Pa	Baltic amber
<i>Heliophanus</i> C. L. Koch, 1833		Palaeogene – Recent
1158. <i>Heliophanus extinctus</i> Berland, 1939	Pa	Aix-en-Provence
<i>Hyllus</i> C. L. Koch, 1846		Quaternary – Recent
= † <i>Parevophrys</i> Petrunkevitch, 1942		
1159. <i>Hyllus succini</i> (Petrunkevitch, 1942)	Qt	Copal
Originally described as Baltic amber		
<i>Lyssomanes</i> Hentz, 1845		Neogene – Recent
1160. <i>Lyssomanes pristinus</i> Wunderlich, 1986	Ne	Dominican amber
i. = <i>Lyssomanes galianoae</i> Reiskind, 1989	Ne	Dominican amber
1161. <i>Lyssomanes pulcher</i> Wunderlich, 1988	Ne	Dominican amber
<i>Maevia</i> C. L. Koch, 1846		?Neogene – Recent
? <i>Maevia</i> sp. in Riquelme & Hill (2013)	Ne	Chiapas amber
† <i>Microlinus</i> Wunderlich, 2004aq		Palaeogene
1162. <i>Microlinus calidus</i> Wunderlich, 2004aq	Pa	Baltic amber
1163. <i>Microlinus folium</i> Wunderlich, 2004aq*	Pa	Baltic amber
<i>Myrmarachne</i> MacLeay, 1839		Quaternary – Recent

	= † <i>Entomocephalus</i> Holl, 1829 [suppressed; see ICZN Opinion 2258]	
1164.	<i>Myrmarachne formicoides</i> (Holl, 1829)	?Qt Copal [?not amber]
Neon Simon, 1876a	Quaternary – Recent
1165.	<i>Neon ?reticulatus</i> (Blackwall, 1853) [Recent]	Qt England
† Paralinus Petrunkevitch, 1942	Palaeogene
1166.	<i>Paralinus crosbyi</i> Petrunkevitch, 1942*	Pa Baltic amber
† Pensacolatus Wunderlich, 1988	Neogene
1167.	<i>Pensacolatus coxalis</i> Wunderlich, 1988*	Ne Dominican amber
1168.	<i>Pensacolatus spinipes</i> Wunderlich, 1988	Ne Dominican amber
1169.	? <i>Pensacolatus tibialis</i> Wunderlich, 2004aq	Ne Dominican amber
	<i>Pensacolatus</i> sp. in Wunderlich (1988)	Ne Dominican amber
Phidippus C. L. Koch, 1846	Palaeogene
1170.	<i>Phidippus impressus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1171.	<i>Phidippus pusillus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† Phlegrata Wunderlich, 1988	Neogene
1172.	<i>Phlegrata pala</i> Wunderlich, 1988*	Ne Dominican amber
† Prolinus Petrunkevitch, 1958	Palaeogene
1173.	<i>Prolinus fossilis</i> Petrunkevitch, 1958*	Pa Baltic amber
† Salticidites Straus, 1967	Neogene
1174.	<i>Salticidites hercynicus</i> Straus 1967*	Ne Willershausen
Sarinda Peckham & Peckham, 1892	Neogene – Recent
	? <i>Sarinda</i> sp. in Wunderlich (2004aq)	Ne Dominican amber
† Steneattus Bronn, 1856	Palaeogene
	= † <i>Leda</i> C. L. Koch & Berendt, 1854 [preoccupied]	
1175.	<i>Steneattus promissa</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
Thiodina Simon, 1900	Neogene
1176.	<i>Thiodina beugelorum</i> Wolff, 1990	Ne Dominican amber
Araneomorphae incertae sedis		
† Elvina Thorell, 1870b	Neogene
1177.	<i>Elvina antiqua</i> (von Heyden, 1859)	Ne Linz am Rhein
Araneae incerte sedis		
	Araneae incertae sedis in Selden et al. (2014)	P Kurty, Kazakhstan
† Amphiclotho Gourret, 1887	Palaeogene
1178.	<i>Amphiclotho breviuscula</i> Gourret, 1887*	Pa Aix-en-Provence
† Amphithomisus Gourret, 1887	Palaeogene
1179.	<i>Amphithomisus barbatus</i> Gourret, 1887*	Pa Aix-en-Provence
† Atocatle Feldmann, Vega, Applegate & Bishop, 1998	[really a spider?].....	Cretaceous
1180.	<i>Atocatle ranulfoi</i> Feldmann, Vega, Applegate & Bishop, 1998*	K Puebla, México
† Cercidiella Gourret, 1887	Palaeogene
1181.	<i>Cercidiella aquisextana</i> Gourret, 1887*	Pa Aix-en-Provence

† <i>Clubionella</i> Gourret, 1887	Palaeogene
1182. <i>Clubionella antiqua</i> Gourret, 1887*	Pa Aix-en-Provence
† <i>Eresoides</i> Gourret, 1887	Palaeogene
1183. <i>Eresoides orbicularis</i> Gourret, 1887*	Pa Aix-en-Provence
† <i>Hersilioides</i> Gourret, 1887	Palaeogene
1184. <i>Hersilioides thanatiformis</i> Gourret, 1887*	Pa Aix-en-Provence
† <i>Opistophylax</i> Menge, 1856	Palaeogene
1185. <i>Opistophylax exarata</i> Menge, 1856*	Pa Baltic amber
† <i>Prodysdera</i> Gourret, 1887	Palaeogene
1186. <i>Prodysdera intermedia</i> Gourret, 1887*	Pa Aix-en-Provence
† <i>Protochersis</i> Gourret, 1887	Palaeogene
1187. <i>Protochersis spinosus</i> Gourret, 1887*	Pa Aix-en-Provence
† <i>Protolachesis</i> Gourret, 1887	Palaeogene
1188. <i>Protolachesis annulata</i> Gourret, 1887*	Pa Aix-en-Provence
† <i>Paralycosa</i> Dunlop & Jekel, 2009	Palaeogene
= † <i>Protolycosa</i> Gourret, 1887 [preoccupied]	
1189. <i>Paralycosa attiformis</i> (Gourret, 1887)*	Pa Aix-en-Provence
† <i>Pseudothomisus</i> Gourret, 1887	Palaeogene
1190. <i>Pseudothomisus articulatus</i> Gourret, 1887*	Pa Aix-en-Provence
† <i>Schellenbergia</i> Heer, 1865	Neogene
1191. <i>Schellenbergia rotundata</i> Heer, 1865*	Ne Öhningen
† <i>Timeropus</i> Thorell, 1891	Palaeogene
= † <i>Lycosoides</i> Gourret, 1887 [preoccupied]	
1192. <i>Timeropus hersiliformis</i> (Gourret, 1887)*	Pa Aix-en-Provence

NOMINA DUBIA

Amaurobius C. L. Koch, 1837 [no currently valid fossil species]

1. *Amaurobius faustus* C. L. Koch & Berendt, 1854
2. *Amaurobius rimosus* C. L. Koch & Berendt, 1854

Auximus Simon, 1892 [now *Lathys* Simon, 1884: Dictynidae; no currently valid fossil species]

3. *Auximus fossilis* Petrunkevitch, 1950
4. *Auximus succini* Petrunkevitch, 1942

† *Clythia* C. L. Koch & Berendt, 1854 (*nomen dubium*)

5. *Clythia alma* C. L. Koch & Berendt, 1854*

† *Corynitoides* Dunlop & Jekel, 2009 (*nomen dubium*)= † *Corynitis* Menge in C. L. Koch & Berendt, 1854 [preoccupied]

6. *Corynitoides spinosa* (Menge in C. L. Koch & Berendt, 1854)*
7. *Corynitoides undulata* (Menge in C. L. Koch & Berendt, 1854)

† *Eocryphoea* Petrunkevitch, 1958 [also contains valid fossil species]

8. *Eocryphoea distincta* Petrunkevitch, 1950
9. *Eocryphoea fossilis* (Petrunkevitch, 1942)

- † *Eometa* Petrunkevitch, 1958 [also contains valid fossil species]
10. *Eometa aberrans* Petrunkevitch, 1958 Pa Baltic amber
 11. *Eometa robusta* Petrunkevitch, 1958 Pa Baltic amber
- Ero* C. L. Koch 1836 [also contains valid fossil species]
12. *Ero setulosa* C. L. Koch & Berendt, 1854 Pa Baltic amber
- † *Fictotama* Petrunkevitch, 1963 (*nomen dubium*) Palaeogene
13. *Fictotama extincta* Petrunkevitch, 1963* Ne Chiapas amber
- † *Memoratrix* Petrunkevitch, 1942 (*nomen dubium*) Palaeogene
- NB: Regarded by Wunderlich (2004p) as a possible pimoid or linyphiid
14. *Memoratrix rydei* Petrunkevitch, 1942 Pa Baltic amber
- † *Mimetarchaea* Eskov, 1992 Palaeogene
15. *Mimetarchaea gintaras* Eskov, 1992* Pa Baltic amber
- NB: Name based on a subadult male
- † *Miropholcus* Petrunkevitch, 1942 (*nomen dubium*) Palaeogene
- = † *Miropholcus* Petrunkevitch, 1942 [*lapsus*]
16. *Miropholcus heteropus* Petrunkevitch, 1942* Pa Baltic amber
- † *Perturbator* Petrunkevitch, 1971 (*nomen dubium*) Neogene
17. *Perturbator corniger* Petrunkevitch, 1971* Ne Chiapas amber
- † *Phalangopus* Menge in C. L. Koch & Berendt, 1854 (*nomen dubium*) Palaeogene
18. *Phalangopus subtilis* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- † *Praeoarces* Wunderlich, 2004q Palaeogene
19. *Praeoarces exitus* Wunderlich, 2004q* Pa Baltic amber
- Segestria* Latreille, 1804 [also contains valid fossil species]
20. *Segestria elongata* C. L. Koch & Berendt, 1854 Pa Baltic amber
 21. *Segestria nana* C. L. Koch & Berendt, 1854 Pa Baltic amber

NOMINA NUDA

- Amaurobius* C. L. Koch, 1837 [no currently valid fossil species]
1. *Amaurobius spinimanus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † *Anatone* Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*) Palaeogene
2. *Anatone hirsuta* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 3. *Anatone marginata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 4. *Anatone spinipes* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Aranea* Clerck, 1757 [now *Araneus* Clerck, 1757; which also contains valid fossil species]
5. *Aranea fossilis* Keferstein, 1834 Pa Aix-en-Provence
- Archaea* C. L. Koch & Berendt, 1854 [also contains valid fossil species]
6. *Archaea incomta* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 7. *Archaea sphinx* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † *Athera* Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*) Palaeogene
8. *Athera exilis* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Attus* Walckenaer, 1805 [now *Salicus* Latreille, 1804; no currently valid fossil species]

9. *Attus fossilis* Walckenaer, 1837 Pa Baltic amber
- Clubiona** Latreille, 1804 [also contains valid fossil species]
10. *Clubiona eseri* Heer, 1865 Ne Öhninge
 11. *Clubiona latifrons* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 12. *Clubiona parvula* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 13. *Clubiona pilosa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Clythia** C. L. Koch & Berendt, 1854 [also contains a *nomen dubium* fossil species]
14. *Clythia funesta* Koch & Berendt, 1854 Pa Baltic amber
 15. *Clythia gracilenta* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 16. *Clythia leptocarena* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Dielacata** Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*) Palaeogene
17. *Dielacata superba* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Drassus** Walckenaer, 1805 [now *Gnaphosa* Latreille, 1804; which also contains valid fossil species]
18. *Drassus oblongus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Dysdera** Latreille, 1804 [also contains valid fossil species]
19. *Dysdera hippopodium* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 20. *Dysdera glabrata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 21. *Dysdera scobiculata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 22. *Dysdera tenera* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Eolinus** Petrunkevitch, 1942 [also contains valid fossil species]
23. *Eolinus bitterfeldensis* Wunderlich, 2004aq Pa Baltic amber
 24. *Eolinus tystschenkoides* Wunderlich, 2004aq Pa Baltic amber
- Epeira** Walckenaer, 1805 [now *Araneus* Clerck, 1757; which also contains valid fossil species]
25. *Epeira eocaenica* Giebel, 1856 Pa Baltic amber
 26. *Epeira eocena* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Epeiridion** Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*) Palaeogene
27. *Epeiridion femoratum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Erithus** Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*) Palaeogene
28. *Erithus applanatus* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Ero** C. L. Koch & Berendt, 1836 [also contains valid fossil species]
29. *Ero coronata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 30. *Ero exculta* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 31. *Ero sphaerica* C. L. Koch & Berendt, 1854 Pa Baltic amber
 32. *Ero quadripunctata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Eyükselus** Özdişmen, 2007 (*nomen nudum*) Palaeogene
- = † *Propetes* Menge, 1854 [preoccupied]
33. *Eyükselus argutus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
 34. *Eyükselus felinus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
 35. *Eyükselus griseus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
 36. *Eyükselus latifrons* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
 37. *Eyükselus pumilus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
- Gea** C. L. Koch, 1843 [also contains valid fossil species]

38. *Gea pubescens* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † *Heteromma* Menge, 1856 (*nomen nudum*) Palaeogene
39. *Heteromma intersecta* Menge, 1856* Pa Baltic amber
- † *Idmonia* Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*) Palaeogene
40. *Idmonia virginea* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Melanophora* C. L. Koch, 1833** [now *Zelotes* Gistel, 1848; which also contains valid fossil species]
41. *Melanophora lepida* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
42. *Melanophora nitida* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Micaria* Westring, 1851** [also contains valid fossil species]
43. *Micaria ovata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
44. *Micaria squamata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
45. *Micaria tenuis* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Micryphantes* C. L. Koch, 1833** [also contains valid fossil species]
46. *Micryphantes globulus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
47. *Micryphantes turritus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † ***Mizalia* C. L. Koch & Berendt, 1854** [also contains valid fossil species]
48. *Mizalia truncata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † ***Ocia* Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** Palaeogene
49. *Ocia hirsuta* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Ocypete* C. L. Koch, 1836** [now *Heteropoda* Latreille, 1804; which also contains valid fossil species]
50. *Ocypete angustifrons* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
51. *Ocypete marginata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † ***Onca* Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** Palaeogene
52. *Onca lepida* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
53. *Onca pumila* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Philodromus* Walckenaer, 1826** [also contains valid fossil species]
54. *Philodromus griseus* Menge, 1856 Pa Baltic amber
55. *Philodromus marginatus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
56. *Philodromus reptans* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
57. *Philodromus redogradus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
58. *Philodromus spinipes* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Pythonissa* C. L. Koch, 1837** [now *Gnaphosa* Latreille, 1804; which also contains valid fossil species]
59. *Pythonissa bipunctata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
60. *Pythonissa discophora* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
61. *Pythonissa glabra* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
62. *Pythonissa villosa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Segestria* Latreille, 1804** [also contains valid fossil species]
63. *Segestria exarata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
64. *Segestria sulcata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
65. *Segestria undulata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † ***Siga* Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** Palaeogene
66. *Siga crinita* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber

- † *Spheconia* Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*) Palaeogene
 67. *Spheconia brevipes* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- † *Syphax* C. L. Koch & Berendt, 1854 [also contains valid fossil species]
 68. *Syphax hirtus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Theridium* Walckenaer, 1805** [now *Theridion* Walckenaer, 1805; which also contains valid fossil species]
 69. *Theridium bifurcum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 70. *Theridium chorius* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 71. *Theridium clavigerum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 72. *Theridium crassipes* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 73. *Theridium setulosum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Thomisus* Walckenaer, 1805** [also contains valid fossil species]
 74. *Thomisus matutinus* Menge, 1856 Pa Baltic amber
- † *Thyelia* C. L. Koch & Berendt, 1854 [also contains valid fossil species]
 75. *Thyelia mengei* Giebel, 1856 Pa Baltic amber
 76. *Thyelia pectinata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 77. *Thyelia spinosa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † *Zilla* C. L. Koch & Berendt, 1834 [also contains valid fossil species]
 78. *Zilla cornumana* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 79. *Zilla spinipalpa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

MISIDENTIFICATIONS

- Aranea* Clerck, 1757** [now *Araneus* Clerck, 1757; which also contains valid fossil species]
 1. *Aranea fusca pilosa* Bloch, 1776 [*nomen dubium*; non Araneae?] Qt Copal
- † ***Araneaovoivius* Dunlop & Braddy, 2011** [ichnogenus] Palaeogene
 2. *Araneaovoivius columbiae* (Scudder 1878)* [fossil egg sac] Pa Canada / USA
- † ***Archaeometa* Pocock, 1911** ?Devonian – Carb.
 3. ?*Archaeometa devonica* Størmer, 1976 [unidentifiable] D Alken an der Mosel
 4. *Archaeometa nephilina* Pocock, 1911* [not identified] C Coseley
- † ***Arachnometa* Petrunkevitch, 1949** Carboniferous
 5. *Arachnometa tuberculata* Petrunkevitch, 1949* [not identified] C Coseley
- † ***Eopholcus* Frič, 1904** Carboniferous
 6. *Eopholcus pedatus* Frič, 1904* [not identified] C Nýřany
- † ***Oichnus* Bromley 1981** [ichnogenus] Palaeogene
 7. *Oichnus bavincourtii* (Vaillant, 1909) [at one stage placed in *Cteniza*] Pa Northern France
- † ***Palpipes* Roth, 1854** Jurassic
 8. *Palpipes cursor* Roth, 1854 [crustacean] J Solnhofen
- † ***Palaeocteniza* Hirst, 1923** Devonian
 9. *Palaeocteniza crassipes* Hirst, 1923* [juvenile trigonotarbid?] D Rhynie chert
- † ***Pleurolycosa* Frič, 1904** Carboniferous
 10. *Pleurolycosa prolifera* (Frič, 1901)* [unidentifiable] C Nýřany

45,143 Recent species according to the WSC (2015)

HAPTOPODA

1 currently valid species of fossil haptopodid

† HAPTOPODA Pocock, 1911	Carboniferous
† PLESIOSIRONIDAE Pocock, 1911	Carboniferous
† Plesiosiro Pocock, 1911	Carboniferous
1. <i>Plesiosiro madeleyi</i> Pocock, 1911	C Coseley

no Recent species

AMBLYPYGI

11 currently valid species of fossil whip spider

AMBLYPYGI Thorell, 1882 Carbon. – Recent

= PHRYNÉIDES Walckenaer, 1837

= PHRYNICHIDA Petrunkevitch, 1945a

PALAEOAMBLYPYGI Weygoldt, 1996 (suborder) Carbon. – Recent

family uncertain

† **Sorellophrynxus Harvey, 2002** Carboniferous

= † *Protophrynxus* Petrunkevitch, 1913 (preoccupied)

1. *Sorellophrynxus carbonarius* (Petrunkevitch, 1913)* C Mazon Creek

† **Thelyphrynxus Petrunkevitch, 1913** Carboniferous

2. *Thelyphrynxus elongatus* Petrunkevitch, 1913 C Mazon Creek

PARACHARONTIDAE Weygoldt, 1996 Carbon. – Recent

† **Graeophonus Scudder, 1890b** Carboniferous

3. *Graeophonus anglicus* Pocock, 1911 C Coseley

4. *Graeophonus carbonarius* (Scudder, 1876)* C Cape Breton

5. *Graeophonus scudderi* Pocock, 1911 C Mazon Creek

† **Paracharonopsis Engel & Grimaldi, 2014** Palaeogene

6. *Paracharonopsis cambayensis* Engel & Grimaldi, 2014* Pa Cambay amber

EUAMBLYPYGI Weygoldt, 1996 (suborder) Cretaceous – Recent

CHARINIDAE Quintero, 1986 Recent

no fossil record

NEOAMBLYPYGI Weygoldt, 1996 (infraorder) Cretaceous – Recent

CHARONTIDAE Simon, 1892a Recent

no fossil record

UNIDISTITARSATA Engel & Grimaldi, 2014 Cretaceous – Recent

† **Kronocharon Engel & Grimaldi, 2014** Cretaceous

7. *Kronocharon prendinii* Engel & Grimaldi, 2014* K Burmese amber

PHRYNOIDEA Blanchard, 1852 Cretaceous – Recent

PHRYNICHIDAE Simon, 1892a Recent

no fossil record

PHRYNIDAE Blanchard, 1852	Cretaceous – Recent
= † ELECTROPHRYNIDAE Petrunkevitch, 1971	
† Britopygus Dunlop & Martill, 2002	Cretaceous
8. <i>Britopygus weygoldti</i> Dunlop & Martill, 2002	K Crato Formation
† Electrophrynus Petrunkevitch, 1971	Neogene
9. <i>Electrophrynus mirus</i> Petrunkevitch, 1971	Ne Chiapas amber
Phrynus Lamarck, 1801	Neogene – Recent
10. <i>Phrynus mexicana</i> Poinar & Brown, 2004	Ne Chiapas amber
11. <i>Phrynus resinae</i> (Schawaller, 1979b)	Ne Dominican amber

NOMEN DUBIUM

1. <i>Phrynus fossilis</i> Keferstein, 1834	Pa Aix-en-Provence
i. = <i>Phrynus marioni</i> Gourret, 1887	Pa Aix-en-Provence

136 Recent species according to Harvey (2003)

UROPYGI

7 currently valid species of fossil whip scorpion

UROPYGI Thorell, 1882 **Carbon. – Recent**

= THELYPHONIDA Latreille, 1804b
 = UROTRICHA C. L. Koch, 1851
 = OXOPOEI Thorell, 1888
 = HOLOPELTIDIA Börner, 1902

Thelyphonida sp. *in* Selden et al. 2014 C Donets Basin

plesion genera

† **Geralinura Scudder, 1884** **Carboniferous**

1. *Geralinura britannica* Pocock, 1911 C Coseley
2. *Geralinura carbonaria* Scudder, 1884* C Mazon Creek
 - i. = *Geralinura gigantea* Petrunkevitch, 1913 C Mazon Creek
 - ii. = *Geralinura similis* Petrunkevitch, 1913 C Mazon Creek

† **Parageralinura Tetlie & Dunlop, 2008** **Carboniferous**

3. *Parageralinura naufraga* (Brauckmann & Koch, 1983) C Hagen-Vorhalle
4. *Parageralinura neerlandicus* Laurentiaux-Viera & Laurentiaux, 1961 C Limburg

† **Proschizomus Dunlop & Horrocks, 1996** **Carboniferous**

5. *Proschizomus petrunkevitchi* Dunlop & Horrocks, 1996 C Coseley

† **Prothelyphonus Frič, 1904** **Carboniferous**

6. *Prothelyphonus bohemicus* (Kušta, 1884b) C Rakovník
 - i. = *Prothelyphonus cordai* Frič, 1904 C Rakovník
 - ii. = *Geralinura crassa* Kušta, 1888 C Rakovník
 - iii. = *Geralinura noctua* Kušta, 1888 C Rakovník
 - iv. = *Geralinura scudderi* Kušta, 1888 C Rakovník

THELYPHONIDAE Lucas 1835 **Cretaceous – Recent**

† **Mesoproctus Dunlop, 1988** **Cretaceous**

7. *Mesoproctus rowlandi* Dunlop, 1998 K Crato Formation
- Mesoproctus* sp. *in* Dunlop & Martill (2002) K Crato Formation

MISIDENTIFICATIONS

1. *Thelyphonus hadleyi* Pierce, 1945 [unidentifiable, ?algal] Ne California

SCHIZOMIDA

6 currently valid species

- the fossil family Calcitronidae cannot be meaningfully compared to the Recent families

SCHIZOMIDA Petrunkevitch, 1945b Palaeogene – Recent

= TARTARIDES Thorell, 1888 (tribe)
 = COLOPYGA Cook, 1899 (order)
 = SCHIZOPELTIDA Börner, 1902 (tribe)

† **CALCITRONIDAE Petrunkevitch, 1945b** Palaeogene – Neogene

† **Calcitro Petrunkevitch, 1945b** Palaeogene – Neogene

1. *Calcitro fisheri* Petrunkevitch, 1945b* Ne Onyx Marble
2. *Calcitro oplonis* Lin in Lin et al., 1988 Pa Shandong, China

HUBBARDIIDAE Cook, 1899 Neogene – Recent

Antilostenochrus Armas and Teruel, 2002 Neogene – Recent

3. *Antilostenochrus pseudoannulatus* (Krüger & Dunlop, 2010) Ne Dominican Amber

† **Calcoschizomus Pierce, 1951** Neogene

4. *Calcoschizomus latisternum* Pierce, 1951 Ne Onyx Marble

† **Onychothelyphonus Pierce, 1950** Neogene

5. *Onychothelyphonus bonneri* Pierce, 1950 Ne Onyx Marble

Rowlandius Reddell & Cokendolpher, 1995 Neogene – Recent

6. *Rowlandius velteni* (Krüger & Dunlop, 2010) Ne Dominican Amber

PROTOSCHIZOMIDAE Rowland, 1975 Recent

no fossil record

References

- Absolon, K. & Kratochvíl, J. 1932. Zur Kenntnis der höhlenbewohnenden Araneeae der illyrischen Karstgebiete. *Mitteilungen über Höhlen- und Karstforschung*, 3: 73–81.
- Agassiz, L. 1844. *Monographie des poisons fossiles du Vieux Gres Rouge ou Systeme Devonian*. Neufchatel, folio: 171 pp.
- Allen, J. G. & Feldman, R. M. 2005. *Panduralimulus babcocki* n. gen. and sp., a new Limulacean horseshoe crab from the Permian of Texas. *Journal of Paleontology*, 79: 594–600.
- Ambrose, T. & Romano, M. 1972. New Upper Carboniferous Chelicerata (Arthropoda) from Somerset, England. *Palaeontology* 15: 569–578.
- Ambrus, B. & Hably, L. 1979. *Eriophyes daphnogene* sp. n. a fossil gall from the Upper Oligocene of Hungary. *Annales Historico-Naturales Musei Nationalis Hungarici*, 71: 55–56.
- Amerling, C. 1862. Naturökonomie der von ihm beobachteten Milben, insbesondere der Trombidieen. *Sitzungsberichte der Königlich Böhmischen Gesellschaft der Wissenschaften in Prague*, 2: 54–56.
- Ammon, L. von 1901. Ueber *Anthracomartus* aus dem Pfälzischen Carbon. *Geognostische Jahresshefte*, 13: 1–6.
- Anderson, L. I., Dunlop, J. A. & Trewin, N. H. 2000. A Middle Devonian chasmataspid arthropod from Achansarras Quarry, Caithness, Scotland. *Scottish Journal of Geology*, 36: 151–158.
- Andrée, K. 1913. Ueber *Anthracophrynx tuberculatus* nov. gen. nov. spec. aus dem productiven Karbon von Dudweiler im Saar-Revier, nebst einer Liste der bisher im Karbon Deutschland gefundenen Arachnoiden-reste. *Jahres-Bericht und Mitteilungen der Oberrheinischen Geologischen Vereins*, 3: 89–93.
- Aoki, J. 1965. Oribatiden (Acarina) Thailands. I. *Nature and Life in Southeast Asia*, 4: 129–193.
- Aoki, J. 1966a. A remarkable new oribatid mite from South Japan (Cryptostigmata: Tokunocepheidae, fam. nov.). *Acarologia*, 8: 358–364.
- Aoki, J. 1966b. Epizoic symbiosis: an oribatid mite, *Symbioribates papuensis*, representing a new family, from cryptogamic plants growing on backs of Papuan weevils (Acari: Cryptostigmata). *Pacific Insects*, 8: 281–289.
- Aoki, J. 1974. [On the fossil mites in Mizunami amber from Gifu Prefecture, Central Japan.] *Bulletin of the Mizunami Fossil Museum*, 1: 397–399 [in Japanese with English summary].
- Aoki, J. 1976. Oribatid mites from the IBP Study Area, Pasoh Forest Reserve, West Malaysia. *Nature and Life in Southeast Asia*, 7: 39–59.
- Aoki, J., Takaku, G. & Ito, F. 1994. Aribatidae, a new myrmecophilous oribatid mite family from Java. *International Journal of Acarology*, 20: 3–10.

- Arillo, A. & Subías, L.S. 2000. A new fossil oribatid mite, *Arachaeocheistes minguezae* n. gen. n. sp. from Spanish Lower Cretaceous amber. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 84: 231–236.
- Arillo, A. & Subías, L.S. 2002. Second fossil oribatid mite from the Spanish Lower Cretaceous amber. *Eupterogaeus bitranslammellatus* n. sp. (Acariformes, Oribatida, Cepheidae). *Acarologia*, 42: 403–406.
- Arillo, A., Subiás, L. S. & Shtanchaeva, U. 2008. A new fossil oribatid mite, *Ommatocepheus nortoni* sp. nov. (Acariformes, Oribatida, Cepheidae), from a new outcrop of Lower Cretaceous Álava amber (northern Spain). *Systematic and Applied Acarology*, 13: 252–255.
- Arillo, A., Subiás, L. S. & Shtanchaeva, U. 2009. A new fossil species of oribatid mite, *Ametroproctus valeriae* sp. nov. (Acariformes, Oribatida, Ametroproctidae), from the Lower Cretaceous amber of San Just, Teruel Province, Spain. *Cretaceous Research*, 30: 322–324.
- Armas L. F. de & Teruel, R. 2002. Un género nuevo de Hubbardiidae (Arachnida: Schizomida) de las Antillas Mayores. *Revista Ibérica de Aracnología*, 6: 45–52.
- Atyeo, W. T. & Baker, E. W. 1964. Tarsocheylidae, a new family of prostigmatic mites (Acarina). *Bulletin of the University of Nebraska State Museum*, 4: 243–256.
- Atyeo W. T. & Gaud, J. 1979. Ptyssalgidae, a new family of analgoid feather mites (Acarina, Acaridida). *Journal of Medical Entomology*, 16: 306–308.
- Atyeo, W. T. & Peterson, P. C. 1972. The feather mite family Alloptidae Gaud, new status, I. The subfamilies Trouessartiinae Gaud and Thysanocercinae, new subfamily (Analgoidea). *Zoologischer Anzeiger*, 188: 56–60.
- Atyeo W. T., Baker, E. W. & Delfinado M. D. 1974. *Gaudiella minuta*, a new genus and species of mite (Acarina: Acaridia) belonging to the new family Gaudiellidae. *Journal of the Washington Academy of Sciences*, 64: 295–298.
- Audouin, V. 1826. Explication sommaire des planches d'arachnides de l'Égypte et de la Syrie. In *Description de l'Égypt ou Recueil des Observations et des Recherches qui ont été Faites en Égypte Pendant l'Expédition de l'Armée Française, 1st edition*, 1(4), 99–186. C. L. F. Panckoucke, Paris.
- Ausserer, A. 1867. Die Arachniden Tirols nach ihrer horizontalen und verticalen Verbreitung; 1. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 17: 137–170.
- Ausserer, A. 1875. Zweiter Beitrag zur Kenntniss der Arachniden-Familie der Territelariae Thorell (Mygalidae Autor). *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 25: 125–206.
- Ayyildiz, N. & Luxton, M. 1989. Epimerellidae (Acari, Oribatida), a new mite family. *Journal of Natural History*, 23: 1381–1386.
- Baily, W. H. 1863. Remarks on some Coal Measures Crustacea belonging to the genus *Belinurus*, König, with description of two new species from Queen's County, Ireland. *Annals and Magazine of Natural History*, 11: 107–114.

- Baily, W. H. 1869. On fossils obtained at Kiltorcan Quarry, Co. Kilkenny. *British Association Report*, pp. 73–75.
- Baker, E. W. 1949. Pomerantziidae, a new family of prostigmatic mites. *Journal of the Washington Academy of Science*, 39: 269–271.
- Baker, E. W. & Pritchard, A. E. 1953. The family categories of tetranychoid mites, with a review of the new families Linotetranidae and Tuckerellidae. *Annals of the Entomological Society of America*, 46: 243–258.
- Baldwin W. & Sutcliffe, W. H. 1904. *Eoscorpius sparthensis* n. sp. from the Middle Coal Measures of Lancashire. *Quarterly Journal of the Geological Society of London*, 60: 395–398.
- Balogh, J. 1958. Oribatides nouvelles de l'Afrique tropicale. *Revue Zoologie Botanique Africaines*, 58: 1–34.
- Balogh, J. 1968. New oribatids (Acari) from New Guinea. *Acta Zoologica Academiae Scientiarum Hungaricae*, 14: 259–285.
- Balogh, J. 1970. New oribatids (Acari) from New Guinea. II. *Acta Zoologica Academiae Scientiarum Hungaricae*, 16: 291–344.
- Balogh, J. 1972. *The oribatid genera of the world*. Akadémiai Kiadó, Budapest, 188 pp.
- Balogh, J. 1983. A partial revision of the Oppidae Grandjean, 1954 (Acari: Oribatei). *Acta Zoologica Academiae Scientiarum Hungaricae*, 29: 1–79.
- Balogh, J. & Balogh, P. 1984. A review of the Oribatuloidea Thor, 1929 (Acari: Oribatei). *Acta Zoologica Hungarica*, 30: 257–313.
- Balogh, J. & Balogh, P. 1992. *The oribatid mites genera of the world. I*. Hungarian National Museum Press, 263 pp.
- Balzan, L. 1888. *Chernetidae Nonnullae Sud-Americanae*, III. Asuncion.
- Balzan, L. 1892. Voyage de M. E. Simon au Venezuela (Décembre 1887 – Avril 1888). Arachnides. Chernetes (Pseudoscorpiones). *Annales de la Société Entomologique de France*, 60: 497–552.
- Bamber, R. 2007. A holistic re-interpretation of the phylogeny of the Pycnogonida Latreille, 1810 (Arthropoda). *Zootaxa*, 1668: 295–312.
- Banks, N. 1892a. A new genus of Phalangiidae. *Proceedings of the Entomological Society of Washington*, 2(2): 249–251.
- Banks, N. 1893. The Phalanginae of the United States. *The Canadian Entomologist*, 25: 205–211.
- Banks, N. 1895. Notes on the Pseudoscorpionida. *Journal of the New York Entomological Society*, 3: 1–13.
- Banks, N. 1896. New North American spiders and mites. *Transactions of the American Entomological Society*, 23: 57–77.
- Banks, N. 1905. Arachnids from the Cocos Island. *Proceedings of the Entomological Society of Washington*, 7: 20–23.
- Barbour, E. H. 1914. Carboniferous eurypterids of Nebraska. *American Journal of Science*, 4th Series, 38: 507–510.

- Beecher, C. E. 1902. Note on a new xiphosuran from the Upper Devonian of Pennsylvania. *American Geologist*, 29, 143–146.
- Beecher, C. E. 1904. Note on a new Permian xiphosuran from Kansas. *American Journal of Science*, 4th Series, 17: 23–24.
- Beier, M. 1932a. Pseudoscorpionidea I. Subord. Chthoniinea et Neobisiinea. *Tierreich*, 57: i–xx, 1–258.
- Beier, M. 1932b. Pseudoscorpionidea II. Subord. C. Cheliferina. *Tierreich*, 58: i–xxi, 1–294.
- Beier, M. 1937. Pseudoscorpione aus dem baltischen Bernstein. *Festschrift zum 60. Geburtstag von Professor Dr. Embrik Strand, Riga*, 2: 302–316.
- Beier, M. 1947a. Pseudoscorpione im Baltischen Bernstein und die Untersuchung von Bernstein-Einschlüssen. *Mikroskopie, Wien*, 1: 188–199.
- Beier, M. 1947b. Zur Kenntnis der Pseudoscorpionidenfauna des südlichen Afrika, insbesondere der südwest und südafrikanischen Trockengebiet. *Eos, Madrid*, 23: 285–339.
- Beier, M. 1955. Pseudoscorpione im baltischen Bernstein aus dem Geologischen Staatsinstitut in Hamburg. *Mitteilungen aus dem Mineralogisch-Geologischen Staatsinstitut in Hamburg*, 25: 48–54.
- Bell, W. A. 1922. A new genus of Characeae and new Merostomata from the Coal Measures of Nova Scotia. *Transactions of the Royal Society of Canada*, 4: 159–167.
- Bergström, J., Stürmer, W. & Winter, G. 1980. *Palaeoisopus*, *Palaeopantopus* and *Palaeothea*, pycnogonid arthropods from the Lower Devonian Hunsrück Slate, West Germany. *Paläontologische Zeitschrift*, 54: 7–54.
- Berland, L. 1913. Araignées. In *Mission du Service géographique de l'armée pour la mesure d'un arc du méridien équatorial en Amérique du Sud (1899-1906)*. Paris, 10: 78–119.
- Berland, L. 1939. Description de quelques Araignées fossiles. *Revue Française d'Entomologie*, 6: 1–9.
- Berlese, A. 1885. Acarorum Systematis. *Bullettino della Società Entomologica Italiana*, 17: 121–135.
- Berlese, A. 1888. Acari Austro-Americani quos collegit Aloysius Balzan. Manipulus primus. Species novas circiter quinquaginta complectens. *Bullettino della Società Entomologica Italiana*, 20: 171–222.
- Berlese, A. 1896. Acari, Myriapoda et Scorpiones hucusque in Italia reperta. *Acari, Myriapoda et Scorpiones in Italia reperta*, Fasc. 79, 15 pp., 6 pls.
- Berlese, A. 1899. Gli acari agrarii. Puntat II. *Rivista di Patologia Vegetale*, Padova, 7: 312–344.
- Berlese, A. 1908. Elenco di generi e specie nuove di acari. *Redia*, 5: 1–15.
- Berlese, A. 1910. Lista di nuove specie e nuovi generi di Acari. *Redia*, 6: 242–271.
- Berlese, A. 1914. Acari nuovi. *Redia*, 10: 1–150.
- Berlese, A. 1923. Centuria sesta di Acari nuovi. *Redia*, 15: 237–262.
- Bernini, F. 1975. Notulae Oribatologicae XII. Una nuova specie di *Carabodes* affine a *C. minusculus* Berlese 1923 (Acarida, Oribatei). *Redia* 56: 455–471.
- Bertkau, P. 1872. Über die Respirationsorgane der Araneen. *Archiv für Naturgeschichte*, 38: 208–233.

- Bertkau, P. 1878a. Versuch einer natürlichen Anordnung der Spinnen, nebst Bemerkungen zu einzelnen Gattungen. *Archiv für Naturgeschichte*, 44: 351–410.
- Bertkau, P. 1878b. Einige Spinnen und ein Myriapode aus der Braunkohle von Rott. *Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande und Westfalens, Bonn*, 35: 346–360.
- Bertkau, P. 1882. Ueber das Cribellum und Calamistrum. Ein Beitrag zur Histologie, Biologie und Systematik der Spinnen. *Archiv für Naturgeschichte*, 48: 316–362.
- Beyschlag, F. & Fritsch, K. von 1899. Das jüngere Steinkohlengebirge und das Rothliegende in der Provinz Sachsen und den angrenzenden Gebieten. *Abhandlungen der Königlich Preussischen geologischen Landesanstalt*, 10: 1–263.
- Blackwall, J. 1833. Characters of some undescribed genera and species of Araneidae. *London philosophical Magazine and Journal of Science*, 3: 104–112, 187–197, 344–352, 436–443.
- Blackwall, J. 1834a. Characters of some undescribed species of Araneidae. *London philosophical Magazine and Journal of Science*, 5: 50–53.
- Blackwall, J. 1834b. *Researches in Zoology*. London, pp. 229–433.
- Blackwall, J. 1841. The difference in the number of eyes with which spiders are provided proposed as the basis of their distribution into tribes; with descriptions of newly discovered species and the characters of a new family and three new genera of spiders. *Transactions of the Linnean Society, London*, 18: 601–670.
- Blackwall, J. 1853. Descriptions of some newly discovered species of Araneida. *Annals and Magazine of Natural History, series 2*, 11: 14–25.
- Blackwall, J. 1859. Descriptions of newly discovered spiders captured by James Yate Johnson Esq., in the island of Maderia. *Annals and Magazine of Natural History, series 3*, 4: 255–267.
- Blackwall, J. 1862. Descriptions of newly-discovered spiders from the island of Madeira. *Annals and Magazine of Natural History, series 3*, 9: 370–382.
- Blackwall, J. 1864. *A History of the Spiders of Great Britain and Ireland. Part II*. The Ray Society, London, 1864 pp. 175–384.
- Blackwall, J. 1870. Notes on a collection of spiders made in Sicily in the spring of 1868, by E. Perceval Wright, M.D., with a list of the species, and descriptions of some new species and of a new genus. *Annals and Magazine of Natural History, series 4*, 5: 392–405.
- Blanchard, E. 1852. Arachnides. In *L'organisation du règne animal, 2nd Edition*, vol. 2. E. Blanchard, Paris.
- Blaszak, J., Cokendolpher, J. C. & Polyak, V. J. 1995. *Paleozemercon cavernicolous*, a new genus and new species of fossil mite from a cave in the southwestern U.S.A. (Acari, Gamasida: Zerconidae). *International Journal of Acarology*, 21: 253–259.
- Bleicher, M. 1897. Sur la découverte d'une nouvelle espèce de limule dans les marnes irisées de Lorraine. *Bulletin de la Société des Sciences de Nancy*, (2)14: 116–126.

- Bloch, M. [E.] 1776. Naturgeschichte des Kopals. *Beschäftigungen der Berlinischen Gesellschaft Naturforschender Freunde*, 2: 91–196.
- Bode, A. 1951. Ein Liassischer Skorpionide. *Palaeontologische Zeitschrift*, 24: 58–65.
- Bolland, H. R. & Magowski, W. Ł. 1990. *Neophyllobius succineus* n. sp. from Baltic amber (Acari: Raphignathoidea: Camerobiidae). *Entomologische Berichten*, 50: 17–21.
- Bosselaers, J. 2004. A new *Garacops* species from Madagascar copal (Araneae: Selenopidae). *Zootaxa*, 445: 1–7.
- Bosselaers, J., Dierick, M., Cnudde, V., Masschaele, B., Van Hoorebeke, L. & Jacobs, P. 2010. High resolution X-ray computed tomography of an extant new *Donaea* (Araneae: Liocranidae) species in Madagascan copal. *Zootaxa*, 2427: 25–35.
- Bottali, P. 1975. Note su due rari esemplari di Araneidi (Aracnidi) rinvenuti nei depositi diatomitici (facies lacustre) di Riano Flaminio (Roma). *Fragmenta entomologica*, 11: 169–174.
- Braddy, S. J., Aldridge, R. J. & Theron, J. N. 1995. A new eurypterid from the Late Ordovician Table Mountain Group, South Africa. *Palaeontology*, 38: 563–581.
- Braddy, S. J., Selden, P. A. & Doan Nhat T. 2002. A new carcinosomatid eurypterid from the Upper Silurian of Northern Vietnam. *Palaeontology*, 45: 897–915.
- Bradley, W. H. 1931. Origin and microfossils of the oil shale of the Green River Formation. *United States Geological Survey, Professional Paper*, 168: 1–58.
- Brauckmann, C. 1982. Der Schwertschwanz *Euproops* (Xiphosuraida, Limulina, Euproopacea) aus dem Ober-Karbon des Piesbergs bei Osnabrück. *Osnabücker naturwissenschaftliche Mitteilungen*, 9: 17–26.
- Brauckmann, C. 1984. Eine neue Arachniden-Art aus dem Westfalium des Saargebietes (West-Deutschland). *Dortmunder Beiträge zur Landeskunde, naturwissenschaftliche Mitteilungen*, 18: 95–103.
- Brauckmann, C. 1987. Neue Arachniden (Ricinuleida, Trigonotarbida) aus dem Namurium B von Hagen-Vorhalle (Ober-Karbon; West-Deutschland). *Dortmunder Beiträge der Landeskunde, naturwissenschaftliche Mitteilungen*, 21: 97–109.
- Brauckmann, C. & Koch, L. 1983. *Prothelyphonus naufragus* n. sp., ein neuer Geisselskorpion [Arachnida: Thelyphonida: Thelyphonidae] aus dem Namurium unteres Oberkarbon von West-Deutschland. *Entomologica Germania*, 9: 63–74.
- Brauckmann, C., Koch, L. & Kemper, M. 1985. Spinnentiere (Arachnida) und Insekten aus den Vorhalle-Schichten (Namurian B; Ober-Karbon) von Hagen-Vorhalle (West-Deutschland). *Geologie und Paläontologie in Westfalen*, 3: 1–131.
- Brauer, F., Redtenbacher, J. & Ganglbauer, L. 1889. Fossile Insekten aus der Juraformation Ost-Siberiens. *Mémoires de l'Académie Impériale des Sciences de St.-Petersbourg, VII serie*, 36(15): 1–22.
- Braun, C. F. W. 1860. Die Thiere in den Pflanzenschifern der Gegend von Bayreuth. Programm zum Jahresbericht der Königl. Kreis-Landwirtschafts- und Gewerbschule zu Bayreuth für das Schuljahr

- 1859/60. *Jahresbericht von der Königl. Kreis-Landwirtschafts- und Gewerbschule zu Bayreuth für das Schuljahr 1859/60*: 11 pp.
- Brescovit, A. D. 1997. Revisão de Anyphaeninae Bertkau a nível de gêneros na região neotropical (Araneae, Anyphaenidae). *Revista Brasileira de Zoologia*, 13: 1–187.
- Briggs, D. E. G. & Collins, D. 1988. A Middle Cambrian chelicerate from Mount Stephen, British Columbia. *Palaeontology*, 31: 779–798.
- Briggs, D. E. G., Siveter, D. J., Siveter, D. J., Sutton, M. D., Garwood, R. J. & Legg, D. 2012. Silurian horseshoe crab illuminates the evolution of arthropod limbs. *Proceedings of the National Academy of Sciences of the United States of America*, 109: 15702–15703.
- Briggs, T. A. 1971. Relict harvestmen from the Pacific northwest (Opiliones). *Pan-Pacific Entomologist*, 74: 165–178.
- Bristowe, W. S. 1938. The classification of spiders. *Proceedings of the Zoological Society of London*, 108: 285–322.
- Bristowe, W. S. 1939. *The comity of spiders. Volume 1*. London, 228 pp.
- Brongniart, C. 1877. Note sur une Aranéide fossile des terrains tertiaires. *Annales de la Société Entomologique de France*, (5) 7: 221–224.
- Bruce, W. A. & Johnston, D. E. 1976. *Gaudoglyphus* n. gen., based on *Analges minor* Nörner (Acaria: Gaudoglyphidae n. fam.). *International Journal of Acarology*, 2: 29–33.
- Broili, F. 1928. Crustaceenfunde aus dem rheinischen Unterdevon. I. Über Extremitätenreste. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung*, 1928: 197–201.
- Broili, F. 1930. Über ein neues Exemplar von *Palaeopantopus*. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung*, 1930: 209–214.
- Bromley, R.G. 1981. Concepts in ichnotaxonomy illustrated by small round holes in shells. *Acta Geologica Hispànica*, 16: 55–64.
- Bronn, H. G. 1856. *Lethaea Geognostica oder Abbildung und Beschreibung für die Gebirgs-Formationen bezeichnendsten Versteinerungen. Dritter Band*. Schweizerbart'sche Verlagshandlung und Druckerei 1853–1856, pp. 622–639.
- Buckland, W. 1837. *The Bridgewater treatises on the power, wisdom and goodness of God as manifested in the creation. Treatise IV. Geology and mineralogy with reference to natural theology. 2nd Edition*. William Pickering, London.
- Bulanova-Zachvatkina, E. M. 1974. [New genera of oribatid mites from the Upper Cretaceous of Tajmyr.] *Paleontological Journal*, 1974: 141–144. [In Russian]
- Burmeister, H. 1843. *Die Organisation der Trilobiten, aus ihren lebenden Verwandten entwickelt; nebst systematischen Uebersicht aller zeither beschrieben Arten*. G. Reimer, Berlin, 148 pp.

- Cambridge, F. O. P.- 1893. Handbook to the study of British spiders (Drassidae and Agelenidae). *British Nature Supplement*, 3: 117–170.
- Cambridge, F. O. P.- 1899. Arachnida. Araneida. Vol. 2. *Biologia Centrali-Americanana*: pp. 41–88.
- Cambridge, O. P.- 1870. Descriptions and sketches of two new species of Araneida, with characters of a new genus. *Journal of the Linnean Society of London*, 10: 398–405.
- Cambridge, O. P.- 1871. Arachnida (1870). *The Zoological Record*, 7: 207–224.
- Cambridge, O. P.- 1873. On some new genera and species of Araneida. *Proceedings of the Zoological Society of London*, 1873: 112–129.
- Cambridge, O. P.- 1874. On some new genera and species of Araneida. *Annals and Magazine of Natural History, series 4*, 14: 169–183.
- Cambridge, O. P.- 1876. On a new order and some new genera of Arachnida from Kerguelen's Land. *Proceedings of the Zoological Society of London*, 1876: 258–265.
- Cambridge, O. P.- 1877. On some new species of Araneida, with characters of two new genera and remarks on the families Podophthalmides and Dinopides. *Proceedings of the Zoological Society of London*, 1877: 557–578.
- Cambridge, O. P.- 1879a. On some new and rare spiders from New Zealand, with characters of four new genera. *Proceedings of the Zoological Society of London*, 1879: 681–703.
- Cambridge, O. P.- 1879b. On some new and rare British spiders, with characters of a new genus. *Annals and Magazine of Natural History*, 4: 190–215.
- Cambridge, O. P.- 1881. On some new genera and species of Araneidea. *Proceedings of the Zoological Society of London*, 1881: 765–775.
- Cambridge, O. P.- 1882a. On new genera and species of Araneidea. *Proceedings of the Zoological Society of London*, 1882: 423–442.
- Cambridge, O. P.- 1882b. Arachnida (1881). *The Zoological Record*, 18: 1–32.
- Cambridge, O. P.- 1894. Arachnida. Araneida. Vol. 1. *Biologia Centrali-Americanana*: pp. 121–144.
- Cambridge, O. P.- 1895. Arachnida. Araneida. Vol. 1. *Biologia Centrali-Americanana*: pp. 145–160.
- Cambridge, O. P.- 1898. Arachnida. Araneida. Vol. 1. *Biologia Centrali-Americanana*: pp. 233–288.
- Cambridge, O. P.- 1902. On new and rare British Arachnida. *Proceedings of the Dorset Natural History and Antiquarian Field Club*, 23: 16–40.
- Camin, J. H. & Gorirossi, F. E. 1955. A revision of the suborder Mesostigmata (Acarina), based on new interpretations of comparative morphological data. *Chicago Academy of Sciences Special Publication*, 11: 1–70.
- Camin J. H., Moss W. W. & Oliver J. H. 1967. Cloacaridae, a new family of cheyletoid mites from the cloaca of aquatic turtles. *Journal of Medical Entomology*, 4: 261–272.

- Campos, D. R. B. 1986. Premeiro registro fóssil de Scorpinoidea na Chapada do Araripe (Cretáceo Inferior), Brasil. *Anais do Academia Brasileira dos Ciências*, 58: 135–137.
- Canestrini, G. & Fanzago, F. 1877. Intorno agli Acari italiani. - Atti del R. Istituto Veneto Scienze, Lettere ed Arti, Ser. 5 4: 69–208.
- Canestrini, G. & Pavesi, P. 1870. Catalogo sistematico degli Araneida italiani. *Archivo per la zoologia, l'anatomia e la fisiologia*, (2)2: 1–44.
- Caporiacco, L. di 1949. Aracnidi della colonia de Kenya raccolti da Toschi e Meneghetti negli anni 1944–1946. *Commentationes Pontificiae Academiae Scientiarum*, 13: 309–492.
- Carvalho, M. P. G. de & Lourenço, W. R. 2001. A new family of fossil scorpions from the Early Cretaceous of Brazil. *Comptes Rendus de l'Académie de Sciences de Paris, Earth and Planetary Sciences*, 332: 711–716.
- Caster, K. E. & Brooks, H. K. 1956. New fossils from the Canadian–Chazan (Ordovician) hiatus in Tennessee. *Bulletins of American Palaeontology*, 36: 157–199.
- Caster, K. E. & Kjellesvig-Waering, E. N. 1953. *Melbournopterus*, a new Silurian eurypterid from Australia. *Journal of Paleontology*, 27: 153–156.
- Caster, K. E. & Kjellesvig-Waering, E. N. 1955. *Marsupipterus*, an unusual eurypterid from the Downtonian of England. *Journal of Paleontology*, 29: 1040–1041.
- Caster, K. E. & Kjellesvig-Waering, E. N. 1956. Some notes on the genus *Dolichopterus* Hall. *Journal of Paleontology*, 30: 19–28.
- Caster K. E. & Kjellesvig-Waering, E. N. 1964. Upper Ordovician eurypterids of Ohio. *Palaeontographica Americana*, 4 (32): 297–358.
- Chamberlin, J. C. 1923a. The genus *Pseudogarypus* Ellingsen (Pseudoscorpionida – Feallidae). *Entomological News*, 34: 146–149, 161–166.
- Chamberlin, J. C. 1923b. New and little known pseudoscorpions, principally from the islands and the adjacent shores of the Gulf of California. *Proceedings of the California Academy of Science*, (4)12: 353–387.
- Chamberlin, J. C. 1929. A synoptic classification of the false scorpions or chela-spinners, with a report on a cosmopolitan collection of the same. Part I. The Heterosphyronida (Chthoniidae) (Arachnida-Chelonethida). *Annals and Magazine of Natural History*, series 10, 4: 50–80.
- Chamberlin, J. C. 1930. A synoptic classification of the false scorpions or chela-spinners, with a report on a cosmopolitan collection of the same. Part II. The Diplosphyronida (Arachnida-Chelonethida). *Annals and Magazine of Natural History*, series 10, 5: 1–48, 585–620.
- Chamberlin, J. C. 1931a. The arachnid order Chelonethida. *Stanford University Publications, Biological Sciences*, 7: 1–284.
- Chamberlin, J. C. 1931b. A synoptic revision of the generic classification of the chelonethid family Cheliferidae Simon (Arachnida). *Canadian Entomologist*, 64: 289–294.

- Chamberlin, J. C. 1947. The Vachoniidae – a new family of false scorpions represented by two new species from caves in Yucatan (Arachnida, Chelonethida, Neobisioidea). *Bulletin of the University of Utah, Biological Series*, 10(4): 1–15.
- Chamberlin, R. V. 1917. New spiders of the family Aviculariidae. *Bulletin of the Museum of Comparative Zoology*, 61: 25–75.
- Chamberlin, R. V. 1922. Two new American arachnids of the order Pedipalpida. *Proceedings of the Biological Society of Washington*, 235: 11–12.
- Chamberlin, R. V. & Ivie, W. 1943. New genera and species of North American linyphiid spiders. *Bulletin of the University of Utah*, 33(10): 1–39.
- Chamberlin, R. V. & Mulaik, S. 1942. On a new family in the Notostigmata. *Proceedings of the Biological Society of Washington*, 55: 125–132.
- Chang A.-c. 1957. On the discovery of the Wenlockian *Eurypterus*-fauna from south China. *Acta Palaeontologica Sinica*, 5: 446–450.
- Chang J.-p. 2004. Some new species of spider and Sacculinidae fossils in Jehol biota. *Global Geology*, 23: 313–320.
- Chapman, F. 1932. Two new Australian fossil king-crabs. *Proceedings of the Royal Society of Victoria, New Series*, 44: 100–102.
- Charbonnier, S., Vannier, J. & Riou, B. 2007. New sea spiders from the Jurassic La Voulte-sur-Rhône Lagerstätte. *Proceedings of the Royal Society B*, 274: 2555–2561.
- Cheng X.-d., Meng Q.-j., Wang X.-r. & Gao C.-l. 2008. [New discovery of Nephilidae in Jehol biota (Araneae, Nephilidae).] *Acta zootaxonomica Sinica*, 33: 330–334. [in Chinese with English summary]
- Cheng X.-d., Shen C.-z. & Gao C.-l. 2009. [A new fossil spider of the Philodromidae from the Yixian Formation of western Liaoning Province, China (Arachnida, Araneae).] *Acta Arachnologica Sinica*, 18: 23–27. [in Chinese with English summary]
- Chernyshev, B. I. 1928. Nouvelles données sur les Xiphosura du bassin Donetz. *Bulletin du Comité Géologique*, 47: 519–531.
- Chernyshev, B. I. 1933. [Arthropoda from the Urals and other regions of the USSR.] *Materials of the Central Scientific and Prospecting Institute Paleontology and Stratigraphy, Magazine*, 1: 15–25. [in Russian with English summary]
- Chernyshev, B. I. 1948. New representative of Merostomata from the Lower Carboniferous. *State of Kiev, Geological Collections*, 2: 119–130.
- Chlupáč, I. 1994. Pterygotid eurypterids (Arthropoda, Chelicerata) in the Silurian and Devonian of Bohemia. *Journal of the Czech Geological Society*, 39: 147–162.
- Chlupáč, I. 1995. Lower Cambrian anthropods from the Paseky Shale (Barrandian area, Czech Republic). *Journal of the Czech Geological Society*, 40: 9–36.

- Chlupáč, I. & Havlíček, V. 1965. *Kodymirus* n. g., a new aglaspid merostome of the Cambrian of Bohemia. *Sborník Geologických Věd. Paleontologie*, 6: 7–20.
- Ciurca Jr., S. J. & Tetlie, O. E. 2007. Pterygotids (Chelicerata; Eurypterida) from the Silurian Vernon Formation of New York. *Journal of Paleontology*, 81: 725–736.
- Clarke, J. M. 1902. Notes on Paleozoic crustaceans. *New York State Museum Report*, 54: 83–110.
- Clarke, J. M. 1907. The *Eurypterus* shales of the Shawangunk Mountains in Eastern New York. *New York State Museum Bulletin* 107: p. 295.
- Clarke, J. N. & Ruedemann, R. 1912. The Eurypterida of New York. – *New York State Museum, Memoir*, 14, 1–439.
- Clarke, J. M. 1919. *Bunaia woodwardi*, a new merostome from the Silurian waterlimes of New York. *Geological Magazine, Decade 6*, 6: 531–532.
- Claypole, E. W. 1890a. Palaeontological notes from Indianapolis (A. A. A. S.) *Pterichthys – Castoroides – Eurysoma* g. n. *American Geologist*, 6: 255–260.
- Claypole, E. W. 1890b. *Carcinosoma newlini*. *American Geologist*, 6: 400.
- Clerck, C. 1757. *Araneae suecici, descriptionibus et figuris oeneis illustrati, ad genera subalterna redacti speciebus ultra LX determinati. Svenska Spindlar, uti sina hufvud-slagter indelte samt...* - Stockholm, 154 pp.
- Cockerell, T. D. A. 1905. Two Carboniferous genera of xiphosurans. *American Geologist*, 36: 330.
- Cockerell, T. D. A. 1907. Some fossil arthropods from Florissant, Colorado. *Bulletin of the American Museum of Natural History*, 23: 605–616.
- Cockerell, T. D. A. 1916. The uropods of *Acanthotelson stimpsoni*. *Journal of the Washington Academy of Science*, 6: 234–236.
- Cockerell, T. D. A. 1917a. Arthropods in Burmese amber. *American Journal of Science, series 4*, 44: 360–368.
- Cockerell, T. D. A. 1917b. Arthropods in Burmese amber. *Psyche*, 24: 40–45.
- Cockerell, T. D. A. 1920. Fossil arthropods in the British Museum. I. *Annals and Magazine of Natural History, series 9*, 5: 273–279.
- Cockerell, T. D. A. 1925. Fossil insects in the United States National Museum. *Proceedings of the U. S. National Museum*, 64: 1–15.
- Coddington, J. 1986. The genera of the spider family Theridiosomatidae. *Smithsonian Contributions to Zoology*, 422: 1–96.
- Coineau, Y. 1974. Un type nouveau d'Acariens Prostigmates libres: les Saxidromoidea, nouvelle super-famille. *Comptes rendus de l'Académie des Sciences, Paris série D*, 278: 1059–1062.
- Coineau, Y. & Magowski, W. Ł. 1994. Caeculidae in amber. *Acarologia*, 35: 243–246.
- Coineau, Y. & Poinar Jr., G. O. 2001. Un Caeculidae de l'ambre de la République Dominicaine. *Acarologia*, 41: 141–144.

- Coineau, Y & Theron, P. 1983. Les Micropsammidae, n. fam. d'Acariens Endeostigmata des sables fin. *Acarologia*, 24: 275–280.
- Cokendolpher, J. C. 1987. A new species of fossil *Pellobunus* from Dominican Republic amber (Arachnida: Opiliones: Phalangodidae). *Caribbean Journal of Science*, 22: 205–211.
- Cokendolpher, J. C. & Poinar Jr., G. O. 1992. Tertiary harvestmen from Dominican Republic amber (Arachnida: Opiliones: Phalangodidae). *Bulletin of the British arachnological Society*, 9: 53–56.
- Cokendolpher, J. C. & Poinar Jr., G. O. 1998. A new fossil harvestman from Dominican Republic amber (Opiliones, Samoidae, *Hummelinckiolus*). *Journal of Arachnology*, 26: 9–13.
- Comstock, J. H. 1940. *The spider book, revised and edited by Willis J. Gertsch*. Ithaca, New York, 729 pp.
- Condé, B. 1996. Les Palpigrades, 1885–1995: acquisitions et lacunes. *Revue suisse de Zoologie*, hors série 1: 87–106.
- Cook, D.R. 1967. Water mites from India. *Memoirs of the American Entomological Institute*, 9: 1–411.
- Cooke, J. A. L. 1965. Spider genus *Dysdera* (Araneae, Dysderidae). *Nature*, 205: 1027–1028.
- Corda, A. J. C. 1835. Ueber den in der Steinkohlenformation bei Cholme gefundenen fossilen Scorpion. *Verhandlungen der Gesellschaft des vaterländischen Museums in Böhmen, Prag*: 36.
- Corda, A. J. C. 1839. Ueber eine fossile Gattung der Afterscorpione. *Verhandlungen der Gesellschaft des vaterländischen Museums in Böhmen, Prag*: 14–18.
- Corronca, J. A. 2003. New genus and species of Selenopidae (Arachnida, Araneae) from Madagascar and neighbouring islands. *African Zoology*, 38: 387–392.
- Crônier, C. & Courville, P. 2005. New xiphosuran merostomata from the Upper Carboniferous of the Graissessac Basin (Massif Central, France). *Comptes Rendus Palevol*, 4: 123–133.
- Crosby, C. R. & Bishop, S. C. 1925. A new genus and two new species of spiders collected by *Bufo quercicus* (Holbrook). *Florida Entomologist* 9: 33–36.
- Cross, E. A. 1965. The generic relationships of the family Pyemotidae (Acarina: Trombidiformes). *Kansas University Science Bulletin*, 45: 29–275.
- Cunliffe, F. 1958. *Pyroglyphus morlani*, a new genus and species of mite forming a new family, Pyroglyphidae, in the Acaridae. *Proceedings of the Entomological Society of Washington*, 60: 85–86.
- Currie, L. D. 1927. On *Cyamocephalus*, a new synziphosuran from the Upper Silurian of Lesmahagow, Lanarkshire. *Geological Magazine*, 64: 153–157.
- Cutler, B. 1970. A fossil crab spider from West-ventral Wyoming (Araneae: Thomisidae). *Entomological News*, 81: 38–40.
- Daber, R. 1990. Arachnidenrest aus dem Westfal D von Zwickau-Oelsnitz. *Zeitschrift für geologische Wissenschaft, Berlin*, 18: 679–682.
- Dabert, J. 1994. Kiwilichidae fam. nov. eine neue Federfamilie (Astigmata, Pterolichoidea). *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, 11: 101–110.

- Daday, E. 1888. A Magyar nemzeti Muzeum àlskorpiónak áttekintése. *Természetrájzi Füzetek*, 11: 111–136, 165–192.
- Dahl, F. 1908. Die Lycosiden oder Wolfsspinnen Deutschlands und ihre Stellung im Haushalt der Natur. Nach statistischen Untersuchungen dargestellt. *Nova Acta Academiae Caesareae Leopoldino-Carolinae*, 88: 175–678.
- Dahl, F. 1912. Arachnoidea. In Korschelt, E. et al. (eds). *Handwörterbuch der Naturwissenschaften*, 1: 485–514.
- Dahl, F. 1913. *Vergleichende Physiologie und Morphologie der Spinnentiere unter besonderer Berücksichtigung der Lebensweise. 1. Die Beziehungen des Körperbaues und der Farben zur Umgebung*. Jena, 1913: 113 pp.
- Dalla Vecchia, F. M. & Selden, P. A. 2013. A Triassic spider from Italy. *Acta Palaeontologica Polonica*, 58: 325–330.
- Dalman, J. W. 1826. Om Insekter inneslutne I Copal, jemte beskrifning på några deribland förekommande nya slägten och arter. *Kungliga Svenska Vetenskapsakademiens Handlingar*, 46: 375–410.
- Dalmas, R. de 1916. Révision du genre *Orchestina* E.S., suivie de la description de nouvelles espèces du genre *Oonops* et d'une étude sur les Dictynidae su genre *Scotolathys*. *Annales de la Société Entomologique de France*, 85: 203–258.
- Dalmas, R. de 1917. Araignées de Nouvelle Zélande. *Annales de la Société Entomologique de France*, 86: 317–430.
- Dana, J. D. 1853. Crustacea, pt. II, Arachnopoda or Pycnogonida. In United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842. Under the command of Chales Wilkes, U.S.N.C. Sherman, Philadelphia, 1382–1391.
- Davies, V. T. 1978. A new family of spiders (Araneae: Teemanaaridae). *Symposium of the Zoological Society of London*, 42: 293–302.
- Davies, V. T. 1980. *Malkara loricata*, a new spider (Araneidae: Malkarinae) from Australia. *Verhandlungen des 8. Internationalen Arachnologen-Kongresses*. Wien, 1980: 377–382.
- Deeleman-Reinhold, C. L. 1995. The Ochyroceratidae of the Indo-Pacific region (Araneae). *Raffles Bulletin of Zoology Supplement*, 2: 1–103.
- Delle, N. 1937. Zemgales lidzenuma, Augszemes un Lietuvas devona nogulumi. *Acta Universitatis Latviensis, Matēmatikas un Dabas Zinātnu Fakultātes Serija* 2(5): 105–384.
- De Geer, C. 1778. *Mémoires pour Servir à l'Histoire des Insectes*, vol. 7. Stockholm.
- De Kay, J. E. 1825. Observations on a fossil crustaceous animal of the order Branchiopoda. *Annals of the New York Lyceum of Natural History*, 1: 375–377.
- Delfinado, M. D. & Baker, E. W. 1974. Varroidae, a new family of mites on honeybees (Mesostigmata: Acarina). *Journal of the Washington Academy of Science*, 64: 4–10.

- De Lima, W. 1890. Note sur un nouval *Eurypterus* du Rothliegendas de Bussaco. *Comunicações da Comissão dos Trabalhos Geológicos da Portugal*, 2: 153–157.
- Desmarest, A.-G. 1822. Les crustacés proprement dits. 66–154. In *Histoire naturelle des crustacés fossiles, sous les rapports zoologiques et géologiques*. F.-G. Levrault, Paris, Strasbourg, **xx** pp.
- Diener, C. 1924. Eurypterida. In Diener, C. (ed.). *Fossilium Catalogus I : Animalia*. W. Junk, Berlin, pp. 1–26.
- Dix, E. & Pringle, J. 1929. On the fossil Xiphosura from the South Wales Coalfield with a note on the myriapod *Euphoberia*. *Summary of Progress, Geological Survey of Great Britain*, 1928: 90–113.
- Dix, E. & Pringle, J. 1930. Some Coal Measures arthropods from the South Wales Coalfield. *Annals and Magazine of Natural History*, 6: 136–144.
- Dohrn, A. 1881. Die Pantopoden des Golfes von Neapel und der angrenzenden Meeresabschnitte. *Monographie der Fauna und Flora des Golfes von Neapel*, 3: 1–252.
- Doleschall, L. 1852. Systematisches Verzeichnis der im Kaiserthum Österreich vorkommenden Spinnen. *Sitzungsberichte der Mathematisch-Naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaft*, 9: 622–651.
- Donnadieu, A. L. 1875. *Recherches pour servir à l'histoire des Tétranyques*. – These. Faculté des Sciences de Lyon: 134 pp. [Thesis also published in a regular journal in 1876.]
- Dresco, E. 1970. Recherches sur la variabilité et la phylogénie chez les Opiliones du genre *Ischyropsalis* C. L. Koch (Fam. Ischyropsalidae), avec la création de la famille nouvelle des Sabaconidae. *Bulletin du Muséum National d'Histoire Naturelle*, 2^e Serie, 41: 1200–1213.
- Dubey, D. P. 1985. A preliminary note on the eurypterid and trilobitid remains from the Upper Vidhyan rocks around Rewa, Madhya Pradesh. *Current Trends in Geology (IV Indian Geological Congress)*, 6: 63–78.
- Dubinin, V. B. 1953. Feather mites (Analgesoidea). II Families Epidermoptidae and Freyanidae. *Fauna SSSR. Paukoobrazynye* 6 (6): 3–411. [In Russian].
- Dubinin, V. B. 1957. On the orientation of the cephalic end of the Devonian pycnogonids of the genus *Palaeoisopus* and their systematic position in the Arthropoda. *Doklady Akademii Nauk SSSR*, 117: 881–884. [In Russian].
- Dufour, L. 1820. Description de cinq Arachnides nouvelles. *Annales générales des sciences physiques*, 5: 198–209.
- Dugès, A. 1834. Recherches sur l'ordre des Acariens et la famille des Trombidés en particular. *Annales des Sciences Naturelles, Zoologie*, série 2, 1: 5–46.
- Dujardin, F. 1851. Sur des acariens à quatre pieds, parasites des végétaux et qui doivent former un genre particulier (*Phytoptus*). In *Observations Zoologiques. Annales des Sciences Naturelles*, série 3, 15: 158–175.
- Dunbar, C. O. 1923. Kansas Permian insects, Part 2. *Paleolimulus*, a new genus of Paleozoic Xiphosura, with notes on other genera. *American Journal of Science*, 5th series, 5: 443–454.

- Dunbar, C. O. 1924. Kansas Permian insects. Part 1. The geologic occurrence and the environment of the insects. *American Journal of Science*, 5th series, 7: 171–209.
- Dunlop, J. A. 1995. Redescription of the Pennsylvanian trigonotarbid arachnid *Lissomartus* Petrunkevitch 1949 from Mazon Creek, Illinois. *Journal of Arachnology*, 23: 118–124.
- Dunlop, J. A. 1996. A trigonotarbid arachnid from the Upper Silurian of Shropshire. *Palaeontology*, 39: 605–614.
- Dunlop, J. A. 1998. A fossil whipscorpion from the Lower Cretaceous of Brazil. *Journal of Arachnology*, 26: 291–295.
- Dunlop, J. A. 1999. A replacement name for the trigonotarbid arachnid *Eotarbus* Dunlop. *Palaeontology*, 42: 191.
- Dunlop, J. A. 2002. Arthropods from the Lower Devonian Severnya Zemlya Formation of October Revolution Island, Russia. *Geodiversitas*, 24: 349–379.
- Dunlop, J. A. 2004. A spiny harvestman (Arachnida: Opiliones) from the Upper Carboniferous of Missouri, USA. In Logunov, D. V. & Penney, D (eds). Proceedings of the 21st European Colloquium of Arachnology, St.-Petersburg, 4–9 August 2003. *Arthropoda Selecta*, Special Issue No. 1: 67–74.
- Dunlop, J. A. 2007. A large parasitengonid mite (Acari, Erythraeoidea) from the Early Cretaceous Crato Formation of Brazil. *Fossil Record*, 10: 91–98.
- Dunlop, J. A. & Anderson, L. I. 2005. A fossil harvestman (Arachnida, Opiliones) from the Mississippian of East Kirkton, Scotland. *Journal of Arachnology*, 33: 482–489.
- Dunlop, J. A. & Bernardi, L. F. de O. 2014. An opilioacarid mite in Cretaceous Burmese amber. *Naturwissenschaften*, 101: 759–763.
- Dunlop, J. A. & Bertrand, M. 2011. Fossil labidostomatid mites (Prostigmata: Labidostommatidae) from Baltic amber. *Acarologia*, 51: 191–198.
- Dunlop, J. A. & Braddy, S. J. 2011. *Cteniza bavincourti* and the nomenclature of arachnid related trace fossils. *The Journal of Arachnology*, 39: 250–257.
- Dunlop, J. A. & Brauckmann, C. 2006. A new trigonotarbid from the Coal Measures of Hagen Vorhalle, Germany. *Fossil Record*, 9: 130–136.
- Dunlop, J. A. & Falkenhagen, R. 2014. Raubmilbe in Aragonit. *Fossilien*, 2014(3): 53–55.
- Dunlop, J. A. & Giribet, G. 2003. The first fossil cyphophthalmid (Arachnida, Opiliones) from Bitterfeld amber, Germany. *The Journal of Arachnology*, 31: 371–378.
- Dunlop, J. A. & Horrocks, C. A. 1996. A new Upper Carboniferous whip scorpion (Arachnida: Uropygi: Thelyphonida) with a revision of the British Carboniferous Uropygi. *Zoologischer Anzeiger*, 234: 293–306.
- Dunlop, J. A. & Horrocks, C. A. 1997. Phalangiotarbid arachnids from the Coal Measures of Lancashire, UK. *Geological Magazine*, 134: 369–381.
- Dunlop, J. A. & Jekel, D. 2009. Nomenclatural notes on fossil spiders. *Bulletin of the British arachnological Society*, 14: 357–360.

- Dunlop, J. A. & Martill, D. M. 2002. The first whipspider (Arachnida: Amblypygi) and three new whipscorpions (Arachnida: Thelyphonida) from the Lower Cretaceous Crato Formation of Brazil. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 92: 325–334.
- Dunlop, J. A. & Mammitzsch, L. 2010. A new genus and species of harvestman from Baltic amber. *Palaeodiversity*, 3: 23–32.
- Dunlop, J. A. & Mitov, P. G. 2009. Fossil harvestmen (Arachnida, Opiliones) from Bitterfeld amber. *ZooKeys*, 16: 347–375.
- Dunlop, J. A. & Mitov, P. G. 2011. The first fossil cyphophthalmid harvestman from Baltic amber. *Arachnologische Mitteilungen*, 40: 47–54.
- Dunlop, J. A. & Penney, D. 2012. *Fossil arachnids*. Siri Scientific Press, Manchester, 192 pp.
- Dunlop, J. A. & Poschmann, M. 1997. On the Emsian (Lower Devonian) arthropods of the Rhenish Schiefergebirge: 1. *Xenarachne*, an enigmatic arachnid from Willwerath, Germany. *Paläontologische Zeitschrift*, 71: 231–236.
- Dunlop, J. A. & Rößler, R. 2003. An enigmatic, solifuge-like fossil arachnid from the Lower Carboniferous of Kamienna Góra (Intra-Sudetic Basin), Poland. *Paläontologische Zeitschrift*, 77: 389–400.
- Dunlop, J. A. & Rößler, R. 2013. The youngest trigonotarbid *Permotarbus schuberti* n. gen., n. sp. from the Permian Petrified Forest of Chemnitz in Germany. *Fossil Record*, 16: 229–243.
- Dunlop, J. A. & Selden, P. A. 2004. A trigonotarbid arachnid from the Lower Devonian of Tredomen, Wales. *Palaeontology*, 47: 1469–1476.
- Dunlop, J. A. & Selden, P. A. 2013. Scorpion fragments from the Silurian of Powys, Wales. *Arachnology*, 16: 27–32.
- Dunlop, J. A., Anderson, L. I. & Braddy, S. J. 1999. A new chasmataspid (Chelicerata: Chasmataspida) from the Lower Devonian of the Midland Valley of Scotland. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 89: 161–165.
- Dunlop, J. A., Anderson, L. I. & Braddy, S. J. 2004. A redescription of *Chasmataspis laurencii* Caster & Brooks (Chelicerata: Chasmataspida) from the Middle Ordovician of Tennessee, USA, with remarks on chasmataspid phylogeny. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 94: 207–205.
- Dunlop, J. A., Bartel, C. & Mitov, P. G. 2012. An enigmatic spiny harvestman from Baltic amber. *Fossil record*, 15: 91–101.
- Dunlop, J. A., Harms, D., Penney, D. 2008. A fossil tarantula (Araneae: Theraphosidae) from Miocene Chiapas amber, Mexico. *Revista Ibérica de Aracnología*, 15: 9–17.
- Dunlop, J. A., Kontschán, J. & Zwanzig, M. 2013. Fossil mesostigmatid mites (Mesostigmata: Gamasina, Microgyniina, Uropodina), associated with longhorn beetles (Coleoptera: Cerambycidae) in Baltic amber. *Naturwissenschaften*, 100: 337–344.

- Dunlop, J. A., Sempf, C. & Wunderlich, J. 2010. A new opilioacarid mite in Baltic amber. In Nentwig, W., Entling, M. & Kropf, C. (eds). *European Arachnology 2008*, pp. 59–70.
- Dunlop, J. A., Wunderlich, J. & Poinar Jr., G. O. 2004. The first fossil opilioacariform mite (Acari: Opilioacariformes) and the first Baltic amber camel spider (Solifugae). *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 94: 261–273.
- Dunlop, J. A., Anderson, L. I., Kerp, H. & Hass, H. 2004. A harvestman (Arachnida: Opiliones) from the Early Devonian Rhynie cherts, Aberdeenshire, Scotland. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 94: 341–354.
- Dunlop, J. A., Fayers, S. F., Hass, H. & Kerp, H. 2006. A new arthropod from the early Devonian Rhynie chert, Aberdeenshire (Scotland), with a remarkable feeding device in the mouthparts. *Paläontologische Zeitschrift*, 80: 296–306.
- Dunlop, J. A., Kutschán, J., Walter, D. E. & Perrichot, V. 2014. An ant-associated mesostigmatid mite in Baltic amber. *Biology Letters*, 10: 20140531.
- Dunlop, J. A., Wang, Y., Selden, P. A. & Krautz, P. 2014. A trigonotarbid arachnid from the Pennsylvanian Astrasado Formation of the Kinney Brick Quarry, New Mexico. *Palaeontological Contributions*, 9: 1–6.
- Dunlop, J. A., Wirth, S., Penney, D., McNeil, A., Bradley, R. S., Withers, P. J. & Preziosi, R. F. 2012. A minute fossil phoretic mite recovered by phase-contrast X-ray computed tomography. *Biology Letters*, 8: 475–460.
- Ebert, T. 1892. *Prestwichia (Euproops) scheeleana*. – *Abhandlung und Jahrbuch Königliche Preußische Geologisches Landesanstalt*, 10: 215–220.
- Edgecombe, G. D. 1998. Early myriapodous arthropods from Australia: *Maldybulakia* from the Devonian of New South Wales. *Records of the Australian Museum*, 50: 293–314.
- Ehlers, G. M. 1935. A new eurypterid from the Upper Devonian of Pennsylvania. *Contributions from the Museum of Palaeontology, University of Michigan*, 4 (18): 291–295.
- Eichwald, E. 1854. Die Grauwackenschichten von Live- und Esthland. *Bulletin de la Société Imperiale des Naturalistes de Moscou*, 27: 1–211.
- Eichwald, E. 1860. *Lethaea Rossica*. Vol. 1. Seconde section de l'ancienne Période. Librairie et Imprimerie de E. Schweizerbart, Stuttgart, 1657 pp.
- Eldredge, N. 1974. Revision of the suborder Synziphosurina (Chelicerata, Merostomata), with remarks on merostome phylogeny. *American Museum Novitates*, 2543: 1–41.
- Elias, M. K. 1936. Character and significance of the late Paleozoic flora of Garnett, Kansas. *Journal of Geology*, 44: 9–23.
- Eller, E. R. 1938a. A review of the xiphosuran genus *Belinurus* with the description of a new species, *B. allegayensis*. *Annals of the Carnegie Museum*, 27: 129–150.

- Eller, E. R. 1938b. A new xiphosuran, *Euproops morani*, from the Upper Devonian of Pennsylvania. *Annals of the Carnegie Museum*, 27: 152–153.
- Eller, E. R. 1940. *Belinurus carteri* a new xiphosuran from the Upper Devonian of Pennsylvania. *Annals of the Carnegie Museum*, 28: 133–136.
- Ellingsen, E. 1906. Report on the pseudoscorpions of the Guinea Coast (Africa) collected by Leonardo Fae. *Annali del Museo Civico de Storia Naturale di Genova*, (3)2: 243–265.
- Ellingsen, E. 1909. On some North American pseudoscorpions collected by Dr. F. Silvestri. *Bollettino del Laboratorio di Zoologia Generale e Agraria della R. Scuola sup. d'Agricoltura, Portici*, 3: 216–221.
- Elzinga, R. J. 1993. Larvamimidae, a new family of mites (Acari: Dermanysoidea) associated with army ants. *Acarologia*, 34: 95–103.
- Emerton, J. H. 1875 Notes on spiders from Caves in Kentucky, Virginia and Indiana. *American Naturalist*, 9: 278–281.
- Emerton, J. H. 1882. New England spiders of the family Theridiidae. *Transactions of the Connecticut Academy of Arts and Sciences*, 6: 1–86.
- Engel, M. S. & Grimaldi, D. A. 2014. Whipspiders (Arachnida: Amblypygi) in amber from the Early Eocene and mid-Cretaceous, including maternal care. *Novitates Paleoentomologicae*, 9: 1–17.
- Eskov, K. Y. 1984. A new fossil spider family from the Jurassic of Transbaikalia from (Araneae: Chelicerata). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1984: 645–653.
- Eskov, K. Y. 1987. A new archaeid spider (Chelicerata: Araneae) from the Jurassic of Kazakhstan, with notes on the so-called “Gondwanan” ranges of recent taxa. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 175: 81–106.
- Eskov, K. Y. 1992. Archaeid spiders from Eocene Baltic amber (Chelicerata: Araneida: Arachaeidae) with remarks on the so-called “Gondwanan” ranges of Recent taxa. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 185: 311–328.
- Eskov, K. Y. & Marusik, Y. M. 1992. [Fossil spiders of the family Nesticidae.] *Palaeontologicheskii Zhurnal*, 2: 87–95. [In Russian]
- Eskov, K. Y. & Selden, P. A. 2005. First record of spiders from the Permian period (Araneae: Mesothelae). *Bulletin of the British arachnological Society*, 13: 111–116.
- Eskov, K. Y. & Wunderlich, J. 1995 (for 1994). On the spiders of the Taimyr ambers, Siberia, with the description of a new family and with general notes on the spiders from the Cretaceous resins. *Beiträge zur Araneologie*, 4: 95–107.
- Eskov, K. Y. & Zonstein, S. 1990. First Mesozoic mygalomorph spiders from the Lower Cretaceous of Siberia and Mongolia, with notes on the system and evolution of the infraorder Mygalomorphae (Chelicerata: Araneae). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 178: 325–368.

- Eskov, K. Y. & Zonstein, S. L. 2000. The first Ctenizoid Mygalomorph Spiders from Eocene Baltic amber (Araneida: Mygalomorphae: Ctenizidae). *Paleontological Journal*, 34: S268–S274. [Translated into English; original in Russian]
- Etheridge Jr., R. 1877. On the remains of a large crustacean, probably indicative of a new species of *Eurypterus*, or allied genus (*Eurypterus?* *Stevensonii*), from the Lower Carboniferous Series (Cementstone Group) of Berwickshire. *Quarterly Journal of the Geological Society*, 33: 223–228.
- Ewing, H. E. 1922. Studies on the taxonomy and biology of the tarsnemid mites, together with a note on the transformation of *Acarapis (Tarsonemus) woodi* Rennie (Acarina). *Canadian Entomologist*, 54: 104–113.
- Ewing, H. E. 1929. A synopsis of the American arachnids of the primitive order Ricinulei. *Annals of the Entomological Society of America*, 22: 583–600.
- Ewing, H. E. 1930. A fossil arachnid from the Lower Carboniferous shales (Pococno formation) of Virginia. *Annals of the Entomological Society of America*, 23: 641–643.
- Fage, L. 1912. Etudes sur les araignées cavernicoles. I. Revision des Ochyroceratidae (n. fam.). In *Biospelogica*, XXV. *Archives de Zoologie expérimentale et générale*, 10: 97–162.
- Fage, L. 1913. Etudes sur les Araignées cavernicoles. II. Revision des Leptonetidae. In *Biospelogica*, XXIX. *Archives de Zoologie expérimentale et générale*, 10: 479–576.
- Fain, A. 1956. Une nouvelle famille d'acariens endoparasites des chauves-souris: Gastronyssidae fam. nov. *Annales de la Société Belge de Médecine Tropicale*, 36: 87–98.
- Fain, A. 1961. Une nouvelle famille d'acariens, parasites de serpents du genre *Mehelya* au Congo: Omentolaelaptidae Fam. nov. (Mesostigmata). *Revue de Zoologie et de Botanique Africaine*, 64: 283–296.
- Fain, A. 1967a. Nouveaux hypopes vivant dans les follicules pileux de Rongeurs américains. *Revue de Zoologie et de Botanique Africaine*, 76: 157–162.
- Fain, A. 1967b. Un acarien remarquable récolté sur un Tarsier (Heteroptidae f.n. : Sarcoptiformes). – *Zoologischer Anzeiger*, 178: 90–94.
- Fain, A. 1968. Deux nouveaux Acariens Cavernicoles du Gabon (Sarcoptiformes). *Revue Biologia Gabonica*, 4: 195–205.
- Fain, A. 1974. Acariens récoltés par le Dr. J. Travé aux îles subantarctiques. I. Familles Saprolyphidae et Hyadesidae (Astigmates). *Acarologia*, 16: 684–708.
- Fayers, S. R., Dunlop, J. A. & Trewin, N. H. 2005. A new early Devonian trigonotarbid arachnid from the Windyfield chert, Rhynie, Scotland. *Journal of Systematic Palaeontology*, 2: 269–284.
- Feider, Z. 1955. Arachnida, Acarina Trombidioidea. *Fauna RPR*, 5: 1–187.
- Feider, Z. & Vasiliu, N. 1969. Révision critique de la famille des Nicoletiellidae. In Proc. 2nd International Congress of Acarology, Sutton Bonington, England 1967. Acad. Kiado, Budapest: pp. 202–207.

- Feldmann, R. M., Schweitzer, C. E., Dattilo, B. & Farlow, J. O. 2011. Remarkable preservation of a new genus and species of limuline horseshoe crab from the Cretaceous of Texas, USA. *Palaeontology*, 54: 1337–1346.
- Feldmann, R. M., Vega, F. J., Applegate, S. P., & Bishop, G. A. 1998. Early Cretaceous arthropods from the Tlayua Formation at Tepexi de Rodriguez, Puebla, México. *Journal of Paleontology*, 72: 79–90.
- Fet, V. & Bechly, G. 2001. Case 3120a. Liochelidae, fam. nov. (Scorpiones): proposed introduction as a substitute name for Ischnuridae Simon, 1879, as an alternative to the suggested emendment of Ischnurinae Fraser, 1957 (Insecta, Odonata) to Ischnurinae in order to remove homonymy. *Bulletin of Zoological Nomenclature*, 58: 280–281.
- Fischer de Waldheim, G. 1839. Notes sur un crustacé fossile du genre *Eurypterus* de Podolie. *Bulletin de la Societe Imperiale des Naturalistes de Moscou*, 11: 125–128.
- Flower, R. H. 1945. A new Deepkill eurypterid. *American Midland Naturalist*, 34: 717–719.
- Flower, R. [H.] 1969. Merostomes from a Cotter horizon of the El Paso Group. *New Mexico Bureau of Mines and Mineral Resources Memoir*, 22: 35–44.
- Fraipont, J. 1889. Euryptérides nouveaux du Dévonien Supérieur de Belgique (Psammites du Condroz). *Annales de la Société Géologique de Belgique*, 17: 53–62.
- Forsslund, K.-H. 1941. Schwedische Arten der Gattung *Suctobelba* Paoli (Acari, Orbatei). *Zoologiska bidrag från Uppsala*, 20: 381–396.
- Forsslund, K.-H. 1947. Über die Gattung *Autogneta* Hull (Acari, Oribatei). *Zoologiska bidrag från Uppsala*, 25: 111–117.
- Forsslund, K.-H. 1956. Schwedische Oribatei (Acari). III. *Entomologisk Tidskrift*, 77: 210–218.
- Forster, R. R. 1948. A new sub-family and species of New Zealand Opiliones. *Records of the Auckland Institute and Museum*, 3: 313–318.
- Forster, R. R. 1954. The New Zealand harvestmen (sub-order Laniatores). *Canterbury Museum Bulletin*, 2: 1–329.
- Forster, R. R. 1955. A new family of spiders of the sub-order Hypochilomorphae. *Pacific Science*, 9: 277–285.
- Forster, R. R. & Forster, L. 1999. *Spiders of New Zealand and their worldwide kin*. University of Otago Press, Dunedin, vi + 270 pp.
- Forster, R. R. & Platnick, N. I. 1984. A review of archaeid spiders and their relatives, with notes on the superfamily Palpimanoidea (Arachnida: Araneae). *Bulletin of the American Museum of Natural History*, 178: 1–106.
- Forster, R. R. & Wilton, C. L. 1973. The spiders of New Zealand. Part IV. *Otago Museum Bulletin*, 4: 1–309.
- Frič, A. 1873. Fauna der Steinkohlenformation Böhmens. *Archiv für Naturwissenschaftliche Landesdurchforschung von Böhmen*, 2(2): 1–16.
- Frič, A. 1899a. On *Prolimulus woodwardi*. *Geological Magazine*, 6: 57–58.

- Frič, A. 1899b. *Fauna der Gaskohle und der Kalksteine der Permformation Böhmens. Vol. IV*: pp. 33–64.
- Frič, A. 1901. *Fauna der Gaskohle und der Kalksteine der Permformation Böhmens. Vol. IV, part 2. Myriopoda pars II. Arachnoidea*, pp. 56–63, pls 153, 154, Prague.
- Frič, A. 1904. *Palaeozoische Arachniden*. A Frič, Prague, 85 pp.
- Fritsch, K. von 1906. Beitrag zur Kenntnis der Tierwelt der deutschen Trias. *Abhandlungen der naturforschender Gesellschaft Halle*, 24: 220–285.
- Fry, W. G. 1978. A classification within the pycnogonids. *Zoological Journal of the Linnean Society*, 63: 35–58.
- García-Villafuerte, M. Á. 2006a. A new fossil *Episinus* (Araneae, Theridiidae) from Tertiary Chiapas amber, Mexico. *Revista Ibérica de Aracnología*, 13: 120–125.
- García-Villafuerte, M. Á. 2006b. Selenopidae y Thomisidae (Arachnida: Araneae) en ámbar de Chiapas, México. *Boletín Sociedad Entomológica Aragonesa*, 38: 209–212.
- García-Villafuerte, M. Á. 2008. Primer registro fósil del género *Hemirraghus* (Araneae, Theraphosidae) en ámbar del Terciario, Chiapas, México. *Revista Ibérica de Aracnología*, 16: 43–47.
- Garwood, R. J., Dunlop, J. A., Giribet, G. & Sutton, M. D. 2011. Anatomically modern Carboniferous harvestmen demonstrate early cladogenesis and stasis in Opiliones. *Nature Communications*, 2:444: 1–7.
- Garwood, R. J., Sharma, P. P., Dunlop, J. A., Giribet, G. 2014. A Paleozoic stem group to mite harvestmen revealed through integration of phylogenetics and development. *Current Biology*, 24: 1–7.
- Gaud, J. & Atyeo, W. T. 1975. Gabuciniidae, famille nouvelle de Sarcoptiformes plumicoles. *Acarologia*, 16: 522–561.
- Gaud, J. & Atyeo, W. T. 1976. Ascouracarinae, n. sub-fam. des Syringobiidae, Sarcoptiformes plumicoles. *Acarologia*, 18: 143–162.
- Gaud, J. & Atyeo, W. T. 1977. A new name for *Ovacarus* and Ovacaridae (Acarina: Analgoidea). *Acarologia*, 18: 568–569.
- Gaud, J. & Atyeo, W. T. 1978. Nouvelles superfamilles pour les Acariens astigmates parasites d'oiseaux. *Acarologia*, 19: 678–685.
- Gaud, J., Atyeo, W.T. & Berla, H.F. 1972. Acariens Sarcoptiformes plumicoles parasites des Tinamous. *Acarologia*, 14: 393–453.
- Gaud, J., Atyeo, W. T. & Klompen, J. S. H. 1989. Oconnoriidae, a new family of feather mites (Acarina, Pterolichoidea). *Journal of Entomological Science*, 24: 417–421.
- Geinitz, H. B. 1882. *Kreischeria wiedei*, ein Pseudoskorpion aus der Steinkohlenformation von Zwickau. *Zeitschrift der Deutschen geologischen Gesellschaft*, 34: 238–242.
- Gerecke, R., Smith, I. M. & Cook, D. R. 1999. Three new species of *Apheviderulix* gen. nov. and proposal of Apheviderulicidae fam. nov. (Acari: Hydrachnidia: Eylaoidea). *Hydrobiologia*, 397: 133–147.

- Gerson, U. & Walter, D. E. 1998. Transfer of *Mecognatha* Wood from Stigmeidae to Mecognathidae, fam. nov., a new synonymy, and a key to families of Raphignathoidea (Acari: Prostigmata). *Systematic and Applied Acarology*, 3: 145–147.
- Gerstaecker, C. E. A. 1863. Pantopoda. 248–350. In Carus, J. V. & Gerstaecker, C. E. A. (eds). *Handbuch der Zoologie*, 2. W. Engelmann, Leipzig, 642 pp.
- Gertsch, W. J. 1941. Report on some arachnids from Barro Colorado Island, Canal Zone. *American Museum Novitates*, 1146: 1–14.
- Gertsch, W. J. & Davis, L. I. 1946. Report on a collection of spiders from Mexico. V. *American Museum Novitates*, 1313: 1–11.
- Gervais, P. M. 1844. Remarques sur la famille des Scorpiones et descriptions des plusiers espèces nouvelles de la collection du Muséum. *Archives du Muséum d'Histoire Naturelle, Paris*, 4: 201–240.
- Gess, R. W. 2013. The earliest record of terrestrial animals in Gondwana: A scorpion from the Famennian (Late Devonian) Witpoort Formation of South Africa. *African Invertebrates*, 54: 373–379.
- Giebel, C. G. 1856. *Die Insekten und Spinnen der Vorwelt mit steter Berücksichtigung der lebenden Insekten und Spinnen; monographisch dargestellt*. Leipzig, 511 pp.
- Gill, E. L. 1909. An arachnid from the Coal Measures of the Tyne Valley. *Transactions of the Natural History Society of Northumberland, Durham and Newcastle-upon-Tyne, new series*, 3(2): 3–16.
- Gill, E. L. 1911. A Carboniferous arachnid from Lancashire. *Geological Magazine*, 48: 395–398.
- Gill, E. L. 1924. Fossil arthropods from the Tyne Coalfield. *Geological Magazine*, 61: 445–471.
- Giribet, G. & Dunlop, J. A. 2005. First identifiable Mesozoic harvestman (Opiliones: Dyspnoi) from Cretaceous Burmese amber. *Proceedings of the Royal Society B*, 272: 1007–1013.
- Giribet, G., Tourhino, A. L., Shih, C.-k. & Ren, D. 2012. An exquisitely preserved harvestman (Arthropoda, Arachnida, Opiliones) from the Middle Jurassic of China. *Organisms, Diversity & Evolution*, 12: 51–56.
- Giribet, G., Sharma, P. P., Benavides, L. R., Boyer, S. L., Clouse, R. M., De Bivort, B. L., Dimitrov, D., Kawauchi, G. Y., Murienne, J., Schwendinger, P. J. 2012. Evolutionary and biogeographical history of an ancient and global group of arachnids (Arachnida: Opiliones: Cyphophthalmi) with a new taxonomic arrangement. *Biological Journal of the Linnean Society*, 105: 92–130.
- Gistel, J. 1848. *Naturgeschichte des Thierreichs für höhere Schulen*. Stuttgart, pp. 155–156.
- Gjelstrup, P. & Solhøy, T. 1994. Oribatid mites (Acari). In The Zoology of Iceland. *Steenstrupia*, (3) 57: 1–78.
- Glushenko, N. V. & Ivanov, V. K. 1961. [Paleolimulus from the Lower Permian of the Donetz Basin.] *Paleontologiceskij Žurnal*, 1861: 128–130. [in Russian]
- Goldenberg, F. 1873. *Fauna Sarapontana Fossilis. Die fossilen Thiere aus der Steinkohlenformation von Saarbrücken. Erstes Heft*. Chr. Möllinger Verlag, Saarbrücken, 26 pp.
- Goodnight, J. C. & Goodnight, M. L. 1942. Phalangids from Central America and the West Indies. *American Museum Novitates*, 1184: 1–23.

- Gonzalez, R. H. 1978. A new species of xenocaligonellid mite from the Galapagos Islands (Acari). *Proceedings of the Entomological Society of Washington*, 80: 191–196.
- González-Sponga, M. A. 1997. Arácnidos de Venezuela. Una nueva familia, dos nuevos géneros y dos nuevas especies de Opiliones Laniatores. *Acta Biologica Venezolana*, 17: 51–58.
- Gourret, P. 1887. Recherches sur les Arachnides tertiaires d'Aix en Provence. *Recueil Zoologique Suisse*, 4: 431–496.
- Grabau, A. W. 1920. A new species of *Eurypterus* from the Permian of China. *Bulletin of the Geological Survey of China*, 2: 61–68.
- Grandjean, F. 1931. Observations sur les Oribates (1^{re} Série). *Bulletin du Muséum National d'Histoire Naturelle*, 3: 131–144.
- Grandjean, F. 1932a. Observations sur les Oribates (3^e série). *Bulletin du Muséum National d'Histoire Naturelle*, 4: 292–306.
- Grandjean, F. 1932b. Au sujet des *Palaearcariformes* Trägårdh. *Bulletin du Muséum National d'Histoire Naturelle*, 4: 411–426.
- Grandjean, F. 1933. Études sur les Développement des Oribates. *Bulletin de la Société zoologique de France*, 58: 30–61.
- Grandjean, F. 1934. La notation des poils gastronotiques et des poils dorsaux du propodosoma chez les Oribates (Acariens). *Bulletin de la Société zoologique de France*, 59: 12–44.
- Grandjean, F. 1936a. Les Microzetidae n. fam. (Oribates). *Bulletin de la Société zoologique de France*, 61: 60–93.
- Grandjean, F. 1936b. Les Oribates de Jean Frédéric Hermann et de son père [Arachn. Acar.]. *Annales Société Entomologique de France*, 105: 27–110.
- Grandjean, F. 1936c. Observations sur les Oribates (10^e Série). *Bulletin du Muséum National d'Histoire Naturelle*, 8: 246–253.
- Grandjean, F. 1937. Le Genre *Pachygnathus* Dugès (*Alycus* Koch) (Acariens). Cinquième et dernière partie. *Bulletin du Muséum National d'Histoire Naturelle*, 9: 262–269.
- Grandjean, F. 1939. Quelque genres d'Acariens appartenant au groupe des Endeostigmata. *Annales des Sciences Naturelles – Zoologie et Biologie Animale*, Série 11, 2: 1–122.
- Grandjean, F. 1947a. Études sur les Smarisidae et quelques autres Èrythroïdes (Acariens). *Archives de Zoologie Expérimental et Générale*, 85: 1–126.
- Grandjean, F. 1947b. Les Enarthronota (Acariens). Première série. *Annales des Sciences Naturelles – Zoologie et Biologie Animale*, 8: 213–248.
- Grandjean, F. 1948. Les Enarthronota (Acariens). (2^e série). *Annales des Sciences Naturelles – Zoologie et Biologie Animale*, 10: 29–58.

- Grandjean, F. 1950. Les Enarthronota (Acariens). (3^e série). *Annales des Sciences Naturelles – Zoologie et Biologie Animale*, 12: 85–107.
- Grandjean, F. 1951. Observations sur les Oribates (22^e Série). *Bulletin du Muséum National d'Histoire Naturelle*, 23: 91–98.
- Grandjean, F. 1953. Observations sur les Oribates (25^e Série). *Bulletin du Muséum National d'Histoire Naturelle*, 25: 155–162.
- Grandjean, F. 1954a. Observations sur les Oribates (28^e série). *Bulletin du Muséum National d'Histoire Naturelle*, 26: 204–211.
- Grandjean, F. 1954b. Essai de classification des Oribates (Acariens). *Bulletin de la Société zoologique de France*, 78: 421–446.
- Grandjean, F. 1954c. Etude sur les Palaeacaroïdes (Acariens, Oribates). *Mémoires du Muséum National d'Histoire Naturelle*, 7: 179–274.
- Grandjean, F. 1956a. Sur deux espèces nouvelles d'oribates (Acariens) apparentées à *Oripoda elongata* Banks 1904. *Archives de Zoologie Expérimentale et Générale*, 93: 185–218.
- Grandjean, F. 1956b. Galumnidae sans carènes lamellaires (Acariens, Oribates), 1^{re} série. *Bulletin de la Société zoologique de France*, 81: 134–150.
- Grandjean, F. 1958a. *Perlohmannia dissimilis* (Hewitt) (Acarien, Oribate). *Mémoires du Muséum National d'Histoire Naturelle*, 16: 57–120.
- Grandjean, F. 1958b. *Charassobates cavernosus* Grandj. 1929 (Acarien, Oribate). *Mémoires du Muséum National d'Histoire Naturelle*, 16: 121–140.
- Grandjean, F. 1959. *Polypterozetes cherubin* Berl. 1916 (Oribate). *Acarologia*, 1: 147–180.
- Grandjean, F. 1960a. Les Mochlozetidae n. fam. (Oribates). *Acarologia*, 2: 101–148.
- Grandjean, F. 1960b. Les Autognetidae n. fam. (Oribates). *Acarologia*, 2: 575–609.
- Grandjean, F. 1961a. Les Plasmobatidae n. fam. (Oribates). *Acarologia*, 3: 96–129.
- Grandjean, F. 1961b. Les Amerobelbidae (Oribates). (1^{re} partie). *Acarologia*, 3: 303–343.
- Grandjean, F. 1963. Les Autognetidae (Oribates). Deuxième partie. *Acarologia*, 4: 632–689.
- Grandjean, F. 1965a. Nouvelles observations sur les Oribates (4^e série). *Acarologia*, 7: 91–112.
- Grandjean, F. 1965b. Oribates mexicains (2^e série). *Stelechobates megalotrichus* n.g., n.sp. *Acarologia*, 7: 532–563.
- Grandjean, F. 1965c. Complément à mon travail de 1953 sur la classification des Oribates. *Acarologia*, 7: 713–734.
- Grandjean, F. 1966. Les Staurobatidae n. fam. (Oribates). *Acarologia*, 8: 696–727.
- Grandjean, F. 1967. Nouvelles observations sur les Oribates (5^e série). *Acarologia*, 9: 242–272.
- Grandjean, F. 1969. Considérations sur le classement des Oribates. Leur division en 6 groupes majeurs. *Acarologia*, 11: 127–153.

- Grandjean, F. 1970. Nouvelles observations sur les Oribates (8^e série). *Acarologia*, 12: 849–876.
- Grassi, B. & Calandruccio, S. 1885. Intorno ad un nuovo Aracnide Artrogastro (*Koenenia mirabilis* [sic]) che crediamo rappresentante d'un nuovo ordine (Microteliphonida). *Naturalista Siciliano*, 4: 127–133, 162–168.
- Griffiths, D. A. 1977. A new family of astigmatid mites from the Iles Crozet, sub-Antarctica, introducing a new concept relating to ontogenetic development of idiosomal setae. *Journal of Zoology, London*, 182: 291–308.
- Griswold, C., Audisio, T. & Ledford, J. 2012. An extraordinary new family of spiders from caves in the Pacific Northwest (Araneae, Trogloraptoridae, new family). *ZooKeys*, 215: 77–102.
- Gromov, A.V. 1998. [A new family, genus and species of scorpions (Arachnida, Scorpiones) from southern Central Asia.] – *Zoologichesky Zhurnal*, 77: 1003–1009. [In Russian.]
- Grote, A. R. & Pitt, W. H. 1875. I. Description of a new Crustacean from the Water Lime Group at Buffalo. *Bulletin of the Buffalo Society of Natural Sciences*, 3: 1–2.
- Gross, W. 1933. Die unterdevonischen Fische und Gigantostraken von Overath. *Abhandlungen der Preußischen Geologischen Landesanstalt (N. F.)*, 145: 41–77.
- Guérin-Méneville, F. E. 1839. Gastéracanthes sculptée et de Feisthamel, nouvelles espèces d'aranéides. *Revue zoologique*. 1839: 109–111.
- Guthörl, P. 1934. Die Arthropoden aus dem Carbon und Perms des Saar-Nahe-Pfalz-Gebietes. *Abhandlungen der Preußischen Geologischen Landesanstalt (N.F.)*, 164: 1–219.
- Guthörl, P. 1938. *Eophrynus waechteri* n. sp. (Arac., Anthracom.) aus der Tiefbohrung Stangenmühle, Saar-Karbon. *Senckenbergiana*, 20: 465–470.
- Guthörl, P. 1964. Zur Arthropoden-Fauna des Karbons und Perms. 20. Neue Arachniden-Funde (Anthracom.) aus dem Westfal A des Aachener Karbons. *Paläontologische Zeitschrift*, 38: 98–103.
- Guthörl, P. 1965. Zur Arthropoden-Fauna des Karbons und Perms. 19. Weiteres über die Arachniden aus dem Westfal und Stefan des saar-lothringischen und pfälzischen Karbons. *Annales Universitatis Saraviensis*, 4: 10–24.
- Haase, E. 1890. Beitrag zur Kenntniss der fossilen Arachniden. *Zeitschrift der Deutsche geologische Gesellschaft*, 1890: 629–657.
- Haeckel, E. 1866. *Generale Morphologie der Organismen. Band 2*. Berlin, 574 pp.
- Hadži, J. 1931. Skorpionreste aus dem tertiären Sprudelsinter von Böttingen (Schwäbische Alb). *Paläontologische Zeitschrift*, 13: 134–148.
- Hadži, J. 1935. Ein eigentümlicher neuer Höhlen-Opilionid aus Nord-Amerika, *Cladonychium corii* g.n. sp. n. *Biologia Generalis*, 11: 49–72.
- Halbert, J. N. 1915. Clare Island Survey, 39. Acarinida. Section II. Terrestrial and marine Acarina. *Proceedings of the Royal Irish Academy*, 31: 45–136.

- Hall, J. 1859. *Natural History of New York: Palaeontology, III.* New York State Museum, 532 pp.
- Hall, C. E. 1877. Contributions to Palaeontology from the Museum of the Second Geological Survey. *Proceedings of the American Philosophical Society*, 16: 621??.
- Hall, J. 1884a. Description of a New Species of *Stylonurus* from the Catskill Group. *New York State Museum (36th Annual Report)*: 76–77.
- Hall, J. 1884b. Note on Eurypteridae of the Devonian and Carboniferous Formations of Pennsylvania, with a supplementary note on the *Stylonurus excelsior*. *Proceedings of the American Association for the Advancement of Science*, 33: 420–422.
- Hall, J. 1884c. Eurypteridae from the Lower Productive Coal Measures in Beaver County, and the Lower Carboniferous Pithole Shale in Venango County. *2nd Geological Survey of Pennsylvania. Report of Progress PPP*: 23–39.
- Halliday, R. B. 2006. New taxa of mites associated with Australian termites (Acaria: Mesostigmata). *International Journal of Acarology*, 32: 27–38.
- Hall, J. & Clarke, J. M. 1888. *Paleontology of New York*. New York, 236 pp.
- Hall, J. & Clarke, J. M. 1888. Trilobites and other Crustacea of the Oriskany, Upper Helderberg, Hamilton, Portage, Chemung, and Catskill Groups. *Geological Survey of the State of New York, Palaeontology*, 7: XXX.
- Hammer, M. 1966. Investigations on the Oribatid Fauna of New Zealand, Part 1. *Biologiske Skrifter udgivet af Det Kongelige Danske Videnskabernes Selskab*, 15(2): 1–108.
- Hammer, M. 1967. Some oribatids from Kodiak Island near Alaska. *Acta Arctica*, 14: 5–25.
- Hammer, M. 1973. Oribatids from Tongatapu and Eua, the Tonga Islands, and from Upolu, Western Samoa. *Biologiske Skrifter udgivet af Det Kongelige Danske Videnskabernes Selskab*, 20(3): 1–70.
- Hansen, H. J. 1894. Arthogastra Danica: en monographisk fremstilling af de i Danmark levende Meiere og Mosskorptioner med bidrag til sidstnævnte underordens systematic. *Naturhistorisk Tidsskrift*, (3) 14: 491–554.
- Hansen, H. J. & Sørensen, W. 1904. *On two orders of Archanaida*. Cambridge University Press, Cambridge, xi + 178 pp.
- Harger, O. 1874. Notice of a new spider from the Coal Measures of Illinois. *American Journal of Science*, 7: 219–223.
- Harlan, R. 1834. Critical notices of various organic remains hitherto discovered in North America. *Transactions of the Geological Society of Pennsylvania*, 1: 46–112.
- Harvey, M. S. 1990. Pezidae, a new freshwater mite family from Australia (Acarina: Halacaroidea). *Invertebrate Taxonomy*, 3: 771–781.
- Harvey, M. S. 1991. *Catalogue of the Pseudoscorpionida*. Manchester University Press, Manchester, vi + 726.

- Harvey, M. S. 1992. The phylogeny and classification of the Pseudoscorpionida (Chelicerata: Arachnida). *Invertebrate Taxonomy*, 6: 1373–1435.
- Harvey, M. S. 2002. Nomenclatural notes on Solifugae, Amblypygi, Uropygi and Araneae (Arachnida). *Records of the Western Australian Museum*, 20: 449–459.
- Harvey, M. S. 2003. *Catalogue of the smaller arachnid orders of the world*. CSIRO Publishing, Collingwood VC, xi + 385 pp.
- Harvey, M. A. & Selden, P. A. 1995. *Nyranytarbus*, replacement name for *Hemiphrynx* Frič, 1901 (Trigonotarbida: Eophrynidae). *Bulletin of the British arachnological Society*, 10: 74.
- Haupt, H. 1956. Beitrag zu Kenntnis der eozanen Arthropodenfauna des Geiselthals. *Nova Acta Leopoldina n.s.*, 128: 1–90.
- Haupt, H. 1957. Eine spinnenartige Arthropode aus dem Rotliegenden: *Rhabdotarachnoides simoni* n. gen. n. sp. *Hallesches Jahrbuch für Mitteldeutsche Erdgeschichte*, 2(4): 246–247.
- Haupt, J. 1983. Vergleichende Morphologie der Genitalorgane und Phylogenie der liphistiomorphen Webspinnen (Araneae: Mesothelae). I. Revision der bisher bekannten Arten. *Zeitschrift für zoologische Systematik und Evolutionsforschung*, 21: 275–293.
- Hauschke, N. & Wilde, V. 1987. *Paleolimulus fuchsbergensis* n. sp. (Xiphosura, Merostomata) aus der oberen Trias von Nordwestdeutschland, mit einer Übersicht zur Systematik und Verbreitung rezenter Limuliden. *Paläontologische Zeitschrift*, 61: 87–108.
- Hauschke, N. & Wilde, V. 1989. Ein Limulide aus dem Zechstein (Oberes Perm) der Korbacher Bucht (Hessen, Bundesrepublik Deutschland). *Geologisches Jahrbuch Hessen*, 117: 17–21.
- Hauschke, N. & Wilde, V. 2000. Limulidenreste aus dem Unteren Buntsandstein (Benberg-Formation) von Beesenlaublingen (Sachsen-Anhalt). *Hallesches Jahrbüch für Geowissenschaften, Reihe B*, 22: 87–90.
- Hauschke, N. & Wilde, V. 2004. Palaeogene limulids (Xiphosura) from Saxony-Anhalt (Germany) – systematics and palaeobiogeography. *Hallesches Jahrbuch für Geowissenschaften, Reihe B*, 18: 161–168.
- Hauschke, N. & Wilde, V. 2008. Limuliden aus dem Oberen Buntsandstein von Süddeutschland. *Hallesches Jahrbuch für Geowissenschaften*, 30: 21–26.
- Hauschke, N., Osterink, H. W. & Wilde, V. 2009. Erster Nachweis eines Limuliden (Xiphosura, Limulacea) im Muschelkalk von Winterswijk (Niederlande). *Der Aufschluss*, 60: 13–23.
- Hauschke, N., Wilde, V. & Brauckmann, C. 2004. Triassic limulids from Madagascar – missing links in the distribution of Mesozoic Limulacea. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 2004(2): 87–94.
- Hauschke, N., Wilde, V. & Pietrzeniuk, E. 1992. Ein Limulide aus dem Muschelkalk (mittlere Trias) von Rüdersdorf bei Berlin. *Zeitschrift für geologische Wissenschaft*, 20: 461–466.
- Hedgpeth, J. W. 1978. A reappraisal of the Palaeopantopoda with description of a species from the Jurassic. *Zoological Journal of the Linnean Society*, 63: 23–34.

- Heer, O. 1865. *Die Urwelt der Schweiz*. Friedrich Schultheß, Zürich, xxix + 622 pp.
- Heetoff, M., Helfen, L. & Norton, R. A. 2009. Description of *Neoliodes dominicus* n. sp. (Acari, Oribatida) from Dominican Amber, aided by synchrotron X-ray microtomography. *Journal of Paleontology*, 83: 153–159.
- Heide, S. van der 1951. Les arthropodes du terrain houiller du Limbourg meridionale (excepte les scorpions et les insectes). *Mededeelingen van de Geologische Stichting Serie C-IV-3 5*: 1–84.
- Heineken C. & Lowe R. T. 1832. Descriptions of two species of Araneidae, natives of Madeira. *Zool. Journ.*, 5: 320–323.
- Henderickx, H. 2005. A new *Geogarypus* from Baltic amber (Pseudoscorpiones: Geogarypidae). *Phegea*, 33: 87–92.
- Henderickx, H. & Boone, M. 2014. The first fossil *Feaella* Ellingsen, 1906, representing an unexpected pseudoscorpion family in Baltic amber (pseudoscorpiones, Feaellidae). *Entomo-Info*, 25: 5–11.
- Henderickx, H., Tafforeau, P. & Soriano, C. 2012. Phase contrast synchrotron microtomography reveals the morphology of a partially visible new *Pseudogarypus* in Baltic amber (Pseudoscorpiones: Pseudogarypidae). *Palaeontologia Electronica*, 15: 2;17A,11 p.
- Henderickx, H., Cnudde, V., Masschaele, B., Dierick, M., Vlassenbroeck, J. & Hoorebeke, L. van 2006. Description of a new fossil *Pseudogarypus* (Pseudoscorpiones: Pseudogarypidae) with the use of X-ray micro-CT to penetrate opaque amber. *Zootaxa*, 1305: 41–50.
- Hentz, N. M. 1832. On North American spiders. *American Journal of Science*, 21: 99–109.
- Hentz, N. M. 1845. Descriptions and figures of the Araneides of the United States. *Boston Journal of Natural History*, 5: 189–202.
- Hentz, N. M. 1847. Descriptions and figures of the Araneides of the United States. *Boston Journal of Natural History* 5: 443–478.
- Hentz, N. M. 1850. Descriptions and figures of the Araneides of the United States. *Boston Journal of Natural History*, 6: 18–35, 271–295.
- Herbst, J. F. W. 1798. *Naturgeschichte der Ungeflügelten Insekten. Zweytes Heft*. Berlin, xx pp.
- Hermann, J. F. 1804. *Mémoire Apterologique*. F. G. Levrault, Strasbourg, 144 pp.
- Heyden, C. H. G. von 1826. Versuch einer systematischen Eintheilung der Acariden. *Isis von Oken*, 18: 609–613.
- Heyden, C. H. G. von 1859. Fossile Insekten aus der Rheinischen Braunkohle. *Palaeontographica*, 8: 1–15.
- Hibbert, S. 1836. On the fresh-water limestone of Burdiehouse in the neighbourhood of Edinburgh belonging to the Carboniferous Group of rocks. With supplementary notes on other fresh-water limestones. *Transactions of the Royal Society of Edinburgh*, 13: 169–282.
- Hickman, V. V. 1931. A new family of spiders. *Proceedings of the Zoological Society of London (B)*, 1931: 1321–1328.
- Hickman, V. V. 1944. On some new Australian Apneumonomorphae with notes on their respiratory system. *Papers and Proceedings of the Royal Society of Tasmania*, 1943: 179–195.

- Hickmann, V. V. 1945. A new group of apneumone spiders. *Transactions of the Connecticut academy of Arts and Sciences*, 36: 135–148.
- Hickman, V. V. 1949. Tasmanian littoral spiders with notes on their respiratory systems, habits and taxonomy. *Papers and Proceedings of the Royal Society of Tasmania*, 1948: 31–43.
- Hickman, V. V. 1957. A fossil spider from Tertiary resin from Allendale Victoria. *Proceedings of the Royal Society of Victoria, N.S.*, 69: 25–27.
- Hilton, W. A. 1942. Pantopoda (continued) II. Family Callipallenidae. *Journal of Entomology and Zoology, Pomona College, Claremont*, 34: 38–41.
- Hirschmann, W. 1971. A fossil mite of the genus *Dendrolaelaps* (Acarina, Mesostigmata, Digamasellidae) found in amber from Chiapas, Mexico. *University of California Publications in Entomology*, 63: 69–70.
- Hirst, S. 1923. On some arachnid remains from the Old Red Sandstone (Rhynie Chert bed, Aberdeenshire). *Annals and Magazine of Natural History, Series 9*, 12: 455–474.
- Hoek, P. C. C. 1881. Report on the Pycnogonida dredged by HMS Challenger 1873–76. *Reports of the Scientific Results of the Exploring Vessel HMS Challenger*, 3(10): 1–167.
- Hoff, C. C. 1963. Sternophorid pseudoscorpions, chiefly from Florida. *American Museum Novitates*, 1875: 1–36.
- Holl, F. 1829. *Handbuch der Petrefactenkunde*. Hilscher, Dresden, 489 pp.
- Holland F. D., Jr., Erickson, J. M. & O'Brien, D. E. 1975. *Casterolimulus*: a new Late Cretaceous generic link in Limulid lineage. Studies in Paleontology and Stratigraphy. *Bulletin of American Paleontology*, 62: 235–249.
- Holmberg, E. L. 1882. Observations à propos du sous-ordre des araignées territélaires (Territelariae), spécialement du genre nordaméricain *Catadysas* Hentz et de la sous-famille Mecicothrioidae, Holmberg. *Boletín de la Academia Nacional de Ciencias en Córdoba (Argentina)*, 4: 153–174.
- Holmberg, E. L. 1883. *Neothereutes darwini* Holmb., representante de una nueva familia de Citrigadas. *Boletín de la Academia Nacional de Ciencias en Córdoba (Argentina)*, 5: 35–48.
- Hong Y.-c. 1982. [Study on new spider genus in amber.] *Science in China*, 24(12): 1500–1515. [In Chinese]
- Hong Y.-c. 1983a. [Discovery of a Miocene scorpion from the diatoms of Shanwang in Shandong Province.] *Bulletin of the Tianjin Institute of Geology and Mineral Resources*, 8, 17–21. [In Chinese]
- Hong Y.-c. 1983b. [Discovery of new fossil pseudoscorpiononods in amber.] *Bulletin of the Tianjin Institute of Geology and Mineral Resources*, 8: 24–29. [In Chinese]
- Hong Y.-c. 1984. Arachnida. 185–187 In Tianjin Institute of Geology and Mineral Resources (eds). *Palaeontological Atlas of North China II. Mesozoic Volume*. Geological Publishing House, Beijing. [In Chinese with English summary]
- Hong Y.-c. 1985. *Fossil Insects, scorpionids and araneids in the diatoms of Shanwang*. Geological Publishing House, Beijing, 80 pp.

- Hopkins, D. M., Giterman, R. E. & Matthews, J. V. 1976. Interstadial mammoth remains and associated pollen and insect fossils, Kotzebue Sound area, northwestern Alaska. *Geology*, 4: 169–173.
- Hradská, I. & Dunlop, J. A. 2013. New records of Pennsylvanian trigonotarbid arachnids from West Bohemia, Czech Republic. *The Journal of Arachnology*, 41: 335–341.
- Huang D.-y., Selden, P. A. & Dunlop, J. A. 2009. Harvestmen (Arachnida: Opiliones) from the Middle Jurassic of China. *Naturwissenschaften*, 96: 955–962.
- Huber, B. A. 2003. Southern African pholcid spiders revision and cladistic analysis of *Quamtana gen. nov.* and *Spermophora* Hentz (Araneae: Pholcidae), with notes on male–female covariation. *Zoological Journal of the Linnean Society*, 139: 477–527.
- Huber, B. A. & Wunderlich, J. 2006. Fossil and extant species of the genus *Leptopholcus* in the Dominican Republic, with the first cases of egg-parasitism in pholcid spiders (Araneae: Pholcidae). *Journal of Natural History*, 40: 2341–2360.
- Hull, J. E. 1920. The spider family Linyphiidae: an Essay in Taxonomy. *Vasculum*, 6: 7–11.
- Hünicken, M. A. 1980. A giant fossil spider (*Megarachne servinei*) from Bajo de Véliz, Upper Carboniferous, Argentina. *Boletín de la Academia Nacional de Ciencias, Córdoba*, 53: 317–341.
- Hunter, J. R. S. 1886. Notes on the discovery of a fossil scorpion (*Paleophonus caledonicus*) in the Silurian strata of Logan water. *Transactions of the Geological Society of Glasgow*, 8: 169–170.
- Jackson, R. T. 1906. A new species of fossil *Limulus* from the Jurassic of Sweden. *Arkiv för Zooogi*, 3(11): 1–7.
- Jaekel, O. 1914. Ein großer *Pterygotus* aus dem rheinischen Unterdevon. *Palaeontologische Zeitschrift*, 1: 379–382.
- Jävi, T. H. 1912/14. Das Vaginalsystem der Sparassiden. *Annales Academiae Scientiarum Fennicae*, A4: 1–248.
- Jell, P. A. & Duncan, P. M. 1986. Invertebrates, mainly insects, from the freshwater Lower Cretaceous Koonwarra fossil bed (Korumburra Group), South Gippsland, Victoria. *Memoirs of the Association of Australian Palaeontology*, 3: 111–205.
- Jeram, A.J. 1994a. Scorpions from the Viséan of East Kirkton, West Lothian, Scotland, with a revision of the infraorder Mesoscorpionina. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 84: 283–299.
- Jeram, A.J. 1994b. Carboniferous Orthosterni and their relationship to living scorpions. *Palaeontology*, 37: 513–550.
- Jocqué, R. 1994. Halidae, a new spider family from Madagascar (Araneae). *Bulletin of the British arachnological Society*, 9: 281–289.
- Jocqué, R. 2001. Chummidae, a new spider family (Arachnida, Araneae) from South Africa. *Journal of Zoology, London*, 254: 481–493.
- Jones, T. R. & Woodward, H. 1888. On some Scandinavian Phyllocarida. *Geological Magazine, New Series, Decade 3*, 5: 145–150.

- Jones, T. R. & Woodward, H. 1899. Contributions to fossil Crustacea. *Geological Magazine, New Series, Decade 4*, 6: 388–395.
- Jordan, H. & Meyer, H. von 1854. Ueber die Crustaceen der Steinkohlenformation von Saarbrücken. *Palaeontographica*, 4: 1–15.
- Judson, M. [L. I.] 2003. Baltic amber pseudoscorpions (Arachnida: Chelonethi): a new species of *Neobisium* (Neobisiidae) and the status of *Obisium rathkii* Koch and Berendt. *Geodiversitas*, 25: 445–450.
- Judson, M. L. I. 2007. First fossil record of the pseudoscorpion family Pseudochiridiidae (Arachnida, Chelonethi, Cheirioidea) from Dominican amber. *Zootaxa*, 1393: 45–51.
- Judson, M. L. I. 2009. Cheliferoid pseudoscorpions (Arachnida, Chelonethi) from the Lower Cretaceous of France. *Geodiversitas*, 31: 61–71.
- Judson, M. L. I. 2010. Redescription of *Chelifer eucarpus* Dalman (Arachnida, Chelonethi, Withiidae) and first records of pseudoscorpions in copal from Madagascar and Colombia. *Palaeodiversity*, 3: 33–42.
- Judson, M. L. I. & Mąkol, J. 2009. A mite of the family Tanaupodidae (Arachnida, Acari, Parasitengona) from the Lower Cretaceous of France. *Geodiversitas*, 31: 41–47.
- Judson, M. [L. I.] & Wunderlich, J. 2003. Rhagidiidae (Acari, Eupodoidea) from Baltic amber. *Acta zoologica cracoviensa*, 46 (suppl.–Fossil Insects): 147–152.
- Jux, U. 1982. *Somaspidion hammapheron* n.gen. n.sp. – ein Arachnid aus dem Oberkarbon der subvaristischen Saumsenke NW Deutschlands. *Paläontologische Zeitschrift*, 56: 77–86.
- Kaddumi, H. F. 2007. *Amber of Jordan: the oldest prehistoric insects in fossilized resin. Second edition*. Eternal River Museum of Natural History, Amman, Jordan, 224 pp.
- Karpinen, E. & Koponen, M. 1973. The subfossil oribatid fauna of Piilonsuo, a bog in southern Finland. *Annales entomologici Fennici*, 39: 22–32.
- Karpinen, E. & Koponen, M. 1974. Further observations on subfossil remains of oribatids (Acar., Oribatei) and insects in Piilonsuo, a bog in southern Finland. *Annales entomologici Fennici*, 40: 172–175.
- Karpinen, E., Krivolutsky, D. A., Koponen, M., Kozlovskaja, L. S., Laskova, L. M. & Viitalsaari, M. 1979. List of subfossil oribatid mites (Acarina, Oribatei) of northern Europe and Greenland. *Annales entomologici Fennici*, 45: 103–108.
- Karsch, F. 1879. Arachnologische Beiträge. *Zeitschrift für die gesammten Naturwissenschaften*, 52: 534–562.
- Karsch, F. 1880a. Arachnologische Blätter. I. Ueber *Corinna* (C. L. Koch) und ihre Verwandtschaften. *Zeitschrift für die gesammten Naturwissenschaften*, 53: 373–378.
- Karsch, F. 1880b. Arachnologische Blätter. X. Scorpionologische Fragmente. *Zeitschrift für die gesammten Naturwissenschaften*, 53: 404–409.
- Karsch, F. 1882. Ueber ein neues Spinnenthier aus der Schlesischen Steinkohle und die Arachnoiden überhaupt. *Zeitschrift der Deutschen geologischen Gesellschaft*, 34: 556–561.
- Karsch, F. 1884. Neue Milben in Bernstein. *Berliner Entomologische Zeitschrift*, 28: 175–176.

- Keegan, H. L., Yunker, C. E. & Baker, E. W. 1960. Malaysian parasites. XLVI. *Hystrichonyssus turneri*, n.sp. n.g. representing a new subfamily of Dermasyddidae (Acarina) from a Malayan porcupine. *Studies from the Institute for Medical Research Federation of Malaya*, 107: 455–473.
- Keferstein, C. 1834. *Die Naturgeschichte des Erdkörpers in ihren ersten Grundzügen*, Vol. 2. F. Fleischer, Leipzig, 896 pp.
- Keifer, H. H. 1966. [untitled.] *Californian Department of Agriculture. Eriophyid Series*, B-21: 1–20.
- Keirans, J. E., Lane, R. S. & Cauble, R. 2002. A series of larval *Amblyomma* species (Acari : Ixodidae) from amber deposits in the Dominican Republic. *International Journal of Acarology*, 28: 61–66.
- Kethley, J. B. 1974. Developmental chaetotaxy of a paedomorphic celaenopsoid, *Neotenogynium malkini* n.g., sp. (Acari: Parasitiformes: Neotenogyniidae, n. fam.) associated with millipedes. *Annals of the Entomological Society of America*, 67: 571–579.
- Kethley, J. B. 1977a. The Status of *Hybalicus* Berlese, 1913 and *Oehsercheste* Jacot, 1939 (Acari: Acariformes: Endeostigmata). *Fieldiana Zoology*, 72: 59–64.
- Kethley, J. B. 1977b. An unusual Parantennuloid, *Philodana johnstoni* n.g., n.sp. (Acari: Parasitiformes: Philodanidae, n. fam.) associated with *Neatus tenebrioides* (Coleoptera: Tenebrionidae) in North America. *Annals of the Entomological Society of America*, 70: 487–494.
- Kethley, J. B. 1979. A cladistic analysis of the Trigynaspida (Acari: Parasitiformes) with a review of the higher categories and nominate taxa. In Piffl, E. (ed). *Proceedings of the 4th International Congress of Acarology – Saalfelden (Austria)*. Akadémiai Kiakó, Budapest, pp. 459–466.
- Kethley, J. B. 1989. Proteonematalycidae (Acari: Acariformes), a new mite family from fore-dune sand of Lake Michigan. *International Journal of Acarology*, 15: 209–217.
- Kethley, J. B., Norton, R. A., Bonamo, P. M. & Shear, W. A. 1989. A terrestrial alicorhagiid mite (Acari: Acariformes) from the Devonian of New York. *Micropaleontology*, 35: 367–373.
- Kew, H. W. 1911. A synopsis of the false scorpions of Britain and Ireland. *Proceedings of the Royal Irish Academy (B)*, 29: 38–64.
- Keyserling, E. 1877. Ueber amerikanische Spinnenarten der Unterordnung Citigradae. *Verhandlungen der Zoologisch-Biologischen Gesellschaft in Wien*, 26: 609–708.
- Keyserling, E. 1880a. *Die Spinnen Amerikas, I. Laterigradae*. Nürnberg, 1, 283 pp.
- Keyserling, E. 1880b. Neue Spinnen aus Amerika. I. *Verhandlungen der Zoologisch-Biologischen Gesellschaft in Wien*, 29: 293–349.
- Keyserling, E. 1882. Neue Spinnen aus Amerika. III. *Verhandlungen der Zoologisch-Biologischen Gesellschaft in Wien*, 31: 269–314.
- Keyserling, E. 1884. *Die Spinnen Amerikas. Theridiidae*. Nürnberg, 2, 222 pp.
- Khaustov A. A. 2000. Bembidiacaridae, a new family of mites (Acari: Heterostigmata) associated with carabid beetles of the genus *Bembidion* (Coleoptera: Carabidae). *Acarina*, 8: 3–8.

- Khaustov, A. A. & Perkovsky, E. E. 2010. The first fossil record of mites of the family Pyemotidae (Acari: Heterostigmata), with description of a new species from Rovno Amber. *Palaeontological Journal*, 44: 418–421.
- Khaustov, A. A. & Poinar jr., G. O. 2010. *Protoresinacarus brevipedis* gen. n., sp. n. from Early Cretaceous Burmese amber: the first fossil record of mites of the family Resinacaridae (Acari: Heterostigmata: Pyemotoidea). *Historical Biology*: 23: 219–222.
- Kirchner, H. 1923. *Limulus Sandbergi* n. sp. aus dem fränkischen oberen Buntsandstein (Plattensandstein). *Centralblatt für Mineralogie, Geologie und Paläontologie*, 20: 634–639.
- Kim, C. M. 2008. Euphsalozerconidae, a new mesostigmatid mite family (Acari: Mesostigmata: Trigynaspida: Aenictequoidea). *Acarologia*, 48: 33–38.
- Kim, J.-p. & Nam, K.-s. 2008. [Mesozoic spider (Araneae: Pisauridae) from Korea.] *Korean Arachnology*, 24: 119–125. [in Korean with English summary]
- Kim, J.-p. & Nam, K.-s. 2008. [Mesozoic spider (Aranea:Lycosidae) from China.] *Korean Arachnology*, 28: 35–45. [in Korean with English summary]
- Kin, A. & Błażejowski, B. 2014. The horseshoe crab of the genus *Limulus*: living fossil or stabilomorph? *PLoS ONE*, 9(10): e108036.
- Kishida, K. 1930. A new scheme of classification of spider families and genera. *Lansania*, 2: 33–43.
- Kjellesvig-Waering, E. N. 1934. Note on a new eurypterid from the Moscow Shales of New York. *American Journal of Science, 5th Series*, 27: 386–387.
- Kjellesvig-Waering, E. N. 1948a. Two new eurypterids from the Silurian of Indiana. *Journal of Paleontology*, 22: 465–472.
- Kjellesvig-Waering, E. N. 1948b. The Mazon Creek Eurypterid: A revision of the genus *Lepidoderma*. *Scientific Papers, Illinois*, 3(4): 1–48.
- Kjellesvig-Waering, 1950a. A new Silurian Hughmilleria from West Virginia. *Journal of Paleontology*, 24: 226–228.
- Kjellesvig-Waering, 1950b. A new Silurian Eurypterid from Florida. *Journal of Paleontology*, 24: 229–231.
- Kjellesvig-Waering, E. N. 1951. Downtonian (Silurian) Eurypterida from Perton, near Stoke Edith, Herefordshire. *Geological Magazine*, 88: 1–24.
- Kjellesvig-Waering, E. N. 1954. Note on a new Silurian (Downtonian) scorpion from Shropshire, England. *Journal of Palaeontology*, 28: 485–486.
- Kjellesvig-Waering, E. N. 1955. A new phyllocarid and eurypterid from the Silurian of Florida. *Journal of Paleontology*, 29: 295–297.
- Kjellesvig-Waering, E. N. 1958. The genera, species and subspecies of the family Eurypteridae Burmeister, 1845. *Journal of Paleontology*, 32: 1107–1148.
- Kjellesvig-Waering, E. N. 1959. A taxonomic review of some late Paleozoic Eurypterida. *Journal of Palaeontology*, 33: 251–256.

- Kjellesvig-Waering, E. N. 1961a. Eurypterida of the Devonian Holland Quarry Shale of Ohio. *Fieldiana, Geology*, 14(5): 79–98.
- Kjellesvig-Waering, E. N. 1961b. The Silurian Eurypterida of the Welsh Boderland. *Journal of Paleontology*, 35: 251–256.
- Kjellesvig-Waering, E. N. 1963a. Revision of some Upper Devonian Styloinuridae (Eurypterida) from New York and Pennsylvania. *Journal of Paleontology*, 37: 490–495.
- Kjellesvig-Waering, E. N. 1963b. Pennsylvanian invertebrates of the Mazon Creek area, Illinois, Eurypterida. *Fieldiana, Geology*, 14(9): 169–197.
- Kjellesvig-Waering, E. N. 1964a. A synopsis of the Family Pterygotidae Clarke and Ruedemann 1912 (Eurypterida). *Journal of Paleontology*, 38: 331–361.
- Kjellesvig-Waering, E. N. 1964b. Eurypterida: Notes on the subgenus *Hughmilleria* (*Nanahughmilleria*) from the Silurian of New York. *Journal of Paleontology*, 38: 410–412.
- Kjellesvig-Waering, E. N. 1966b. Silurian scorpions of New York. *Journal of Paleontology*, 40: 359–375.
- Kjellesvig-Waering, E. N. 1966c. The scorpions of Trinidad and Tobago. *Caribbean Science*, 6: 123–135.
- Kjellesvig-Waering, E. N. 1969. A new phalangiotarbid (Arachnida) from the Pennsylvanian of Oklahoma. *Journal of Paleontology*, 43: 1280–1282.
- Kjellesvig-Waering, E. N. 1971. A new Downtonian stylonurid from Central England (Silurian, Eurypterida). *Journal of Paleontology*, 45: 538–539.
- Kjellesvig-Waering, E. N. 1972. *Brontoscorpius anglicus*: a giant Lower Palaeozoic scorpion from central England. *Journal of Paleontology*, 46: 39–42.
- Kjellesvig-Waering, E. N. 1973. A new Silurian *Slimonia* (Eurypterida) from Bolivia. *Journal of Paleontology*, 47: 549–550.
- Kjellesvig-Waering, E. N. 1979. Eurypterids. In Jaanusson, V., Laufeld, S. & Skoglund, R. (eds). Lower Wenlock faunal and floral dynamics – Vattenfallet section, Gotland. *Sveriges Geologiska Undersökning, Serie C*, NR 762, Årsbok 73 NR, 3: 121–136.
- Kjellesvig-Waering, E. N. 1986. A restudy of the fossil Scorpionida of the world. *Palaeontographica Americana*, 55: 1–287.
- Kjellesvig-Waering, E. N. & Caster, K. E. 1955. The Pterygotidae of the Silurian Vernon Shales of New York. *Journal of Paleontology*, 29: 1041–1047.
- Kjellesvig-Waering, E. N. & Heubusch, C. A. 1962. Some Eurypterida from the Ordovician and Silurian of New York. *Journal of Paleontology*, 36: 211–221.
- Kjellesvig-Waering, E. N. & Leutze, W. P. 1966. Eurypterida from the Silurian of West Virginia. *Journal of Paleontology*, 40: 1109–1122.
- Kjellesvig-Waering, E. N. & Størmer, L. 1952. The *Dolichopterus*–*Strobilopterus* group in the Eurypterida. *Journal of Palaeontology*, 26: 659–661.

- Klompen, H. & Grimaldi, D. 2001. First Mesozoic record of a parasitiform mite: a larval argasid tick in Cretaceous amber (Acari: Ixodida: Argasidae). *Annals of the Entomological Society of America*, 94: 10–15.
- Kobayashi, T. 1933. On the occurrence of Xiphosuran remains in Chosen (Korea). *Japanese Journal of Geology and Geography*, 10: 175–182.
- Koçak, A. Ö. & Kemal, M. 2008. New synonyms and replacement names in the genus group taxa of Araneida. *Centre for entomological Studies, Miscellaneous Papers*, 139–140: 1–4.
- Koch, C. L. 1829–1844. Arachniden. In Panzer (ed). *Faunae Insectorum Germaniae initia. Fortgesetzt von Herrich-Schäffer, Hefte 111–190*. Regensburg. [1833, Hefte 119–121]
- Koch, C. L. 1834. Arachniden. In Panzer (ed). *Faunae Insectorum Germaniae initia. Hefte 122–125*, 127. Regensburg.
- Koch, C. L. 1835. Arachniden. In Panzer (ed). *Faunae Insectorum Germaniae initia. Hefte 128–131*. Regensburg.
- Koch, C. L. 1837. *Uebersicht des Arachnidensystems* 1. C. H. Zeh'sche Buchhandlung, Nürnberg, 39 pp.
- Koch, C. L. 1839a. *Uebersicht des Arachnidensystems* 2. C. H. Zeh'sche Buchhandlung, Nürnberg, 38 pp.
- Koch, C. L. 1839b. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Sechster Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 156 pp.
- Koch, C. L. 1839c. *Deutschlands Crustaceen, Myriapoden und Arachniden. Hefte 23–30*.
- Koch, C. L. 1842a. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Neunter Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 108 pp.
- Koch, C. L. 1842b. *Uebersicht des Arachnidensystems* 3. C. H. Zeh'sche Buchhandlung, Nürnberg, 130 pp.
- Koch, C. L. 1843a. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Zehnter Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 142 pp.
- Koch, C. L. 1843b. *Uebersicht des Arachnidensystems* 3. C. H. Zeh'sche Buchhandlung, Nürnberg, 130 pp
[continuation of 1842b; see above].
- Koch, C. L. 1844. Systematische Übersicht über die Ordnung der Zecken. *Archiv für Naturgeschichte*, 1: 217–239.
- Koch, C. L. 1846. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Dreizehnter Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 234 pp.
- Koch, C. L. 1847. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Vierzehnter Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 210 pp.
- Koch, C. L. 1851. *Übersicht des Arachnidensystems* 5. C. H. Zeh'sche Buchhandlung, Nürnberg, 104 pp.
- Koch, C. L. & Berendt, G. C. 1854. Die im Bernstein befindlichen Myriapoden, Arachniden und Apteriden der Vorwelt. In Berendt, G. C. *Die in Bernstein befindlichen organischen Reste der Vorwelt gesammelt in Verbindung mit mehreren bearbeitet und herausgegeben* 1. Berlin, Nicolai, 124 pp.
- Koch, L. 1866. *Die Arachniden-Familie der Drassiden*. 1–6. J. L. Lotzbeck, Nürnberg, 352 pp.

- Koch, L. 1871–1883. *Die Arachniden Australiens nach der Natur beschrieben und abgebildet*. Bauer & Raspe, Nürnberg, 1489 pp.
- Koch, L. 1873. *Uebersichtliche Darstellung der europäischen Chernetiden (Pseudoscorpione)*. Bauer und Raspe, Nürnberg, **xx** pp.
- Konikiewicz, M & Mąkol, J. 2014. A fossil Paratrombiinae mite (Actinotrichida: Trombidioidea) from the Rovno amber, Ukraine. *Zootaxa*, 3847: 583–589.
- Kraepelin, K. 1899. Zur Systematik der Solifugen. *Mitteilungen aus dem Naturhistorischen Museum in Hamburg*, 16: 195–258.
- Kraepelin, K. 1901. Palpigradi und Solifugae. *Tierreich*, 12: i–x, 1–159.
- Kraepelin, K. 1905. Die geographische Verbreitung der Skorpione. - *Zoologische Jahrbücher, Abtheilung für Systematik*, 22: 321–364.
- Kramer, P. 1879. Neue Acariden. *Archiv für Naturgeschichte*, 45: 13–16.
- Kramer, P. 1885. Ueber Halarachne Halichoeri, Allm. *Zeitschrift für Naturwissenschaften*, 58: 1–31.
- Krause, T., Hauschke, N. & Wilde, V. 2009. Ein Limulide aus den Gelben Basisschichten des Oberen Muschelkalks von Ohrdruf bei Gotha (Thüringen). *Geowissenschaftliche Mitteilungen von Thüringen*, 13: 163–168.
- Kratochvíl, J. 1958. Höhlenweberknechte Bulgariens (Palpatores – Nemastomatidae). *Acta Academiae Scientiarum Čechoslovenicae Basis Brunensis*, 30: 523–576.
- Krivolutsky, D. A. & Krasilov, B. A. 1977. Oribatid mites from Upper Jura deposits of USSR. 16–24. In Skarlato, O. A. & Balashov, Y. S. (eds) *Morphology and Diagnostics of Mites*. Zoological Institute, Leningrad, 85 pp. [in Russian]
- Krüger, J. & Dunlop, J. A. 2010. Schizomids (Arachnida: Schizomida) from Dominican Republic amber. *Alavesia*, 3: 43–53.
- Kues, B. S. & Kietzke, K. K. 1981. A large assemblage of a new eurypterids from the Red Tanks Member, Madera Formation (Late Pennsylvania - Early Permian) of New Mexico. *Journal of Paleontology*, 55: 709–729.
- Kühl, G., Poschmann, M. & Rust, J. 2013. A ten-legged sea spider (Arthropoda: Pycnogonida) from the Lower Devonian Hunsrück Slate (Germany). *Geological Magazine*, 150: 556–564.
- Kühl, G., Bergamnn, A., Dunlop, J. A., Garwood, R. J. & Rust, J. 2012. Redescription and palaeobiology of *Palaeoscorpius devonicus* Lehmann, 1944 from the Lower Devonian Hunsrück Slate of Germany. *Palaeontology*, 55: 775–787.
- Kulczynski, L. 1902. Species Oribatinarum (Oudms.) (Damaeinorum Michael) in Galicia collectae. *Dissertationum mathematicarum et physicarum Academiae Litterarum Cracoviensis*, 42: 1–50.
- Kulicka, R. 1990. The list of animal inclusions in Baltic amber from collection of the Museum of Earth in Warsaw. *Prace Muzeum Ziemi*, 41: 144–146.

- Kury, A. B. 2003. Annotated catalogue of the Laniatores of the New World (Arachnida, Opiliones). *Revista Ibérica de Aracnología*, Volumen especial monográfico 1: 1–337.
- Kury, A. B. & Pérez González, A. 2002. A new family of Laniatores from northwestern South America (Arachnida, Opiliones). *Revista Ibérica de Aracnología*, 6: 3–11.
- Kušta, J. 1883. *Anthracomartus krejci*, eine neue Arachnide aus dem Böhmischem Karbon. *Sitzungsberichte der Königlich Böhmischen Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1883: 7.
- Kušta, J. 1884a. Neue Arachniden aus der Steinkohlenformation von Rakonitz. *Sitzungsberichte der Königlich Böhmischen Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1884: 398–401.
- Kušta, J. 1884b. *Thelyphonus bohemicus* n. sp., ein fossiler Geisselscorpion aus der Steinkohlenformation von Rakonitz. *Sitzungsberichte der Königlich Böhmischen Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1884: 186–191.
- Kušta, J. 1885. Neue fossile Arthropoden aus dem Noeggarathienschiefer von Rakonitz. *Sitzungsberichte der Königlich Böhmischen Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1885: 1–7.
- Kušta, J. 1888. O nových arachnidech z karbonu Rakovnického. (Neue Arachniden aus der Steinkohlenformation bei Rakonitz). *Sitzungsberichte der Königlich Böhmischen Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1888: 194–208.
- Kutorga, S. 1838. *Beitrag zur Kenntnis der organischen Überreste des Kupfersandsteins am westlichen Abhange des Urals*. St. Petersburg, 38 pp.
- Kuznetsov, N. N., Khaustov, A. A. & Perkovsky, E. E. 2010. First record of mites of the family Stigmaeidae (Acari, Raphignathoidea) from Rovno amber with description of a new species of the genus *Mediolata*. *Vestnik zoologii*, 44: 545–547.
- Lamarck, J. B. P. A. 1801. *Système des animaux sans vertèbres*. Lamarck and Deterville, Paris, xx pp.
- Lamont, A. 1955. Scottish Silurian Chelicerata. *Transactions of the Edinburgh Geological Society*, 16: 200–216.
- Lamsdell, J. C. 2011. The eurypterid *Stoermeropterus conicus* from the Lower Silurian of the Pentland Hills, Scotland. *Monographs of the Palaeontographical Society*, 165: 1–84.
- Lamsdell, J. C. 2012. Redescription of *Drepanopterus pentlandicus* Laurie, 1892, the earliest known mycteroiid (Chelicerata: Eurypterida) from the early Silurian (Llandovery) of the Pentland Hills, Scotland. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 103: 77–103.
- Lamsdell, J. C. 2013a. Revised systematics of the Palaeozoic ‘horseshoe crabs’ and the myth of the monophyletic Xiphosura. *Zoological Journal of the Linnaean Society*, 167: 1–27.
- Lamsdell, J. C. 2013b. Redescription of *Drepanopterus pentlandicus* Laurie, 1892, the earliest known mycteroiid (Chelicerata: Eurypterida) from the early Silurian (Llandovery) of the Pentland Hills,

- Scotland. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 103: 77–103.
- Lamsdell, J. C. & Selden, P. A. 2013. Babes in the wood – a unique window into sea scorpion ontogeny. *BMC Evolutionary Biology* 13: 98.
- Lamsdell, J. C., Braddy, S. J. & Tetlie, O. E. 2010. The systematics and phylogeny of the Stylocephala (Arthropoda: Chelicerata: Eurypterida). *Journal of Systematic Palaeontology*, 8: 49–61.
- Lamsdell, J. C., Hoşgör, İ & Selden, P. A. 2013. A new Ordovician eurypterid (Arthropoda: Chelicerata) from southeast Turkey: evidence for a cryptic Ordovician record of Eurypterida. *Gondwana Research*, 23: 354–366.
- Lamsdell, J. C., Simonetta, L. & Selden, P. A. 2013. First eurypterid from Italy: a new species of *Adelophthalmus* (Chelicerata: Eurypterida) from the Upper Carboniferous of the Carnic Alps (Friuli, NE Italy). *Revista Italiana di Paleontologia et Stratigrafia*, 119: 147–151.
- Lamsdell, J. C., Xue, J.-h. & Selden, P. A. 2013. A horseshoe crab (Arthropoda: Chelicerata: Xiphosura) from the Lower Devonian (Lochkovian) of Yunnan, China. *Geological Magazine*, 150: 367–370.
- Lamsdell, J. C., Braddy, S. J., Loeffler, E. J. & Dineley, D. L. 2010. Early Devonian stylocephalan eurypterids from Arctic Canada. *Canadian Journal of Earth Sciences*, 47: 1405–1415.
- Lane, R. S. & Poinar jr., G. O. 1986. First fossil tick (Acarina: Ixodidae) in new world amber. *International Journal of Acarology*, 12: 75–78.
- Latreille, P. A. 1795. Observations sur la variété des organes de la bouche des tiques, et distribution méthodique des insectes de cette famille d'après les caractères établis sur la conformation de ces organes. *Magasin Encyclopédique, ou Journal des Sciences, des Lettres et des Arts*, 4: 15–20.
- Latreille, P. A. 1796. *Précis de caractères génériques des insectes, disposés dans un ordre naturel*. Prévot, Paris, **xx** pp.
- Latreille, P. A. 1802. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes*. Dufart, Paris, **xx** pp.
- Latreille, P. A. 1804a. Tableau méthodique des Insectes. *Nouveau Dictionnaire d'histoire naturelle*, 24: 129–200.
- Latreille, P. A. 1804b. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes*, Vol. 7. F. Dufart, Paris, pp. 144–305.
- Latreille, P. A. 1806. *Genera Crustaceorum et Insectorum*. Vol. 1. A. Koenig, Paris, pp. 82–127.
- Latreille, P. A. 1809. *Genera Crustaceorum et Insectorum*. Vol. 4. Paris, pp. 73–371.
- Latreille, P. A. 1810. *Considérations générales sur l'Ordre Naturel des Animaux composant les Classes des Crustacés, des Arachnides et des Insectes*. Paris, 446 pp.
- Latreille, P. A. 1819. [Articles sur les Araignées]. *Nouveau Dictionnaire d'histoire naturelle* 30-35 : ?? pp.
- Latreille, P. A. 1829. Les Arachnides. In Cuvier, G (ed.) *Le règne animal, nouv. éd.* Paris, pp. 206–291.

- Laurentiaux-Viera, F. & Laurentiaux, D. 1961. *Prothelyphonus neerlandicus*, nov. sp., Uropyge du Westphalien du Limbourg Hollandais. *Mededelingen van de Geologische Stichting, N.S.*, 13: 29–34.
- Laurentiaux-Viera, F. & Laurentiaux, D. 1963. Sur quelques restes nouveaux d'Arachnides du terrain houiller. *Annales de la Société Géologique du Nord*, 83: 23–29.
- Laurie, M. 1892. On some eurypterid remains from the Upper Silurian rocks of the Pentland Hills. *Transactions of the Royal Society of Edinburgh*, 37: 151–162.
- Laurie, M. 1896. Further notes on the anatomy and development of scorpions, and their bearing on the classification of the order. *Annals and Magazine of Natural History, series 6*, 17: 185–193.
- Laurie, M. 1899. On a Silurian scorpion and some additional eurypterid remain from the Pentland Hills. *Transactions of the Royal Society of Edinburgh*, 39: 575–590.
- Leach, W. E. 1815. A tabular view of the external characters of four classes of animals which Linné arranged under Insecta; with the distribution of the genera composing three of these classes into orders, andc. And descriptions of several new genera and species. *Transactions of the Linnean Society of London*, 11: 306–400.
- Leach, W. E. 1819. *Dictionnaire des Sciences Naturelles*, Vol. 14. Paris, pp. 537–538.
- Leary, R.L. 1980. *Labriscorpio alliedensis*, a new Carboniferous scorpion from Rock Island County, Illinois. *Journal of Paleontology*, 54: 1255–1257.
- Leech, R. & Matthews Jr., J. V. 1971. *Xysticus archaeopalpus* (Arachnida: Thomisidae), a new species of crab spider from Pliocene sediments in western Alaska. *Canadian Entomologist*, 103: 1337–1340.
- Legg, D. A. 2014. *Sanctacaris uncata*: the oldest chelicerate (Arthropoda). *Naturwissenschaften*, 101: 1065–1073.
- Lehmann, W.M. 1944. *Palaeoscorpius devonicus* n. g., n. sp., ein Skorpion aus dem rheinischen Unterdevon. *Neues Jahrbuch für Paläontologie, Monatshefte, B*: 177–185.
- Lehtinen, P. T. 1967. Classification of the cribellate spiders and some allied families, with notes on the evolution of the suborder Araneomorpha. *Annales Zoologici Fennici*, 4: 199–468.
- Lehtinen, P. T. 1981. New Holothyrida (Arachnida, Anactinotrichida) from New Guinea and South America. *Acarologia*, 22: 3–13.
- Lenz, H. 1886. Beiträge zur Kenntniss der Spinnenfauna Madagascars. *Zoologische Jahrbücher, Systematik*, 1: 379–408.
- Leutze, W. P. 1958. Eurypterids from the Silurian Tymochtee dolomite of Ohio. *Journal of Paleontology*, 32: 937–942.
- Leutze, W. P. 1961. Arthropods from the Syracuse Formation, Silurian of New York. *Journal of Paleontology*, 35: 49–64.
- Levy, G. 2007. The first troglobite scorpion from Israel and a new chactoid family (Arachnida: Scorpiones). *Zoology in the Middle East*, 40: 91–96.

- Li S.-q. & Wunderlich, J. 2008. Sinopimoidae, a new spider family from China (Arachnida, Araneae). *Acta zoootaxonomica sinica*, 33: 1–6.
- Lin Q.-b., Zhang, Z.-f. & Wang, B.-z. 1989. New evidences for Miocene climatic optimum event—review on the Miocene spider fossils from Shanwang collection. *Proceedings of International Symposium on Pacific Neogene and Marine Events*. Nanjing University Press, pp. 137–147.
- Lin Q.-b., Yao Y.-m., Xiang W.-d. & Xia Y.-r. 1988. An Oligocene micropalaeoentomofauna from Gubei district of Shandong and its ecological environment. *Acta Micropalaeontologica Sinica*, 5: 331–345.
- Lindquist, E. E. & Moraza, M. L. 1993. Pyrosejidae, a new family of trigynaspid mites (Acari: Mesostigmata: Cercomegistina) from Middle America. *Acarologia*, 34: 283–307.
- Lindquist, E. E. & Palacios-Vargas, J. G. 1991. Proterorhagiidae (Acari: Endeostigmata), a new family of rhagidiid-like mites from Mexico. *Acarologia*, 32: 341–363.
- Linnaeus, C. 1758. *Systema naturae*, 10th edition. Vol 1. L. Salvii, Holmiae.
- Loman, J. C. C. 1900. Ueber die geographische Verbreitung der Opilioniden. *Zoologische Jahrbücher, Systematik*, 16: 71–104.
- Lourenço, W. R. 1995. Description de trois nouveaux genres et quatre nouvelles espèces de scorpions Buthidae de Madagascar. *Bulletin du Muséum National d'Histoire Naturelle* (4), 17A: 95–106.
- Lourenço, W. R. 1996a. Faune de Madagascar. 87. Scorpions (Chelicerata, Scorpiones). Muséum National d'Histoire Naturelle, Paris, 102 pp.
- Lourenço, W. R. 1996b. Premier cas connu d'un sub-fossile de scorpion dans le copal de Madagascar. *Compte Rendus de l'Académie des Sciences, Paris, Sér. IIa*, 323: 889–891.
- Lourenço, W. R. 1998. Panbiogeographie, les distributions disjointes et le concept de famille relictuelle chez les Scorpions. *Biogeographica*, 74: 133–144.
- Lourenço, W. R. 2000a. More about the Buthoidea of Madagascar, with special references to the genus *Tityobuthus* Pocock (Scorpiones, Buthidae). *Revue suisse de Zoologie*, 107: 721–736.
- Lourenço, W. R. 2000b. Premier cas d'un sub-fossile d'araignée appartenant au genre *Archaea* Koch and Berendt (Archaeidae) dans le copal de Madagascar. *Comptes rendus de l'Académie des Sciences Paris, Sciences de la Terre et des planètes*, 330: 509–512.
- Lourenço, W. R. 2001. A remarkable scorpion fossil from the amber of Lebanon. Implications for the phylogeny of Buthoidea. *Comptes rendus de l'Académie des Sciences Paris, Sciences de la Terre et des planètes*, 332: 641–646.
- Lourenço, W. R. 2002. The first scorpion fossil from the Cretaceous amber of Myanmar (Burma). New implications for the phylogeny of Buthoidea. *Comptes Rendus Palevol*, 1: 97–101.
- Lourenço, W. R. 2003. The first scorpion fossil from the Cretaceous amber of France. New implications for the phylogeny of Chactoidea. *Comptes Rendus Palevol*, 2: 213–219.

- Lourenço, W. R. 2004. Description of a further species of fossil scorpion in Baltic amber. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 3: 1886–1889.
- Lourenço, W. R. 2009a. A new sub-fossil scorpion of the genus *Microcharmus* Lourenço from Malagasy copal (Scorpiones, Microcharmidae). *Boletín Sociedad Entomológica Aragonesa*, 44: 135–137.
- Lourenço, W. R. 2009b. A new species of *Tityus* C. L. Koch, 1836 (subgenus *Brazilotityus* Lourenço, 2006) from the Dominican amber (Scorpiones: Buthidae). *Euscorpius*, 83: 1–5.
- Lourenço, W. R. 2012. Further considerations on scorpions found in Baltic amber, with a description of a new species (Scorpiones: Buthidae). *Euscorpius*, 146: 1–7.
- Lourenço, W. R. 2013. A new species of *Tityus* C. L. Koch, 1836 (Scorpiones: Buthidae) from Dominican amber. *Euscorpius*, 156: 1–5.
- Lourenço, W. R. 2013. A new species of scorpion from Chiapas amber, Mexico (Scorpiones: Buthidae). *Revista Ibérica de Aracnología*, 24: 59–63.
- Lourenço, W.R. & Gall, J.-C. 2004. Fossil scorpions from the Buntsandstein (Early Triassic) of France. *Comptes Rendus Palevol*, 3: 369–378.
- Lourenço, W. R. & Henderickx, H. 2012. Another new sub-fossil species of scorpion of the genus *Palaeogroosphus* Lourenço, 2000 from Malagasy copal (Scorpiones: Buthidae). *Euscorpius*, 137: 1–4.
- Lourenço, W. R. & Weitschat, W. 1996. More than 120 years after its description, the enigmatic status of the genus of the Baltic amber scorpion “*Tityus eogenus*” Menge, 1869 can finally be clarified. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 79: 183–188.
- Lourenço, W. R. & Weitschat, W. 2000. New fossil scorpions from the Baltic amber – implications for Cenozoic biodiversity. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 84: 247–260.
- Lourenço, W. R. & Weitschat, W. 2001. Description of another fossil scorpion from Baltic amber with considerations on the evolutionary levels of Cenozoic Buthoidea. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 85: 277–283.
- Lourenço, W. R. & Weitschat, W. 2005a. A new genus and species of fossil scorpion from a different kind of Baltic amber (Scorpiones, Buthidae). – *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 89, 183–188.
- Lourenço, W. R. & Weitschat, W. 2005b. First sub-fossil scorpion of genus *Chactas* Gervais from Colombian copal (Scorpiones, Chactidae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 89: 179–182.
- Lourenço, W. R. & Weitschat, W. 2009. A new species of *Palaeoananteris* Lourenço & Weitschat, 2001, fossil scorpion from Ukrainian amber (Scorpiones, Buthidae). *Boletín Sociedad Entomológica Aragonesa*, 45: 231–235.

- Lourenço, W. R., Henderickx, H. & Weitschat, W. 2005. A new genus and species of fossil scorpion from Baltic amber (Scorpiones, Buthidae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 89: 159–166.
- Lucas, H. 1835. Sur une monographie du genre Thélyphone. *Magasin de Zoologie*, 5: Classe VIII, pls. 8–10.
- Lucas, H. 1846. Histoire naturelle des Animaux articulés. *In Exploration scientifique de l'Algérie pendant les années 1840, 1841, 1842, publiée par ordre du Gouvernement et avec le concours d'une commission académique. Sciences physiques, Zoologie, 5 tomes, Paris, 1846–1850*. Vol. 1: 89–271.
- Luxton, M. 1985. Cryptostigmata (Arachnida: Acari): a concise review. *Fauna of New Zealand*, 7: 1–112.
- Luxton, M. 1988. A new mite superfamily (Acari: Cryptostigmata). *Zoological Journal of the Linnean Society*, 93: 71–91.
- Lyubarsky, G. Y. & Perkovsky, E. E. 2012. The first Eocene species of the genus *Cryptophagus* (Coleoptera, Clavicornia, Cryptophagidae). *Vestnik zoologii*, 46: 36–40.
- MacLeay, W. S. 1839. On some new forms of Arachnida. *Annals and Magazine of Natural History*, 2: 1–14.
- Magowski, W. Ł. 1994. Discovery of the first representative of the mite subcohort Heterostigmata (Arachnida: Acari) in the Mesozoic Siberian amber. *Acarologia*, 35: 229–241.
- Magowski, W. Ł. 1995. Fossil heterostigmatid mites in amber – 85 million year-old arthropod mite Relationships 53–58. *In Kropczynska, D., Boczek, J. & Tomczyk, A. (eds) The Acari: Physiological and Ecological Aspects of Acari – Host Relationships* Dabor, Warsaw, 698 pp.
- Malz, H. & Poschmann, M. 1993. Erste Süßwasser-Limuliden (Arthropoda, Chelicerata) aus dem Rotliegenden der Saar-Nahe-Senke. *Osnabrücker naturwissenschaftliche Mitteilungen*, 19: 21–24.
- Mahnert, V. 1979. Pseudoskopione (Arachnida) aus dem Amazonas-Gebiet (Brasilien). *Revue suisse de Zoologie*, 86: 719–810.
- Mahunka, S. 1986. A survey of the family Carabodidae C. L. Koch, 1836 (Acari: Oribatida). *Acta Zoologica Hungarica*, 32: 73–135.
- Mahunka, S. 1990. A survey of the superfamily Euphthiracaroidea Jacot, 1930 (Acari: Oribatida). *Folia Entomologica Hungarica*, 51: 37–80.
- Marshall, D. J., Lamsdell, J. C., Shpinev, E. & Braddy, S. J. 2014. A diverse chasmataspidid (Arthropoda: Chelicerata) fauna from the Early Devonian (Lochkovian) of Siberia. *Palaeontology*, 57, 631–655.
- Martens, J. 1976. Genitalmorphologie, System und Phylogenie der Weberknechte (Arachnida: Opiliones). *Entomologica Germanica*, 3: 51–68.
- Martens, J. 1988. Fissiphalliidae, a new family of South American laniatorean harvestmen (Arachnida: Opiliones). *Zeitschrift für zoologische Systematik und Evolutionsforschung*, 26: 114–127.
- Martin, W. 1809. *Petrificata Derbiensis* 1, [Wigan](#), xx pp.
- Marusik, Y. M. & Penney, D. 2004. A survey of Baltic amber Theridiidae (Araneae) inclusions, with descriptions of six new species. *In Logunov, D. V. & Penney, D (eds). European Arachnology 2003 (Proceedings of the*

- 21st European Colloquium of Arachnology, St.-Petersburg, 4–9 August 2003). *Arthropoda Selecta*, Special Issue No. 1: 201–208.
- Marx, G. 1888. On a new and interesting spider. *Entomologica Americana*, 4: 160–162.
- Marx, G. 1890a. Arachnida. In Howard, L. O. (ed.) Scientific results of the explorations by the U. S. Fish Commission Steamer Albatross. No. V. – Annotated catalogue of the insects collected in 1887–'88. – *Proceedings of the United States National Museum*, 12: 207–211.
- Marx, G. 1890b. Catalogue of the described Araneae of temperate North America. *Proceedings of the United States National Museum*, 12: 497–594.
- Matthew, G. F. 1888. On some remarkable organisms of the Silurian and Devonian rocks in Southern New Brunswick. *Transactions of the Royal Society of Canada*, 1888: 49–61.
- Matthew, G. F. 1895. Organic remains of the Little River Group, No. IV. *Transactions of the Royal Society of Canada*, 2nd Ser., 1: 273–279.
- McAlpine, J. F. & Martin, J. E. H. 1969. Canadian amber – a paleontological treasure chest. *Canadian Entomologist*, 101: 819–838.
- McCook, H. C. 1888. A new fossil spider, *Eoatypus woodwardii*. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1888: 200–202.
- Meek, F. B. 1867. Notes on a new genus of fossil Crustacea. *Geological Magazine, Decade 4*, **xx**: 320–321.
- Meek, F. B. & Worthen, A. H. 1865. Notice of some new types of organic remains from the Coal Measures of Illinois. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 17: 41–45.
- Meek, F.B & Worthen, A.H. 1868a. Preliminary notice of a scorpion, a *Eurypterus*? and other fossils from the Coal Measures of Illinois and Iowa. *American Journal of Science and Arts, series 2*, 45: 25.
- Meek, F.B. & Worthen, A.H. 1868b. Palaeontology of Illinois. In *Geological Survey of Illinois*, 3: 289–565.
- Melander, A. L. 1903. Some additions to the Carboniferous terrestrial fauna of Illinois. *Journal of Geology*, 11: 178–198.
- Melendez, B. 1971. Un novel Euryptéride du Westphalien des Asturies (NW Espagne). In Krefeld (ed.) *Septième Congrès de Stratigraphie et de Géologie du Carbonifère*, 3: 415–417.
- Mello-Leitão, C. F. de 1932. Notas sobre as Micratheneas do Brasil. *Anais do Academia Brasileira dos Ciências*, 4: 73–97.
- Mello-Leitão, C. F. de 1937. Distribution et Phylogénie des Faucheurs Sud-Américains. *Proceedings of the 12th International Congress of Zoology, Lisbon*, 2(5): 1217–1228.
- Mello-Leitão, C. F. de 1940. Arañas de las islas Juan Fernandez, recogidas por el Señor R. Wagenknecht. *Revista Chilena de Historia Natural*, 44: 236–239.
- Menge, A. 1854. Footnotes in Koch, C. L. & Berendt, G. C. Die im Bernstein befindlichen Myriapoden, Arachniden und Apteran der Vorwelt. In Berendt, G. C. *Die in Bernstein befindlichen organischen Reste der Vorwelt gesammelt in verbindung mit mehreren bearbeitetet und herausgegeben* 1. Berlin, Nicolai, 124 pp.

- Menge, A. 1855. Ueber die Scheerenspinnen, Chernetidae. *Neueste Schriften der Naturforschenden Gesellschaft*, 5: 1–43.
- Menge, A. 1856. Lebenszeichen vorweltlicher, im Bernstein eingeschlossener Thiere. *Programm der Petrischule zu Danzig*, 8: 32 pp.
- Menge, A. 1866. Preussische Spinnen. Erste Abtheilung. *Schriften der Naturforschenden Gesellschaft in Danzig (Neue Folge)*, 2: 1–152.
- Menge, A. 1868. Preussische Spinnen. II. Abtheilung. *Schriften der Naturforschenden Gesellschaft in Danzig (Neue Folge)*, 2: 153–218.
- Menge, A. 1869. Ueber einen Scorpion und zwei Spinnen im Bernstein. *Schriften der Naturforschenden Gesellschaft in Danzig (Neue Folge)*, 2: 1–9.
- Mesquita, M. V. 1996. *Cretaraneus matensnetoi* n.sp. (Araneoidea) da Formação Santana, Cretáceo Inferior da Bacia do Araripe. *Revista Universidade Guarulhos, Série Geociências*, 1(3): 24–31.
- Miller, S. A. 1874. Notes and descriptions of Cincinnati Group fossils. *Cincinnati Quarterly Journal of Science*, 1: 343–351.
- Miller, S. A. & Gurley, W. F. E. 1896. New species of Echinodermata and a new crustacean from the Palaeozoic rocks. *Illinois State Museum Natural History Bulletin*, 10: 1–91.
- Millot, J. 1947. Une araignée malgache énigmatique, *Gallieniella mygaloides* n. g., n. sp. *Bulletin du Muséum National d'Histoire Naturelle, 2^e Série*, 19: 158–160.
- Millot, J. 1948. Faits nouveaux concernant les Archaea [Aranéides]. *Mémoires de l'Institut Scientifique de Madagascar*, 1(A1): 3–14.
- Moberg, J. C. 1892. Om en nyupptäckt fauna i block af kambrisk sandsten, insamlade af Dr N.O. Holst. *Geologiska Föreningens i Stockholm Förhandlingar*, 14: 103–120.
- Moore, J. I. 1923. A review of the present knowledge of fossil scorpions, with the description of a new species from the Pottsville Formation of Clay County, Indiana. *Proceedings of the Indiana Academy of Science*, 38: 125–134.
- Moore, R. A., McKenzie, S. C. & Lieberman, B. S. 2007. A Carboniferous synziphosurine (Xiphosura) from the Bear Gulch Limestone, Montana, USA. *Palaeontology*, 50: 1013–1019.
- Moore, R. A., Briggs, D. E. G., Braddy, S. J. & Shultz, J. W. 2011. Synziphosurines (Xiphosura: Chelicerata) from the Silurian of Iowa. *Journal of Paleontology*, 85: 83–91.
- Moore, R. A., McKenzie, S. C., Braddy, S. J., Anderson, L. I., Mikulic, D. G. & Kluessendorf, J. 2005. A new synziphosurine (Chelicerata: Xiphosura) from the Late Llandovery (Silurian) Waukesha Lagerstätte, Wisconsin, USA. *Journal of Paleontology*, 79: 242–250.
- Moran, R. J. 1986. The Sternodidae (Araneae, Araneomorpha), a new family of spiders from eastern Australia. *Bulletin of the British Arachnological Society*, 7: 87–96.

- Moraza, M. L. & Lindquist, E. E. 1999. Coprozerconidae, a new family of zeronoid mites from North America (Acari: Mesostigmata: Zerconoidea). *Acarologia*, 39: 291–313.
- Müller, O. F. 1785. *Entomastraca, seu, Insecta testacea quae in aquis Daniae et Norvegiae reperit, descriptis et iconibus illustravit*. Hauniae, Thiele, xx pp.
- Müller, A. H. 1957. Ein Arachnidenrest (*Brachylycosa ? manebachensis* n. sp.) aus dem Unteren Rotliegenden (Manebacher Schichten) von Thüringen. *Geologie*, 6: 95–98.
- Münster, G. Graf zu 1839. Die Rhyncholiten des Muschelkalks mit ihrem Fortsätzen. In Münster, G. Graf zu (ed.) *Beiträge zur Petrefacten-Kunde* 1: 48–51.
- Münster, G. Graf zu 1840. Über die fossilen Arten *Limulus* in den lithographischen Schiefern von Bayern. In Münster, G. Graf zu (ed.) *Beiträge zur Petrefacten-Kunde* 3: 26–27.
- Murdoch, J.B. 1893. Proceedings for Session 1890–91. *Transactions of the Geological Society of Glasgow*, 9: 414–422.
- Murray, A. 1877. *Economic Entomology, Aptera*. South Kensington Museum Handbooks, 433 pp.
- Nalepa, A. 1898. Eriophyidae (Phytoptidae). In *Das Tierreich. Eine Zusammenstellung und Kennzeichnung der rezenten Tierformen*. 4. Lieferung. Acarina. Deutsche Zoologische Gesellschaft, 4: 72 pp.
- Nicolet, H. 1855. Histoire naturelle des Acariens qui se trouvent aux environs de Paris. *Archives de Museum Nationale d'Histoire Naturelle de Paris*, 7: 381–482.
- Nieszkowski, J. 1859. Zusätze zur Monographie der Trilobiten der Ostseeprovinzen, nebst der Beschreibung einiger neuen obersilurischen Crustaceen. *Archiv für die Naturkunde Liv.-, Ehst.- und Kurlands* (Ser. 1), 1: 345–384.
- Nindel, F. 1955. Die tierischen Reste aus dem Karbon von Karl-Marx-Stadt und Hainichen i.S. *Geologie*, 4: 673–694.
- Nishikawa, Y. 1974. [Amber spiders from Mizunami, Japan.] *Bulletin of the Mizunami Fossil Museum*, 1: 401–406. [in Japanese with English summary]
- Norton, R. A. 2006. First record of *Collohmannia* (*C. schusteri* n. sp.) and *Hermannia* (*H. sellnicki* n. sp.) from Baltic amber, with notes on Sellnick's genera of fossil oribatid mites (Acari: Oribatida). *Acarologia*, 46: 111–125.
- Norton R.A. & Metz, L. 1980. Nehyphochthoniidae (Acari: Oribatei), a new family from the southeastern United States. *Annals of the Entomological Society of America*, 73: 54–62.
- Norton, R. A., Bonamo, P. N., Grierson, J. D. & Shear, W. A. 1988. Oribatid mite fossils from a terrestrial Devonian deposit near Gilboa, New York. *Journal of Paleontology*, 62: 259–269.
- Novojilov, N. J. 1959. Mérostomes du Dévonien inférieur et moyen de Sibérie. *Annales de la Société Géologique du Nord*, 78: 241–258.
- Novojilov, N. & Størmer, L. 1963. A new scorpion from the Upper Carboniferous of Siberia. *Norsk Geologisk Tidsskrift* 43: 83–87.

- O'Connell, M. 1916. The habitat of the Eurypterida. *Bulletin of the Buffalo Society of Natural Sciences*, 11: 1–278.
- Olivier, P. A. S. & Theron, P. D.. 2000. Pentapalpidae, a new family of eupodoid mites (Prostigmata:Eupodoidea) from South Africa. *Acarologia*, 40: 385–392.
- Ono, H. 1981. First record of a crab spider (Thomisidae) from Dominican amber (amber collection Stuttgart : Arachnida, Araneae). *Stuttgarter Beiträge zur Naturkunde (B)*, 73: 1–13.
- Opluštil, S. 1985. New findings of Arachnida from the Bohemian Upper Carboniferous. *Věstník Ústředního ústavu geologického*, 60: 35–42.
- Opluštil, S. 1986. *Promygale janae* sp. n., the new anthracomartid (Arachnida) from the Upper Carboniferous of central Bohemia. *Věstník Ústředního ústavu geologického*, 61: 287–292.
- Oppenheim, P. 1887–1888. Die Insectenwelt des lithographischen Schiefers in Bayern. *Palaeontographica*, 34: 215–247.
- Orr, P. J., Siveter, D. J., Briggs, D. E. G., Siveter, D. J. & Sutton, M. D. 2000. A new arthropod from the Silurian Konservat-Lagerstätte of Herefordshire, UK. *Proceedings of the Royal Society B*, 267: 1497–1504.
- Oudemans, A. C. 1902. Classificatie der Acari. *Tijdschrift voor Entomologie*, 45: 50–64.
- Oudemans, A. C. 1909. Über die bis jetzt genauer bekannten Thrombidium-larven und über eine neue Klassifikation der Prostigmata. *Tijdschrift voor Entomologie*, 52: 19–61.
- Oudemans, A. C. 1916. Acarologische Aanteekeningen LX. *Entomologische berichten*, 4: 308–316.
- Oudemans, A. C. 1923. Studie over de sedert 1977 ontworpen system der Acari; nieuwe classificatie; phylogenerische beschowingen. *Tijdschrift voor Entomologie*, 66: 49–85.
- Özdikmen, H. 2007. Nomenclatural changes for seven preoccupied spider genera (Arachnida: Araneae). *Munis Entomology & Zoology*, 2: 137–142.
- Packard, A. S. 1885. Types of Carboniferous Xiphosura new to North America. *American Naturalist*, 1885: 291–294.
- Packard, A. S. 1886. On the Carboniferous xiphosurous fauna of North America. *Memoirs of the National Academy of Sciences*, 3: 143–157.
- Page, D. 1856. *Advanced textbook of geology*. William Blackwood and Sons, Edinburgh, 326 pp.
- Page, D. 1859. *Advanced textbook of geology*, 2nd edn. William Blackwood and Sons, London, xx pp.
- Palmer, A. R. 1957. Miocene arthropods from the Mojave Desert California. *Geological Survey Professional Paper*, 294-G: 237–280.
- Pampaloni, L. 1902. I resti organici nel disodile di Melilli in Sicilia. *Palaeontographica Italica*, 8: 121–130.
- Patrick, R. R. 1989. A new phalangiotarbid (Arachnida) from the McLeansboro Group (Pennsylvanian) of Indiana. *Journal of Paleontology*, 63: 327–331.
- Peach, R. N. 1882. Further researches among Crustacea and Arachnida. *Transactions of the Royal Society of Edinburgh*, 30: 511–529.

- Peach, R. N. 1883. A new species of fossil scorpions from the Carboniferous rocks of Scotland and the English borders, with a review of the genera *Eoscorpius* and *Mazonia* of Messrs. Meek and Worthen. *Transactions of the Royal Society of Edinburgh*, 30: 397–412.
- Peach, R. N. 1888. On a new eurypterid from the Upper Coal-measures of Radstock, Somersetshire. *Proceedings of the Royal Physical Society, Edinburgh*, 9: 438–445.
- Peckham, G. W. & Peckham, E. G. 1892. Ant-like spiders of the Family Attidae. *Occasional Papers of the Natural History Society of Wisconsin*, 2(1): 1–83.
- Peckham, G. W. & Wheeler, W. H. 1889. Spiders of the subfamily Lyssomanae. *Transactions of the Wisconsin Academy of Science, Arts and Letters*, 7: 222–256.
- Penney, D. 2000. Miocene spiders in Dominican amber (Oonopidae, Mysmenidae). *Palaeontology*, 43: 343–357.
- Penney, D. 2001. Advances in the taxonomy of spiders in Miocene amber from the Dominican Republic (Arthropoda: Araneae). *Palaeontology*, 44: 987–1009.
- Penney, D. 2002. Spiders in Upper Cretaceous amber from New Jersey (Arthropoda: Araneae). *Palaeontology*, 45: 709–724.
- Penney, D. 2003a. *Afrarchaea grimaldii*, a new species of Archaeidae (Araneae) in Cretaceous Burmese amber. *The Journal of Arachnology*, 31: 122–130.
- Penney, D. 2003b. A new deinopid spider from Cretaceous Lebanese amber. *Acta Palaeontologica Polonica*, 48: 569–574.
- Penney, D. 2004a. New spiders in Upper Cretaceous amber from New Jersey in the American Museum of Natural History (Arthropoda: Araneae). *Palaeontology*, 47: 367–375.
- Penney, D. 2004b. Cretaceous Canadian amber spider and the palpimanoidean nature of lagonomegopids. *Acta Palaeontologica Polonica*, 49: 579–584.
- Penney, D. 2004c. A new genus and species of Pisauridae (Araneae) in Cretaceous Burmese amber. *Journal of Systematic Palaeontology*, 2: 141–145.
- Penney, D. 2005a. First fossil Filistatidae: a new species of *Misionella* in Miocene amber from the Dominican republic. *The Journal of Arachnology*, 33: 93–100.
- Penney, D. 2005b. The fossil spider family Lagonomegopidae in Cretaceous ambers with descriptions of a new genus and species from Myanmar. *The Journal of Arachnology*, 33: 439–444.
- Penney, D. 2005c. First Caribbean *Floricomus* (Araneae: Linyphiidae), a new fossil species in Miocene Dominican Republic amber. A new synonymy fro the extant North American fauna. *Geologica Acta*, 3: 59–64.
- Penney, D. 2005d. An annotated systematic catalogue, including synonymies and transfers, of Miocene Dominican Republic amber spiders described up until 2005. *Revista Ibérica de Aracnología*, 12: 25–52.
- Penney, D. 2006a. Fossil oonopid spiders in Cretaceous ambers from Canada and Myanmar. *Palaeontology*, 49: 229–235.

- Penney, D. 2006b. The oldest lagonomegopid spider, a new species in Lower Cretaceous amber from Álava, Spain. *Geologica Acta*, 4: 377–382.
- Penney, D. 2007a. The oldest fossil pholcid and selenopid spiders (Araneae) in lowermost Eocene amber from the Paris Basin France. *The Journal of Arachnology*, 34: 592–598.
- Penney, D. 2007b. A new fossil oonopid spider in lowermost Eocene amber from the Paris Basin, with comments on the fossil spider assemblage. *African Invertebrates*, 48: 71–75.
- Penney, D. 2009. A new spider family record for Hispaniola – a new species of *Plectreurus* (Araneae: Plectreuridae) in Miocene Dominican amber. *Zootaxa*, 2144: 65–68.
- Penney, D. 2011. Grandoculidae: a new fossil spider family from the Upper Cretaceous of Canada. *Bulletin of the British arachnological Society*, 15: 179–180.
- Penney, D. 2014. A fossil ray spider (Araneae: Theridiosomatidae) in Cretaceous amber from Vendée, France. *Paleontological Contributions*, 10B: 1–8.
- Penney, D. & Ortuño, V. N. 2006. Oldest true orb-weaving spider (Araneae: Araneidae). *Biology Letters*, 2: 447–450.
- Penney, D. & Selden, P. A. 2002. The oldest linyphiid spider in Lower Cretaceous Lebanese amber /Araneae, Linyphiidae, Linyphiinae). *The Journal of Arachnology*, 30: 487–493.
- Penney, D. & Selden, P. A. 2006. First fossil Huttoniidae (Arthropoda: Chelicerata: Araneae) in late Cretaceous Canadian amber. *Cretaceous Research*, 27: 442–446.
- Penney, D., Dierick, M., Cnudde, V., Masschaele, B., Vlassenbroeck, J., Hoorebeke, L. van & Jacobs, P. 2007. First fossil Micropholcommatidae (Araneae), imaged in Eocene Paris amber using X-Ray Computed Tomography. *Zootaxa*, 1623: 47–53.
- Penney, D., Green, D. I., Tichner, S. B., Titchner, B. G., Brown, T. A., Preziosi, R. F. 2012c. An unusual palaeobiocoenosis of subfossil spiders in Colombian copal. *Bulletin of the British Arachnological Society*, 15: 241–244.
- Penney, D., McNeil, A., Green D. I., Bradley, R., Marusik, Y. M., Withers, P. J. & Preziosi, R. F. 2011. A new species of anapid spider (Araneae: Araneoidea, Anapidae) in Eocene Baltic amber, imaged using phase contrast X-ray computed micro-tomography. *Zootaxa*, 2742: 60–66.
- Penney, D., McNeil, A., Green D. I., Bradley, R., Withers, P. J. & Preziosi, R. F. 2012a. The oldest fossil pirate spider (Araneae: Mimetidae), in uppermost Eocene Indian amber, imaged using X-ray computed tomography. *Bulletin of the British Arachnological Society*, 15: 299–302.
- Penney, D., Green D. I., McNeil, A., Bradley, R., Marusik, Y. M., Withers, P. J. & Preziosi, R. F. 2012b. A new species of *Craspedisia* (Araneae: Theridiidae) in Miocene Dominican amber, imaged using X-ray computed tomography. *Paleontological Journal* 46: 583–588. [Translation of Russian original]
- Pérez, d'A.V. 1988. Un oribatido del Eoceno (Terciario). Primar acaro fosil de Chile (Arachnida: Acari: Oribatida). *Revista Chilena de Entomología*, 16: 23–24.

- Pérez González, A.. & Kury A. 2007. Kimulidae. In Pinto da Rocha, R., Machado, G. & Giribet, G. (eds). *Harvestmen. The Biology of Opiliones*. Harvard University Press, Cambridge MA, pp. 207–209.
- Perkovsky, E. E., Rasnitsyn, A. P., Vlaskin, A. P., Taraschuk, M. V. 2007. A comparative analysis of the Baltic and Rovno amber arthropod faunas: representative samples. *African Invertebrates*, 48:229–245
- Perry, M. L. 1995. Preliminary description of a new fossil scorpion from the middle Eocene Green River Formation, Rio Blanco County, Colorado. In Dayvault, R. D. & Averett, W. R. (eds). *The Green River Formation in Piceance Creek and Estern Unita Basins Field Trip*. Grand Junction Geological Society, Grand Junction Colorado, pp. 131–133.
- Peters, W. 1861. (Ueber eine neue Eintheilung der Skorpione und ueber die von ihm in Mossambique gesammelten Arten von Skorpionen). *Monatsberichte der Königlichen Preussischen Akademie der Wissenschaft zu Berlin*, 1861: 507–516.
- Petrunkewitch, A. I. 1913. A monograph of the terrestrial Palaeozoic Arachnida of North America. *Transactions of the Connecticut Academy of Arts and Sciences*, 18: 1–137.
- Petrunkewitch, A. I. 1922. Tertiary spiders and opilionids of North America. *Transactions of the Connecticut Academy of Arts and Sciences*, 25: 211–279.
- Petrunkewitch, A. I. 1923. On families of spiders. *Annals of the New York Academy of Science*, 29: 145–180.
- Petrunkewitch, A. I. 1928. Systema Aranearium. *Transactions of the Connecticut Academy of Arts and Sciences*, 29: 1–270.
- Petrunkewitch, A. I. 1942. A study of amber spiders. *Transactions of the Connecticut Academy of Arts and Sciences*, 34: 119–464.
- Petrunkewitch, A. I. 1945a. Palaeozoic Arachnida. An inquiry into their evolutionary trends. *Scientific Papers, Illinois State Museum*, 3(2): 1–76.
- Petrunkewitch, A. I. 1945b. *Calcitro fisheri*. A new fossil arachnid. *American Journal of Science*, 243: 320–329.
- Petrunkewitch, A. I. 1946. Fossil spiders in the collection of the American Museum of Natural History. *American Museum Novitates*, 1328: 1–36.
- Petrunkewitch, A. I. 1949. A study of Palaeozoic Arachnida. *Transactions of the Connecticut Academy of Arts and Sciences*, 37: 69–315.
- Petrunkewitch, A. I. 1950. Baltic amber spiders in the Museum of Comparative Zoology. *Bulletin of the Museum of Comparative Zoology*, 103: 257–337.
- Petrunkewitch, A. I. 1953. Palaeozoic and Mesozoic Arachnida of Europe. *Memoirs of the Geological Society of America*, 53: 1–128.
- Petrunkewitch, A. I. 1955a. Arachnida. 42–162. In Moore, R. C. (ed.) *Treatise on invertebrate paleontology, Part P, Arthropoda 2*. Geological Society of America, Boulder, and University of Kansas Press, Lawrence, xvii + 181 pp.

- Petrunkевич, A. I. 1955b. *Trigonotarbus arnoldi*, a new species of fossil arachnid from Southern France. *Journal of Paleontology*, 29: 475–477.
- Petrunkевич, A. I. 1958. Amber spiders in European collections. *Transactions of the Connecticut Academy of Arts and Sciences*, 41: 97–400.
- Petrunkевич, A. I. 1963. Chiapas amber spiders. *University of California Publications in Entomology*, 31: 1–40.
- Petrunkевич, A. I. 1971. Chiapas amber spiders, II. *University of California Publications in Entomology*, 63: 1–44.
- Pickett, J. W. 1984. A new freshwater limuloid from the middle Triassic of New South Wales. *Palaeontology*, 27: 609–621.
- Pickett, J. W. 1993. A Late Devonian xiphosuran from near Parkes, New South Wales. *Memoirs of the Association of Australian Palaeontologists*, 15: 279–287.
- Pickford, M. 2000. Fossil spider's webs from the Namib Desert and the antiquity of *Seothyra* (Araneae, Eresidae). *Annales de Paléontologie*, 86: 147–155.
- Pictet, F. J. 1846. *Traité élémentaire de paléontologie*. Vol. 4. Paris, 458 pp.
- Pierce, W. D. 1945. A fossil whiptail scorpion from Cabrillo Beach. *Bulletin of the Southern California Academy of Sciences*, 44: 7–8.
- Pierce, W. D. 1950. Fossil arthropods from onyx-marble. *Bulletin of the Southern Californian Academy of Sciences*, 49: 101–104.
- Pierce, W. D. 1951. Fossil arthropods from onyx-marble. *Bulletin of the Southern Californian Academy of Sciences*, 50: 34–49.
- Pinto, I. D. & Hünicken, M. A. 1980. *Gondwanarachne* a new genus of the order Trigonotarbida (Arachnida) from Argentina. *Boletín de la Academia Nacional de Ciencias Córdoba*, 53: 307–315.
- Pirozhnikov, L. P. 1957. [Remains of Gigantostraca from the series of Matakara (Devonian of North Minusinsk Depression).] *Annuaire de la Société paléontologique de Russie*, 16: 207–213. [in Russian]
- Platnick, N. I. 1977. The hypochiloid spiders: a cladistic analysis, with notes on the Atypoidea (Arachnida, Araneae). *American Museum Novitates*, 2627, 1–23.
- Platnick, N. I. 1989. *Advances in Spider Taxonomy 1981-1987: A Supplement to Brignoli's A Catalogue of the Araneae described between 1940 and 1981*. Manchester University Press, 673 pp.
- Pocock, R. I. 1892. *Liphistius* and its bearing upon the classification of spiders. *Annals and Magazine of Natural History*, series 6, 10: 306–314.
- Pocock, R. I. 1893. Notes on the classification of scorpions, followed by some observations on synonymy, with descriptions of new genera and species. *Annals and Magazine of Natural History*, series 6, 12: 303–330.
- Pocock, R. I. 1895. Description of two new spiders obtained by Messrs J. J. Quelch and F. MacConnel on the summit of Mount Roraima, in Demerara; with a note upon the systematic position of the genus *Desis*. *Annals and Magazine of Natural History*, series 6, 16: 139–143.

- Pocock, R. I. 1897. On the genera and species of tropical African Arachnida of the order Solifugae, with notes upon the taxonomy and habits of the group. *Annals and Magazine of Natural History, series 6*, 20: 249–272.
- Pocock, R. I. 1898. The Arachnida from the Province of Natal, South Africa, contained in the collection of the British Museum. *Annals and Magazine of Natural History, series 7*, 2: 197–226.
- Pocock, R. I. 1901. The Scottish Silurian scorpions. *Quarterly Journal of Microscopical Science*, (2) 44: 291–311.
- Pocock, R. I. 1902. *Eophryinus* and allied Carboniferous Arachnida. *Geological Magazine, Decade 4*, 9: 439–448, 487–493.
- Pocock, R. I. 1903a. A new Carboniferous arachnid. *Geological Magazine, Decade 4*, 10: 247–251.
- Pocock, R. I. 1903b. Further remarks upon the Carboniferous arachnid *Anthracosiro*, with the description of a second species of the genus. *Geological Magazine, Decade 4*, 10: 405–408.
- Pocock, R. I. 1903c. On the geographical distribution of spiders of the order Mygalomorphae. *Proceedings of the Zoological Society of London*, 1903: 340–368.
- Pocock, R. I. 1911. A monograph of the terrestrial Carboniferous Arachnida of Great Britain. *Monographs of the Palaeontographical Society*, 64: 1–84.
- Pohlman, J. 1882. Additional Notes on the Fauna of the Water-Lime Group near Buffalo. *Bulletin of the Buffalo Society of Natural Sciences*, 4(2): 41–47.
- Poinar Jr., J. O. 1985. Fossil evidence of insect parasitism by mites. *International Journal of Acarology*, 11: 37–38.
- Poinar Jr., G.O. 1988. Hair in Dominican amber: evidence for Tertiary land mammals in the Antilles. *Experientia*, 44: 88–89.
- Poinar Jr., G. O. 1995. First fossil soft tick, *Ornithodoros antiquus* n. sp. (Acari: Argasidae) in Dominican amber with evidence of their mammalian host. *Experimentia Basel*, 51: 584–587.
- Poinar Jr., G. [O.] 2008. *Palaeosiro burmanicum* n. gen., n. sp., a fossil Cyphophthalmi (Arachnida: Opiliones: Sironidae) in Early Cretaceous Burmese amber. In Makarov, S. E. & Dimitrijević, R. N. (eds) *Advances in Arachnology and Developmental Biology. Papers dedicated to Prof. Dr. Božidar Ćurčić*. Inst. Zool., Belgrade; BAS, Sofia; Fac. Life Sci., Vienna; SASA, Belgrade & UNESCO MAB Serbia. Vienna — Belgrade — Sofia, Monographs, 12: 267–274 .
- Poinar Jr., G. O. & Brown, A. E. 2003. A new genus of hard ticks in Cretaceous Burmese amber (Acari: Ixodida: Ixodidae). *Systematic Parasitology*, 54: 199–205.
- Poinar Jr., G. O. & Brown, A. E. 2004. A new whip spider (Arachnida: Amblypygi), *Phrynx mexicana*, is described from Mexican amber. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 3: 1881–1885.
- Poinar Jr., G. O. & Buckley, R. 2008. *Compluriscutula vetulum* (Acari: Ixodida: Ixodidae), a new genus and species of hard tick from Lower Cretaceous Burmese amber. *Proceedings of the Entomological Society of Washington*, 110: 445–450.

- Poinar Jr., G. O. & Buckley, R. 2012. Predatory behaviour of the social orb-weaver spider, *Geratonephila burmanica* n. gen., n. sp. (Araneae: Nephilidae) with its wasp prey, *Cascoscelio incassus* n. gen., n. sp. (Hymenoptera: Platygastridae) in Early Cretaceous Burmese amber. *Historical Biology*, 24: 519–525.
- Poinar Jr., G. O. & Santiago-Blay, J. A. 1989. A fossil solpugid, *Haplodontus proterus*, new genus, new species (Arachnida: Solpugida) from Dominican amber. *Journal of the New York Entomological Society*, 97: 125–132.
- Ponomarenko, A. G. 1985. King crabs and eurypterids from the Permian and Mesozoic of the USSR. *Paleontological Journal*, 19: 100–104. [Translation of *Paleontologiceskij Žurnal*, 1985: 115–117.]
- Poschmann, M. 2009. Ein fossiler Skorpion aus der Oberkarbon (Westfalium C) des Saar-Nahe-Beckens (SW Deutschland). *Mitteilungen der Pollicchia*, 94: 5–10.
- Poschmann, M. & Dunlop, J. A. 2006. A new sea spider (Arthropoda: Pycnogonida) with a flagelliform telson from the Lower Devonian Hunsrück Slate, Germany. *Palaeontology*, 49: 983–989.
- Poschmann, M. & Dunlop, J. A. 2010. Trigonotarbid arachnids from the Lower Devonian (Lower Emsian) of Alken an der Mosel (Rhineland-Palatinate, SW Germany). *Paläontologische Zeitschrift*, 84: 467–484.
- Poschmann, M. & Dunlop, J. A. 2011. Trigonotarbid arachnids from the Lower Devonian (Siegenian) of Bürdenbach (Lahrbach Valley, Westerwald area, Rhenish Slate Mountains, Germany). *Paläontologische Zeitschrift*, 85: 433–447.
- Poschmann, M. & Dunlop, J. A. 2012. Reassessing *Devonotarbus*, a phalangiotarbid arachnid from the Lower Devonian (Siegenian and Emsian) of the Rheinisches Schiefergebirge (SW Germany). *Paläontologisches Zeitschrift*, 86: 377–387.
- Poschmann, M. & Tetlie, O. E. 2004. On the Emsian (Early Devonian) arthropods of the Rhenish Slate Mountains: 4. The eurypterids *Alkenopterus* and *Vinetopterus* n. gen. (Arthropoda: Chelicerata). *Senckenbergiana lethaea*, 84: 173–193.
- Poschmann, M., Anderson, L. I. & Dunlop, J. A. 2005. Chelicerate arthropods, including the oldest phalangiotarbid arachnid, from the Early Devonian (Siegenian) of the Rhenish Massif, Germany. *Journal of Paleontology*, 79: 110–124.
- Poschmann, M., Dunlop, J. A., Kamenz, C. & Scholtz, G. 2008. The Lower Devonian scorpion *Waeringoscorpio* and the respiratory nature of ist filamentous structures, with a description of a new species from the Westerwald area, Germany. *Paläontologische Zeitschrift*, 82: 418–436.
- Prach, F. K. 1860. Život Pavouků pravých či předoučích (Araneae). *Živa*, 8: 80–93.
- Presl, J. S. 1822. Additamenta ad faunam protogaeam, sistens descriptions aliquot animalium in succino inclusorum. In Presl, J. S. & Presl, C. B. (eds). *Deliciae Pragenses Historiam Naturalem Spectantes. Tome I. Calvae, Pragae*, viii + 244 pp.
- Prestwich, J. 1840. Memoir on the geology of Coalbrook Dale. *Transactions of the Geological Society of London* 5: 413–495.

- Příbyl, A. 1952. On the genus *Adelophthalmus* Jordan and Meyer, 1854 (Euryperida) and its representatives in the Upper Carboniferous of Czechoslovakia. *Bulletin International de l'Académie tchéque des Sciences*, 53: 63–70.
- Příbyl, A. 1958. Some new Carboniferous arachnids from the Ostrava-Karviná coal district. *Časopis pro Mineralogii a Geologii*, 3: 425–434.
- Příbyl, A. 1967. *Moravurus* gen.n. eine neue Xiphosurida Gattung aus dem mährisch-schlesischen Oberkarbon. *Časopis pro Mineralogii a Geologii*, 12: 457–460.
- Pritchard A. E. 1956. A new superfamily of trombidiform mites with the description of a new family, genus and species (Acarina: Iolinoidea: Iolinidae: *Iolina nana*). *Annals of the Entomological Society of America*, 49: 204–206.
- Protescu, O. 1937. Etude géologique et paléobiologique de l'ambre roumain. *Bulletin de la Société române Géologique*, 3: 65–110.
- Prószyński, J. & Żabka, M. 1980. Remarks on Oligocene amber spiders of the family Salticidae. *Acta Palaeontologica Polonica*, 25: 213–223.
- Pruvost, P. 1912. Note sur les Araignées du terrain houiller du Nord de la France. *Annales de la Société Géologique du Nord*, 41: 85–100.
- Pruvost, P. 1919. *Introduction à l'étude du terrain houiller du Nord et du Pas-de-Calais: La faune continentale du terrain houiller de la France*. pp. 339–364. *Classe des Arachnides*. Thèse Université de Lille, Lille.
- Pruvost, P. 1922. Les arachnides fossiles du Houiller de Belgique. *Annales de la Société Scientifique de Bruxelles*, 41: 349–355.
- Pruvost, P. 1926. Description de deux fossiles du terrain houiller de Noeux (*Anthracosiro corsini*, nov. sp. et *Fayolia sterzeli* Weiss). *Annales de la Société Géologique du Nord*, 51: 144–149.
- Pruvost, P. 1930. La Faune continentale du terrain houiller de la Belgique. Arachnides. *Mémoires du Musée royal d'Histoire naturelle de Belgique*, 44: 206–217.
- Pruvost, P. 1939. *Euypterus (Anhraconectes) corneti* du Westphalien A du couchant de Mons. *Annales de la Société Scientifique de Bruxelles*, 59: 56–59.
- Qin, T. K. & Halliday, R. B. 1997. Eriorhynchidae, a new family of Prostigmata (Acarina), with a cladistic analysis of eupodoid species of Australia and New Zealand. *Systematic Entomology*, 22: 151–171.
- Quintero Jr., D. 1996. Revision de la clasification de Amblypygidos pulvanados: creacion de subordenes, una nueva familia y un nuevo genero con tres nuevas especies (Arachnida: Amblypygi). 203–212. In Eberhardt, W. G., Lubin, Y. D. & Robinson, B. C. (eds). *Proceedings of the Ninth International Congress of Arachnology, Panama 1983*. Smithsonian Institution Press, Washington, DC, xx pp.
- Racheboeuf, P. R. 1992. *Valloisella lievinensis* n. g. n. sp.: nouveau Xiphosure carbonifère du nord de la France. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1992(6): 336–342.

- Racheboeuf, P. R., Vannier, J. & Anderson, L. I. 2002. A new three-dimensionally preserved xiphosuran chelicerate from the Montceau-les-Mines Lagerstätte (Carboniferous, France). *Palaeontology*, 45: 125–147.
- Ramírez, M. J. & Grismado, C. J. 1997. A review of the spider family Filistatidae in Argentina (Arachnida: Araneae), with a cladistic reanalysis of filistatid genera. *Entomologica Scandinavica*, 28: 319–349.
- Ramsay, G.W. 1960. Sub-fossil mites from the Hutt Valley. *Transactions of the Royal Society of New Zealand*, 88: 575–576.
- Raymond, P. E. 1944. Late Paleozoic xiphosurans. *Bulletin of the Museum of Comparative Zoology*, 94: 475–508.
- Raven, R. J. 1985. The spider infraorder Mygalomorphae (Araneae): cladistics and systematics. *Bulletin of the American Museum of Natural History*, 182: 1–180.
- Redell, J. R. & Cokendolpher, J. C. 1995. Catalogue, bibliography and generic revision of the order Schizomida (Arachnida). *Texas Memorial Museum, Speleological Monographs*, 4: 1–170.
- Reeside, J. B. & Harris, D. V. 1952. A Cretaceous horseshoe crab from Colorado. *Journal of the Washington Academy of Science*, 42: 174–178.
- Reiskind, J. 1986. A new *Lyssomanes* from the Dominican amber and the possible use of insular fossils in building phylogenies. 423. In Barrientos, J. A. (ed.) *Actas X Congresso Internacional de Aracnologia, Jaca. Espanã*, Volume 1. Barcelona.
- Reiskind, J. 1989. The potential use of amber fossils in the study of the biogeography of spiders in the Caribbean with the description of a new species of *Lyssomanes* from Domican amber (Araneae: Salticidae). 217–228. In Woods, C. A (ed.) *Biogeography of the West Indies, past, present and future*. Sandhill Crane Press, Gainesville, Florida.
- Remy, W. & Remy, R. 1959. Arthropodenfunde im Stefan der Halleschen Mulde. *Monograph-Bericht der Deutschen Akademie Wissenschaft Berlin*, 1: 299–312.
- Reuss, A. E. 1855. Palaeontologische Miscellen. III. Über eine neue Krusterspecies aus der Böhmisichen Steinkohlenformation. *Denkschrift der königlich-kaiserlichen Akademie der Wissenschaft in Wien*, 10: 81–83.
- Richter, R. & Richter, E. 1929. *Weinbergina opitzii* n. g., n. sp., ein Schwerträger (Merost. Xiphos.) aus dem Devon (Rheinland). *Senckenbergiana*, 11: 193–209.
- Ribera, C. 2003. El arácanido del Plesiotoceno inferior de Incaral V (Girona, NE de la Península Ibérica). *Paleontologia i Evolució*, 34: 51–53.
- Riek, E. F. 1955. A new xiphosuran from the Triassic sediments at Brookvale, New South Wales. *Records of the Australian Museum*, 23: 281–282.
- Riek, E. F. & Gill, E. D. 1971. A new xiphosuran genus from Lower Cretaceous Freshwater sediments at Koonwarra, Victoria, Australia. *Palaeontology*, 14: 206–210.

- Riquelme, F. & Hill, D. E. 2013. Insights into amber salticids from the Neogene of Middle America, with the first report of Marpissinae (Araneae: Salticidae) from the Chiapas amber. *Peckhamia*, 106.1: 1–5.
- Risso, A. 1826. Animaux articulés: description de quelques Myriapodes, Scorpionides, Arachnides et Acarides, habitant les Alpes Maritimes. In Risso, A. (ed.). *Histoire Naturelle des Principales Productions de l'Europe Méridionale et Principalement de Celles des Environs de Nice et des Alpes Maritimes*. Levraut, Paris, **xx** pp.
- Ritchie, A. 1968. *Lanarkopterus dolichoshelus* (Størmer) gen. nov., a mixopterid eurypterid from the Upper Silurian of the Lesmahagow and Hagshaw Hills inliers, Scotland. *Scottish Journal of Geology*, 4: 317–338.
- Robineau-Desvoidy, J. B. 1828. *Recherches sur l'organisation vertébrale des Crustacés, Arachnides et Insectes*. Comprè Jeune, Paris, 228 pp.
- Roemer, F. 1866. *Protolycosa anthracophila*, eine fossile Spinne aus dem Steinkohlengebirge Oberschlesiens. *Neues Jahresbuch für Mineralogie, Geologie und Paläontologie*: 136–143.
- Roemer, F. 1878. Auffindung und Vorlegung eines neuen Gliderthieres in dem Steinkohlengebiete der Ferdinandsgrube bei Glatz. *Jahresbericht der Schlesischen Gesellschaft für Vaterländische-Kultur.*, pp. 54–55.
- Roewer, C. F. 1912. Die Familien der Assamiden und Phalangodiden der Opiliones-Laniatores. (=Assamiden, Dampetridenten, Phalangodiden, Epedaniden, Biantiden, Zalmoxiden, Samoiden, Palpipediden anderer Autoren.) *Archiv für Naturgeschichte* 78A (3): 1–242.
- Roewer, C.-F. 1913. Die Familie der Gonyleptiden der Opiliones-Laniatores. *Archiv für Naturgeschichte*, 79A (4, 5): 1–256, 257–473.
- Roewer, C.-F. 1923. *Die Webergänge der Erde. Systematische Bearbeitung der bisher bekannten Opiliones*. Gustav Fischer, Jena, 1116 pp.
- Roewer, C.-F. 1933. Solifugae, Palpigradi. 161–480. In Bronn, H. G. (ed.). *Klassen und Ordnung des Tierreichs*. 5: *Arthropoda IV: Arachnoidea*, vol. 5(IV) (4) (2–3). Akademische Verlagsgesellschaft M.B.H, Leipzig.
- Roewer, C.-F. 1934. Solifugae, Palpigradi. 481–723. In Bronn, H. G. (ed.). *Klassen und Ordnung des Tierreichs*. 5: *Arthropoda IV: Arachnoidea*, vol. 5(IV) (4) (4–5). Akademische Verlagsgesellschaft M.B.H, Leipzig.
- Roewer, C.-F. 1935. Zwei myrmecophile Spinnen-Arten Brasiliens. *Veröffentlichungen aus dem Deutschen Kolonial- und Übersee-Museum in Bremen*, 1: 193–197.
- Roewer, C.-F. 1939. Opilioniden im Bernstein. *Palaeobiologica*, 7(1): 1–4.
- Roewer, C.-F. 1942. *Katalog der Araneae von 1758 bis 1940. 1. Band*. Kommissions-Verlag von „NATURA“: 1040 pp.
- Roewer, C.-F. 1943. Über Gonyleptiden. Weitere Webergänge (Arachn., Opil.) XI. *Senckenbergiana*, 26: 12–68.
- Roewer, C.-F. 1951. Über Nemastomatiden. Weitere Webergänge XVI. *Senckenbergiana*, 32: 95–153.
- Rovainen, H. 1953. Subfamilies of European eriophyid mites. *Annales entomologici Fennici*, 19: 83–87.

- Romero, A. & Via Boada, L. 1977. *Tarracolimulus rieki*, nov. gen., nov. sp., nuevo limulido del Triásico de Montral-Alcover (Tarragona). *Cuadernos de Geología Ibérica*, 4: 239–246.
- Ross, A. J. & Vannier, J. 2002. Crustacea (excluding Ostracoda) and Chelicerata of the Purbeck Limestone Group, southern England: a review. *Special Papers in Palaeontology*, 68: 71–82.
- Rößler, R. & Schneider, J. 1997. Eine bemerkenswerte Paläobiocoenose im Unterkarbon Mitteleuropas – Fossilführung und Paläoenvironment der Hainichen-Subgruppe (Erzgebirge-Becken). *Veröffentlichungen des Museums für Naturkunde Chemnitz*, 20: 5–44.
- Roth, J. R. 1851. Ueber fossile Spinnen des lithographischen Schiefers. *Gelehrte Anzeigen herausgegeben von Mitgliedern der Königlichen Bayerischen Akademie der Wissenschaften in München*, 32: 164–167.
- Rowland, J. M. 1975. A partial revision of Schizomida (Arachnida) with descriptions of new species, genus, and family. *Occasional Papers of the Museum, Texas Tech University*, 31: 1–21.
- Rowland, J. M. & Sissom, W. D. 1980. Report on a fossil palpigrade from the Tertiary of Arizona, and a review of the morphology and systematics of the order (Arachnida: Palpigradida). *The Journal of Arachnology*, 8: 69–86.
- Rudkin, D. M., Young, G. A. & Nowlan, G. S. 2008. The oldest horseshoe crab: a new xiphosurid from late Ordovician Konservat-Lagerstätten deposits, Manitoba, Canada. *Palaeontology*, 51: 1–9.
- Rudkin, D. M., Cuggy, M. B., Young, G. A. & Thompson, D. P. 2013. An Ordovician pycnogonid (sea spider) with serially subdivided ‘head’ region. *Journal of Paleontology*, 87: 395–405.
- Ruedemann, R. 1916. Account of some new or little known species of fossils, mostly from the Palaeozoic rocks of New York. *New York State Museum Bulletin*, 189: 7–112.
- Ruedemann, R. 1921. A recurrent Pittsford (Salina) fauna. *New York State Museum Bulletin*, 219–20: 205–215.
- Ruedemann, R. 1926. The Utica and Lorraine Formations of New York, Part 2, Systematic Paleontology, no. 2, Mollusks, Crustacea and Eurypterids. *New York State Museum Bulletin*, 189: 98–112.
- Ruedemann, R. 1942. Some new eurypterids from New York. *New York State Museum Bulletin*, 327: 24–29.
- Russell, L. S. 1953. A new species of eurypterid from the Devonian of Gaspé. *Annual Report of the National Museum for the Fiscal Year 1952–1953, Bulletin*, 132: 83–91.
- Ryke, P. A. J. 1962. The subfamily Rhodacarinae with notes on a new subfamily Ologamasinae (Acarina: Rhodacaridae). *Entomologische Berichte Amsterdam*, 22: 155–162.
- Salter, J. W. 1856. On some new Crustacea from the uppermost Silurian Rocks. *Quarterly Journal of the Geological Society of London*, 12: 26–34.
- Sanchez, J. P., Nava, S., Lareschi, M., Ortiz, P. E. & Guglielmone, A. A. 2010. Finding of an ixodid tick inside a late Holocene owl pellet from northwestern Argentina. *Journal of Parasitology*, 96: 820–822.
- Santiago-Blay, J. A. & Poinar Jr., G. O. 1988. A fossil scorpion *Tityus geratus* new species (Scorpiones: Buthidae) from Dominican amber. *Historical Biology*, 1: 345–354.

- Santiago-Blay, J. A., Fet, V., Soleglad, M. E. & Anderson, S. R. 2004. A new genus and subfamily of scorpions from Lower Cretaceous Burmese amber (Scorpiones: Chaerilidae). *Revista Ibérica de Aracnología*, 9: 3–14.
- Sarle, C. J. 1903. A new eurypterid fauna from the base of the Salina in western New York. *New York State Museum Bulletin*, 69: 1080–1108.
- Sars, G. O. 1891. Pycnogonidea. *Norwegian North-Atlantic Expedition, 1876–1878*, 6 (Zool. 20): 1–163.
- Saupe, E. E. & Selden, P. A. 2009. First fossil Mecysmaucheniiidae (Arachnida, Chelicerata, Araneae), from Lower Cretaceous (uppermost Albian) amber of Charente-Maritime, France. *Geodiversitas*, 31: 49–60.
- Saupe, E. E., Selden, P. A. & Penney, D. 2010. First fossil *Molinaranea* Mello-Leitão, 1940 (Araneae: Araneidae), from middle Miocene Dominican amber, with a phylogenetic and palaeobiogeographical analysis of the genus. *Zoological Journal of the Linnean Society*, 158: 711–725.
- Saupe, E. E., Pérez-de la Fuente, R., Selden, P. A., Delclòs, X., Tafforeau, P. & Soriano, C. 2012. New *Orchestina* Simon, 1882 (Araneae: Oonopidae) from Cretaceous ambers of Spain and France: First spider described using phase-contrast x-ray synchrotron microtomography. *Palaeontology*, 55: 127–143.
- Savage, T. E. 1916. Alexandrian rocks of northeastern Illinois and eastern Wisconsin. *Bulletin of the Geological Society of America*, 27: 305–324.
- Sayre, R. M., Smiley, R. L. & Walter, D. E. 1992. Report of a teneriffiid mite (Acari) in Baltic amber and notes on recent discoveries. *International Journal of Acarology*, 18: 303–305.
- Scharf, W. 1924. Beitrag zur Geologie des Steinkohlengebietes im Südharz. *Jahrbuch des Halleschen Verbands für die Erforschung der Mitteldeutschen Bodenschätze und ihrer Verwaltung*, 4: 404–437.
- Schawaller, W. 1978. Neue Pseudoscorpione aus dem Baltischen Bernstein der Stuttgarter Bernsteinsammlung (Arachnida: Pseudoscorpionidea). *Stuttgarter Beiträge zur Naturkunde (B)*, 42: 1–21.
- Schawaller, W. 1979a. Erstnachweis eines Skorpions in Dominikanischem Bernstein (Stuttgarter Bernsteinsammlung: Arachnida, Scorpionida). *Stuttgarter Beiträge zur Naturkunde (B)*, 45: 1–15.
- Schawaller, W. 1979b. Erstnachweis der Ordnung Geisselspinnen in Dominikanischem Bernstein (Stuttgarter Bernsteinsammlung: Arachnida, Amblypygi). *Stuttgarter Beiträge zur Naturkunde (B)*, 50: 1–12.
- Schawaller, W. 1980a. Fossile Chthoniidae in Dominikanischem Bernstein, mit phylogenetischen Anmerkungen (Stuttgarter Bernsteinsammlung: Arachnida, Pseudoscorpionidea). *Stuttgarter Beiträge zur Naturkunde (B)*, 63: 1–19.
- Schawaller, W. 1980b. Erstnachweis tertärer Pseudoscorpione (Chernetidae) in Dominikanischen Bernstein. *Stuttgarter Beitrag zur Naturkunde (B)*, 57: 1–20.
- Schawaller, W. 1981. Cheiridiidae in Dominikanischem Bernstein, mit Anmerkungen zur morphologischen Variabilität (Stuttgarter Bernsteinsammlung: Arachnida, Pseudoscorpionidea). *Stuttgarter Beiträge zur Naturkunde (B)*, 75: 1–14.

- Schawaller, W. 1982a. Zwei weitere Skorpione in Dominikanischem Bernstein (Stuttgarter Bernsteinsammlung: Arachnida, Scorpionida). *Stuttgarter Beiträge zur Naturkunde (B)*, 82: 1–14.
- Schawaller, W. 1982b. Der erste Pseudokorpion (Chernetidae) aus Mexicanischem Bernstein. *Stuttgarter Beiträge zur Naturkunde (B)*, 85: 1–9.
- Schawaller, W. 1982c. Spinnen der Familien Tetragnathidae, Uloboridae und Dipluridae in Dominikanischem Bernstein und allgemeine Gesichtspunkte (Arachnida, Araneae). *Stuttgarter Beiträge zur Naturkunde (B)*, 89: 1–19.
- Schawaller, W. 1982d. Zur fossilen Spinnenfauna des Pliozäns von Willershausen in Norddeutschland (Arachnida, Araneae). *Berichte der Naturhistorischen Gesellschaft zu Hannover*, 125: 89–95.
- Schawaller, W. 1984. The family Selenopidae in Dominican amber (Arachnida: Araneae). *Stuttgarter Beiträge zur Naturkunde (B)*, 103: 1–8.
- Schawaller, W., 1991. The first Mesozoic pseudoscorpion, from Cretaceous Canadian amber. *Palaeontology*, 34: 971–976.
- Schawaller, W. & Ono H. 1979. Fossile Spinnen aus miozänen Sedimenten des Randecker Maars in SW-Deutschland (Arachnida: Araneae). *Jahreshefte der Gesellschaft für Naturkunde in Württemberg*, 134: 131–141.
- Schawaller, W., Shear, W. A. & Bonamo, P. M. 1991. The first Paleozoic pseudoscorpions (Arachnida, Pseudoscorpionida). *American Museum Novitates*, 3009: 1–17.
- Schille, F. 1916. Entomologie aus der Mammut- und Rhinoceros-Zeit Galiziens. *Entomologische Zeitschrift*, 30: 42–43.
- Schimkewitsch, W. 1913. Ein Beitrag zur Klassifikation der Pantopoden. *Zoologischen Anzeiger*, 41: 597–615.
- Schimper, W. P. 1853. Paleontologica alsatica ou fragments paléontologiques des différents terrains stratifiés qui se recontrent en Alsace. *Mémoires de la Société du Muséum d'Histoire Naturelle de Strasbourg*, 4: 1–10.
- Schmidt, A. R., Jancke, S., Lindquist, E. E., Ragazzi, E., Roghi, G., Nascimbene, P. C., Schmidt, K., Wappler, T. & Grimaldi, D. A. 2012. Arthropods in amber from the Triassic period. *Proceedings of the National Academy of Science, USA*, doi/10.1073/pnas.1208464109.
- Schmidt, A. R., Perrichot, V., Svojtka, M. Anderson, K. B., Belete, K. H., Bussert, R., Dörfelt, H., Jancke, S., Mohr, B., Mohrmann, E., Nascimbene, P. C., Nel, A., Nel, P., Ragazzi, E., Roghi, G., Saupe, E. E., Schmidt, K., Schneider, H., Selden, P. A., Vávra, N. 2010. Cretaceous life captured in amber. *Proceedings of the National Academy of Sciences, USA*: doi/10.1073/pnas.1000948107.
- Schmidt, F. 1883. Nachtrag zur Monographie der Russischen Leperditen II. Die Crustaceenfauna der Euryptereenschichten von Rootzikull auf Oesel. *Miscellanea silurica III. Memoirs of the Academy of Science de St. Petersbourg*, 31: 28–85.
- Schram, F. R. 1979. Limulines of the Mississippian Bear Gulch Limestone of Central Montana, USA. *Transactions of the San Diego Society of Natural History*, 19: 67–74.

- Schultka, S. 1991. *Trigonotarbus stoermeri* n. sp. – ein Spinnentier aus den Bensberger Schichten (Ems/Unter-Devon) des Rheinischen Schiefergebirge. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 183: 375–390.
- Schuster, R. 1963. *Thalassozetes riparius* n. gen., n. sp., eine litoralbewohnende Oribatide von bemerkenswerter morphologischer Variabilität (Oribatei-Acari). *Zoologischer Anzeiger*, 171: 391–403.
- Scopoli, J. A. 1763. *Entomologia Carniolica, exhibens Insecta Carniolae indigena et distributa in ordines, genera, species, varietates. Methodo Linnaeana. Vindobonae*, 1763: 420 pp.
- Scott, A. G. 2003. Sub-fossil spiders from Holocene peat cores. *Journal of Arachnology*, 31: 1–7.
- Scudder, S. H. 1868. Supplement to descriptions of Articulates. Description of fossil insects found on Mazon Creek and near Morris, Grundy Co., Ill. *Geological Survey of Illinois*, 3: 566–572.
- Scudder, S. H. 1876. New and interesting insects from the Carboniferous of Cape Breton. *Canadian Naturalist and Quarterly Journal of Science*, 8: 88–90.
- Scudder, S. H. 1878. Additions to the Insect-Fauna of the Tertiary Beds at Quesnel, British Columbia. *Geological Survey of Canada. Report of Progress*, 1876–1877: 457–464.
- Scudder, S. H. 1884. A contribution to our knowledge of Paleozoic Arachnida. *Proceedings of the American Academy of Arts and Sciences*, 20: 13–22.
- Scudder, S. H. 1885. 3. Classe. Arachnoidea. Spinnen. Skorpione. 732–746. In Zittel, K. A. (ed), *Handbuch der Paläontologie. I. Abtheilung. Palaeozoologie* 2. R. Oldenbourg, München & Leipzig.
- Scudder, S. H. 1890a. The Tertiary Insects of North America. *Report of the United States Geological Survey*, 13: 734 pp.
- Scudder, S. H. 1890b. Illustrations of the Carboniferous Arachnida of North America, of the orders Anthracomarti and Pedipalpi. *Memoirs of the Boston Society of Natural History*, 4: 443–456.
- Scudder, S. H. 1891. Index to the known fossil insects of the world including myriapods and arachnids. *Bulletin of the United States Geological Survey* 71: 1–744.
- Seemann, F. 1906. Beiträge zur Gigantostrakenfauna Böhmens. *Beiträge zur Paläontologie Österreich-Ungarns und des Orients*, 19: 49–57.
- Selden, P. A. 1990. Lower Cretaceous spiders from the Sierra de Montsech, north-east Spain. *Palaeontology*, 33: 257–285.
- Selden, P. A. 1992. Revision of the fossil ricinuleids. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 83: 595–634.
- Selden, P. A. 1996. First fossil mesothelid spider from the Carboniferous of France. *Revue suisse de Zoologie*, hors série: 585–596.
- Selden, P. A. 2000. *Palaeothelid*, replacement name for the fossil mesothelid spider *Eothelid* non Rowell. *Bulletin of the British arachnological Society*, 11: 292.

- Selden, P. A. 2001. Eocene spiders from the Isle of Wight with preserved respiratory structures. *Palaeontology*, 44: 695–729.
- Selden, P. A. 2002. First British Mesozoic spider, from Cretaceous amber of the Isle of Wight, southern England. *Palaeontology*, 45: 973–983.
- Selden, P. A. 2010. A theridiosomatid spider from the Early Cretaceous of Russia. *Bulletin of the British Arachnological Society*, 15: 69–78.
- Selden, P. A. 2014a. A new spider (Araneae: Haplogynae: Plectreuridae) from the Cretaceous Fossil-Lagerstätte of El Montsec, Spain. *The Journal of Arachnology*, 42: 16–23.
- Selden, P. A. 2014b. Spiders (Arachnida: Araneae) from the Insect Limestone (Bembridge Marls, Late Eocene) of the Isle of Wight, southern England. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 104: 1–8.
- Selden, P. A. & Beattie, R. G. 2013. A spider fossil from the Jurasic Talbragar Fossil Fish Bed of New South Wales. *Alcheringa*, 37: 203–208.
- Selden, P. A. & Drygant, D. M. 1987. A new xiphosuran from the Silurian of Podolia, Ukraine, USSR. *Palaeontology*, 30: 537–542.
- Selden, P. A. & Dunlop, J. A. 2014. The first fossil spider (Araneae: Palpimanoidea) from the Lower Jurassic (Grimmen, Germany). *Zootaxa*, 3894: 161–168.
- Selden, P. A. & Gall, J.-C. 1992. A Triassic mygalomorph spider from the northern Vosges, France. *Palaeontology*, 35: 211–235.
- Selden, P. A. & Huang, D.-y. 2010. The oldest haplogyne spider (Araneae: Plectreuridae), from the Middle Jurassic of China. *Naturwissenschaften*, 97: 449–459.
- Selden, P. A. & Penney, D. 2003. Lower Cretaceous spiders (Arthropoda: Arachnida: Araneae) from Spain. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 2003: 175–192.
- Selden, P. A. & Penney, D. 2009. A fossil spider (Araneae: Pisauridae) of Eocene age from Horsefly, British Columbia, Canada. *Contributions to Natural History*, 12: 1269–1282.
- Selden, P. A. & Shear, W. A. 1996. The first Mesozoic solifuge (Arachnida), from the Cretaceous of Brazil, and a redescription of the Palaeozoic solifuge. *Palaeontology*, 39: 583–604.
- Selden, P. A. & Siveter, D. J. 1987. The origin of the limuloids. *Lethaia*, 20: 383–392.
- Selden, P. A., Baker, A. S. & Phipps, K. J. 2008. An oribatid mite (Arachnida: Acari) from the Oxford Clay (Jurassic: Upper Callovian) of South Cave Station Quarry, Yorkshire, UK. *Palaeontology*, 51: 623–633.
- Selden, P. A., Casado, F. C. & Mesquita, M. V. 2006. Mygalomorph spiders (Araneae: Dipluridae) from the Lower Cretaceous Crato Lagerstätte, Araripe Basin, north-east Brazil. *Palaeontology*, 49: 817–826.
- Selden, P. A., Huang D.-y., Ren D. 2008. Palpimanoid spiders from the Jurassic of China. *Journal of Arachnology*, 36: 306–321.

- Selden, P. A., Shear, W. A. & Bonamo, P. M. 1991. A spider and other arachnids from the Devonian of New York, and reinterpretations of Devonian Araneae. *Palaeontology*, 34: 241–281.
- Selden, P. A., Shear, W. A. & Sutton, M. D. 2008. Fossil evidence for the origin of spider spinnerets, and a proposed arachnid order. *Proceedings of the National Academy of Sciences of the United States of America*, 105: 20781–20785.
- Selden, P. A., Shih, C.-K. & Ren, D. 2011. A golden orb-weaver spider (Araneae: Nephilidae: Nephila) from the Middle Jurassic of China. *Biology Letters*, 7: 775–778.
- Selden, P. A., Shih, C.-K. & Ren, D. 2013. A giant spider from the Jurassic of China reveals greater diversity of the orbicularian stem group. *Naturwissenschaften*, 100: 1171–1181.
- Selden, P. A., Anderson, J. M., Anderson, H. M. & Fraser, N. C. 1999. Fossil araneomorph spiders from the Triassic of South Africa and Virginia. *Journal of Arachnology*, 27: 401–414.
- Selden, P. A., Nam, K.-s., Kim, S. H. & Kim, H. J. 2012. A fossil spider from the Cretaceous of Korea. *Journal of Palaeontology*, 86: 1–6.
- Selden, P. A., Shcherbakov, D. E., Dunlop, J. A. & Eskov, K. Y. 2014. Arachnids from the Carboniferous of Russia and Ukraine, and the Permian of Kazakhstan. *Paläontologische Zeitschrift*, 88: 297–307.
- Sellnick, M. 1918. Die Oribatiden der Bernsteinsammlung der Universität Königsberg I Pr. *Schriften der Physikalisch-Ökonomischen Gesellschaft zu Königsberg (1919)*, 59: 21–42.
- Sellnick, M. 1922. Milben der Sammlung des Deutschen Entomologischen Instituts. I. Oribatidae. *Entomologische Mitteilungen*, 11: 18–20.
- Sellnick, M. 1928. Formenkreis: Hornmilben, Oribatei. In Brohmer, P., Ehrmann, P. & Ulmer, G. (eds). *Die Tierwelt Mitteleuropas*, 3, 4(9): 1–42.
- Sellnick, M. 1931. Milben im Bernstein. *Bersteinforschung*, 2: 148–180.
- Semper, M. 1898. Die Gigantostraken des älteren böhmischen Paläozoicum. *Beiträge zur Paläontologie Österreich-Ungarns und des Orients*, 2: 71–88.
- Shear, W. A., 1980. A review of the Cyphophthalmi of the United States and Mexico, with a proposed reclassification of the suborder (Arachnida, Opiliones). *American Museum Novitates*, 2705: 1–34.
- Shear, W. A., 1986. A cladistic analysis of the opilionid superfamily Ischyropsalidoidea, with description of the new family Ceratolasmatidae, the new genus *Acuclavella* and four new species. *American Museum Novitates*, 2844: 1–29.
- Shear, W. A., 1993. The genus *Troglosiro* and the new family Troglosironidae (Opiliones, Cyphophthalmi). *Journal of Arachnology*, 21: 81–90.
- Shear, W. A. 2000. *Gigantocharinus szatmaryi*, a new trigonotarbid arachnid from the Late Devonian of North America (Chelicerata, Arachnida, Trigonotarbida). *Journal of Paleontology*, 74: 25–31.
- Shear, W. A. 2010. New species and records of ortholasmatine harvestmen from México, Honduras, and the western United States (Opiliones, Nemastomatidae, Ortholasmatinae). *ZooKeys*, 52: 9–45.

- Shear, W. A., Selden, P. A., Rolfe, W. D. I., Bonamo, P. M. & Grierson, J. D. 1987. New terrestrial arachnids from the Devonian of Gilboa, New York. *American Museum Novitates*, 2901: 1–74.
- Sharma, P. P. & Giribet, G. 2011. The evolutionary and biogeographic history of the armoured harvestmen – Laniatores phylogeny based on ten molecular markers, with the description of two new families of Opiliones (Arachnida). *Invertebrate Systematics*, 25: 106–142.
- Sharma, P. P., Prieto, C. E. & Giribet, G. 2011. A new family of Laniatores (Arachnida: Opiliones) from the Afrotropics. *Invertebrate Systematics*, 25: 143–154.
- Shipnev, E. S. 2006. A new species of *Adelophthalmus* (Eurypterida) from the Lower Carboniferous of the Krasnoyarsk Region. *Paleontological Journal*, 40: 431–433. [English translation of Russian original]
- Shipnev, E. S. 2012. On some eurypterids (Eurypterida, Chelicerata) from the Devonian of South Siberia. *Paleontological Journal*, 46: 370–377. [English translation of Russian original]
- Shuler, E. W. 1915. A new Ordovician eurypterid. *American Journal of Science*, 4th Series, 39: 551–554.
- Sidorchuk, E. A. & Bertrand, M. 2013. New fossil labidostomatids (Acari: Labidostomatidae) from Eocene amber and presence of an apustulate species in Europe. *Acarologia*, 53: 25–39.
- Sidorchuk, E. A. & Klimov, P. B. 2011. Redescription of the mite *Glaesacarus rhombeus* (Koch & Berendt, 1854) from Baltic amber (Upper Eocene): evidence for female-controlled mating. *Journal of Systematic Palaeontology*, 9: 183–196.
- Sidorchuk, E. A. & Norton, R. A. 2011. The fossil mite family Archaeorchestidae (Acari, Oribatida) I: redescription of *Stieremaeus illibatus* and synonymy of *Strieremaeus* with *Archaeorchestes*. *Zootaxa*, 2993: 34–58.
- Sidorchuk, E. A., Schmidt, A. R., Ragazzi, E., Roghi, G., Lindquist, E. E. 2015. Plant-feeding mite diversity in Triassic amber (Acari: Tetrapodili). *Journal of Systematic Palaeontology*, online.
- Siebold, C. T. E. von. 1850. Ueber *Eriophyes*. *Jahresbericht der Schlesischen Gesellschaft*, 28: 88–89.
- Siegfried, P. 1972. Ein Schwertschwanz (Merostomata, Xiphosurida) aus dem Oberkarbon von Ibbenburen/Westfalen. *Paläontologische Zeitschrift*, 46, 180–186.
- Šilhavý, V. 1973. Two new systematic groups of the gonyleptomorph phalangids from the Antillean-Caribbean Region. Agoristenidae fam. n. and Caribbantinae subfam. n. *Věstník Československé Společnosti Zoologické*, 37: 110–143.
- Šilhavý, V. 1979. New American representatives of the subfamily Samoinae (Opiliones, Phalangodidae, Arach.). *Annotationes zoologicae et botanicae, Bratislava*, 130: 1–27.
- Simon, E. 1864. *Histoire naturelle des Araignées (Aranéides)*. Paris, 540 pp.
- Simon, E. 1874. *Les arachnides de France. Tome 1*. Paris, 272 pp.
- Simon, E. 1876a. *Les Arachnides de France. Tome 3*. Paris, 360 pp.
- Simon, E. 1876b. Etude sur les Arachnides du Congo. *Bulletin de la Société zoologique de France*, 1: 12–15, 216–224.

- Simon, E. 1879a. *Les Arachnides de France VII. Contenant les ordes des Chernetes, Scorpiones et Opiliones.* Paris, : 332 pp.
- Simon, E. 1879b. Essai d'une classification des Opiliones Mecostethi. Remarques synonymiques et descriptions d'espèces nouvelles. *Annales de la Société Entomologique de Belgique*, 22: 183–241.
- Simon, E. 1880. Études arachnologiques 12e Mémoire(1). XVII. Descriptions de Genres et Espèces de l'ordre des Scorpiones. *Annales de la Société Entomologique de France*, (5)10: 377–398.
- Simon, E. 1881. *Les Arachnides de France. Tome 5, 1^{re} partie.* Paris, 179 pp.
- Simon, E. 1882. Etudes arachnologiques. 13^e Mémoire. 20. Descriptions d'espèces et de genres nouveaux de la famille des Dysderidae. *Annales de la Société Entomologique de France*, (6) 2: 201–240.
- Simon, E. 1884a. Note synonymique sur les genres *Prodidomus* Hentz et *Miltia* E.S. *Annales de la Société Entomologique de Belgique*, 28: 302.
- Simon, E. 1884b. Note complémentaire sur la famille des Archaeidae. *Annali del Museo Civico di Storia Naturale di Genova*, 20: 373–380.
- Simon, E. 1884c. *Les Arachnides de France. Tome 5, 2^e et 3^e parties.* Paris, pp. 180–808.
- Simon, E. 1884d. Description d'une nouvelle famille de l'ordre des Araneae (Bradystichidae). *Annales de la Société Entomologique de Belgique*, 28: 297–301.
- Simon, E. 1885a. Etudes arachnologiques. 17e Mémoire. XXVI. Arachnides recueillis dans la vallée de Templé et sur le mont Ossa (Thessalie). *Annales de la Société Entomologique de France*, 5: 209–217.
- Simon, E. 1885b. Etude sur les Arachnides recueillis en Tunisie en 1883 et 1884 par MM. A. Letourneux, M. Sébillot et Valéry Mayet, membres de la Mission de l'Exploration scientifique de la Tunisie. *In Exploration scientifique de la Tunisie*, Paris, 55 pp.
- Simon, E. 1885c. Etudes arachnologiques. 18e Mémoire. XXVI. Matériaux pour servir à la fauna des Arachnides du Sénégal. (Suivi d'un appendice intitulé: Descriptions de plusieurs espèces africaines nouvelles). *Annales de la Société Entomologique de France*, 5: 345–396.
- Simon, E. 1887. Espèces et genres nouveaux de la famille des Sparassidae. *Bulletin de la Société zoologique de France*, 12: 466–474.
- Simon, E. 1888. Etudes arachnologiques. 21^e Mémoire. 29. Descriptions d'espèces et de genres nouveaux de l'Amérique centrale et des Antilles. *Annales de la Société Entomologique de France*, (6) 8: 203–216.
- Simon, E. 1889a. Etudes arachnologiques. 21^e Mémoire. 31. Descriptions d'espèces et de genres nouveaux de Madagascar et de Mayotte. *Annales de la Société Entomologique de France*, (6) 8: 223–236.
- Simon, E. 1889b. Arachnides. *In Voyage de M. E. Simon au Venezuela (décembre 1887 – avril 1888).* 4^e Mémoire. *Annales de la Société Entomologique de France*, (6) 9: 169–220.
- Simon, E. 1890. Etudes arachnologiques. 22^e Mémoire. 34. Etude sur les Arachnides de l'Yemen. *Annales de la Société Entomologique de France*, 10: 77–124.

- Simon, E. 1891a. Observations biologiques sur les Arachnides. I. Araignées sociables. In Voyage de M. E. Simon au Venezuela (Décembre 1887 – avril 1888). 11^e Mémoire. *Annales de la Société Entomologique de France*, 60: 5–14.
- Simon, E. 1891b. On the spiders of the Island of St. Vincent. Part I. *Proceedings of the Zoological Society of London*, 1891: 549–575.
- Simon, E. 1892a. Arachnides. In Raffray, A., Bolivar, I. & Simon, E. (eds) Etude sur les Arthropodes cavernicoles de île Luzon, Voyage de M. E. Simon aux îles Philippines (Mars et avril 1890). *Annales de la Société Entomologique de France*, 61: 35–52.
- Simon, E. 1892b. *Histoire naturelle des Araignées. Volume 1, part 1*. Roret, Paris, pp. 1–254.
- Simon, E. 1893. *Histoire naturelle des Araignées. Volume 1, part 2*. Roret, Paris, pp. 255–488.
- Simon, E. 1894. *Histoire naturelle des Araignées, Volume 1, part 3*. Roret, Paris, pp. 489–760.
- Simon, E. 1895. *Histoire naturelle des Araignées, Volume 1, part 4*. Roret, Paris, pp. 761–1084.
- Simon, E. 1896. Description d'un Arachnide cavernicole de l'Afrique australe. *Bulletin de la Société Entomologique de France*, 1869: 285–286.
- Simon, E. 1897a. *Histoire naturelle des Araignées, Volume 2, part 1*. Roret, Paris, 1–192.
- Simon, E. 1897b. On the Spiders of the Island of St. Vincent. Part III. *Proceedings of the Zoological Society of London*, 1897: 860–890.
- Simon, E. 1898a. *Histoire naturelle des Araignées, Volume 2, part 2*. Roret, Paris, 1–269.
- Simon, E. 1898b. Etude sur les Arachnides de la région des Maures (Var.) *Feuille des Jeunes Naturalistes*, (3) 29: 2–4.
- Simon, E. 1900. Descriptions d'arachnides nouveaux de la famille des Attidae. *Annales de la Société Entomologique de Belgique*, 44: 381–407.
- Simon, E. 1903. *Histoire naturelle des Araignées, Volume 2, part 4*. Roret, Paris, 669–1080.
- Simon, E. 1929. *Les Arachnides de France. Tome 6*. Paris, pp. 533–772.
- Simon, R. 1971. Neue Arthropodenfunde aus dem Stephan der Halleschen Mulde. *Bericht der Deutschen Gesellschaft für Geologische Wissenschaft, Reihe A: Geologie/Paläontologie*, 16: 53–62.
- Simonetta, A. M. & Delle Cave, L. 1978. Una possibile interpretazione filogenetica degli artropodi paleozoici. *Bollettino di zoologia*, 45: 87–90.
- Simpson, S. 1951. A new Eurypterid from the Upper Old Red Sandstone of Portishead. *Annals and Magazine of Natural History, series 12*, 4: 849–861.
- Siveter, D. J. & Selden, P. A. 1987. A new, giant xiphosurid from the lower Namurian of Weardale, County Durham. *Proceedings of the Yorkshire Geological Society*, 46: 153–168.
- Siveter, D. J., Sutton, M. D., Briggs, D. E. G. & Siveter, D. J. 2004. A Silurian sea spider. *Nature*, 431: 978–980.
- Sivhed, U. & Wallwork, J. A. 1978. An early Jurassic oribatid mite from southern Sweden. *Geologiska Föreningens i Stockholm Förhandlingar*, 100: 65–70.

- Smith, F. P. 1902. The spiders of Epping Forest. *Essex Naturalist*, 12: 181–201.
- Sørensen, W. E. 1884. Opiliones Laniatores (Gonyleptides W. S. Olim) Musei Hauniensis. *Naturhistorisk Tidsskrift, Kjøbenhavn, series 3*, 14: 555–646.
- Sørensen, W. 1886. Opiliones. pp. 53–86. In Koch, L. & Keyserling, E. (eds) *Die Arachniden Australiens nach der Natur Beschrieben und Abgebildet*. Bauer und Raspe, Nürnberg, **xx** pp.
- Sørensen, W. 1932. Descriptiones Laniatorum (Arachnidorum Opilionum Subordinis). (Opus posthum recognovit et editit Kai L. Henriksen). – *Kongelige Danske Videnskabernes Selskabs Skrifter - Naturvidenskab og Mathematiske Afdeling, København*, ser. 9, 3(4): 197–422.
- Soriano, C., Archer, M., Azar, D., Creaser, P., Delclòs, X., Godhelp, H., Hand, S., Jones, A., Nel, A., Nèraudeau, D., Ortega-Blanco, J., Pérez-de la Fuente, R., Perrichot, V., Saupe, E., Solorzano-Kraemer, M., Taffreau, P. 2010. Synchrotron X-ray imaging on inclusions in amber. *Comptes Rendus Palevol*, 9, 361–368.
- Southcott, R. V. 1957. Description of a new Australian raphignathoid mite, with remarks on the classification of the Trombidiformes (Acarina). *Proceedings of the Linnean Society of New South Wales*, 81(3): 306–312.
- Southcott, R. V. & Lange, R. T. 1971. Acarine and other microfossils from the Maslin eocene, South Australia. *Records of the South Australian Museum*, 16(**xx**): 1–21.
- Stahnke, H. L. 1940. The scorpions of Arizona. *Iowa State College Journal of Science*, 15: 101–103. [Thesis abstract.]
- Stainier, X. 1917. On a new eurypterid from the Belgian Coal Measures. *Quarterly Journal of the Geological Society*, 71: 639–647.
- Sterzel, J.T. 1918. Die organischen Reste des Kulms und Rotliegenden der Gegend von Chemnitz. *Abhandlungen der Königlich Sächsischen Gesellschaft der Wissenschaften, Mathematisch-physikalische Klasse*, 35: 1–315.
- Stock, J. H. 1954. Papers from Dr. Th. Mortensen's Pacific expedition 1914–1916. LXXVII. Pycnogonida from Indo-West-Pacific, Australian, and New-Zealand waters. *Videnskabelige Meddelelser fra Dansk naturhistorisk Forening*, 116(1): 1–168.
- Stott, C. A., Tetlie, O. E., Braddy, S. J., Nowlan, G. S., Glasser, P. M. & Devereux, M. G. 2005. A new eurypterid (Chelicerata) from the Upper Ordovician of Manitoulin Island, Ontario, Canada. *Journal of Paleontology*, 79: 1166–1174.
- Størmer, L. 1934a. Downtonian Merostomata from Spitsbergen with remarks on the suborder Synziphosura. *Skrifter utgitt av Det Norske Videnskaps-Akademi i Oslo, I. Matem.-Naturvid. Klasse*, 1933(3): 1–26.
- Størmer, L. 1934b. Merostomata from the Downtonian Sandstones of Ringerike, Norway. *Skrifter utgitt av Det Norske Videnskaps-Akademi i Oslo, I. Matem.-Naturvid. Klasse*, 1933(10): 1–125.
- Størmer, L. 1934c. Über den neuen von W. Gross beschriebenen Eurypteriden aus dem Unterdevon von Overath im Rheinland. *Jahrbuch der Preussischen Geologischen Landesanstalt*, 55: 284–291.

- Størmer, L. 1934d. A new Eurypterid from the Saaremaa-(Oesel-)Beds in Estonia. *Publications of the Geological Institution of the University of Tartu*, 37: 1–8.
- Størmer, L. 1936a. Eurypteriden aus dem Rheinischen Unterdevon. *Abhandlungen der Preussischen Geologischen Landesanstalt, Neue Folge*, 175: 1–74.
- Størmer, L. 1936b. *Mixopterus dolichoshelus* (Laurie MS), a Downtonian eurypterid from Scotland. *Summary of Progress of the Geological Survey for 1934*: 41–46.
- Størmer, L. 1951. A new eurypterid from the Ordovician of Montgomeryshire, Wales. *Geological Magazine*, 88: 409–422.
- Størmer, L. 1952. Phylogeny and taxonomy of fossil horseshoe crabs. *Journal of Paleontology*, 26: 630–639.
- Størmer, L. 1963. *Gigantoscorpio willsi*, a new scorpion from the Lower Carboniferous of Scotland and its associated preying microorganisms. *Skrifter Utgitt av det Norske Videnskaps-Akademi i Oslo. Matematisk-Naturvidenskabelig Klasse*, 8: 1–171.
- Størmer, L. 1969. Eurypterids from the Lower Devonian of Willwerath, Eifel. *Senckenbergiana lethaea*, 50: 21–35.
- Størmer, L. 1970. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 1: Arachnida. *Senckenbergiana lethaea*, 51: 335–369.
- Størmer, L. 1972. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 2: Xiphosura. *Senckenbergiana lethaea*, 53: 1–29.
- Størmer, L. 1973. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 3: Eurypterida, Hughmilleridae. *Senckenbergiana lethaea*, 54: 119–205.
- Størmer, L. 1974. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 4: Eurypterida, Drepanopteridae, and other groups. *Senckenbergiana lethaea*, 54: 359–451.
- Størmer, L. 1976. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 5: Myriapoda and additional forms, with general remarks on the fauna and problems regarding invasion of land by arthropods. *Senckenbergiana lethaea*, 57: 87–183.
- Størmer, L. & Waterston, C. D. 1968. *CyrtocTenus* gen. nov., a large late Palaeozoic arthropod with pectinate appendages. *Transactions of the Royal Society Edinburgh*, 68: 63–104.
- Strand, E. 1926. Miscellanea nomenclatorial zoological et palaeontologica. I–II. *Archiv für Naturgeschichte A*, 92(8): 30–75.
- Strand, E. 1929. Zoological and palaeontological nomenclatorial notes. *Acta Universitatis Latviensis*, 20: 29 pp.
- Strand, E. 1932. Miscellanea nomenklatorica zoologica et palaeontologica, III, IV. *Folia zoologica et hydrobiologica*, 4: 133–147, 193–196.
- Strand, E. 1942. Miscellanea nomenclatoria zoologica et palaeontologica. *Folia Zoologica et Hydrobiologica*, 11: 386–402.
- Straus, A. 1967. Zur Paläontologie des Pliozäns von Willershausen. *Berichte der Naturhistorischen Gesellschaft Hannover*, 111: 15–24.

- Strenzke, K. 1963. Entwicklung und Verwandschaftsbeziehungen der Oribatidengattung *Gehyphochthonius* (Arach., Acari). *Senckenbergiana Biologica*, 44: 231–255.
- Stumm, E. C. & Kjellesvig-Waering, E. N. 1962. A new eurypterid from the Upper Silurian of southern Michigan. *Contributions from the Museum of Paleontology, The University of Michigan*, 17: 195–204.
- Stur, D. 1877. Die Culm-Flora der Ostrauer und Waldenburger Schichten. *Abhandlung der königliche geologische Reichanstalt*, 4: 5.
- Subías, L. S. 2004. Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo. *Graellsia* 60 (número extraordinario), 3–305. Available from: <http://www.ucm.es/info/zoo/Artropodos/Catalogo.pdf>.
- Subías, L. S. & Arillo, A. 2002. Oribatid mite fossils from the Upper Devonian of South Mountain, New York and the Lower Carboniferous of County Antrim, Northern Ireland (Acariformes, Oribatida). *Estudios del Museo de Ciencias Naturales de Alava*, 17: 93–106.
- Sundevall, J.C. 1833. *Conspectus Arachnidium*. C. F. Berling, Londini Gothorum, 39 pp.
- Swartz, C. K. 1923. Order Eurypterida. 716–778. In Swartz, C. K., Prouty, W. F., Ulrich, E. O. & Bassler, R. S. (eds). *Silurian Volume*. Maryland Geological Survey, 795 pp.
- Taczanowski, L. 1879. Les aranéides du Pérou central (suite). *Horae Societatis entomologicae Rossicae*, 15: 102–136.
- Tasch, P. 1961. Paleolimnology: part 2 – Harvey and Sedgwick counties, Kansas: stratigraphy and biota. *Journal of Paleontology*, 35: 836–865.
- Tasch, P. 1963. Paleolimnology: part 3 – Marion and Dickinson counties, Kansas, with additional sections in Harvey and Sedgwick counties: stratigraphy and biota. *Journal of Paleontology*, 37: 1233–1251.
- Tesakov, A. S. & Alekseev, A.S. 1992. Myriapod-like arthropods from the Lower Devonian of central Kazakhstan. *Paleontological Journal*, 26: 18–23.
- Tesakov, A. S. & Alekseev, A.S. 1998. *Maldybulakia* – new name for *Lophodesmus* Tesakov and Alekseev, 1992 (Arthropoda). *Paleontological Journal*, 32: 29.
- Tetlie, O. E. 2002. A new *Baltoeurypterus* (Eurypterida: Chelicerata) from the Wenlock of Norway. *Norwegian Journal of Geology*, 82: 37–44.
- Tetlie, O. E. 2006a. Two new Silurian species of *Eurypterus* (Chelicerata: Eurypterida) from Norway and Canada and the phylogeny of the genus. *Journal of Systematic Palaeontology* 4: 397–412.
- Tetlie, O. E. 2006b. Eurypterida (Chelicerata) from the Welsh Borderlands, England. *Geological Magazine*, 143: 723–735.
- Tetlie, O. E. & Braddy, S.J. 2004. The first Silurian chasmataspid, *Loganamaraspis dunlopi* gen. et sp. nov. (Chelicerata: Chasmataspidida) from Lesmahagow, Scotland, and its implications for eurypterid phylogeny. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 94: 227–234.

- Tetlie, O. E. & Briggs, D. E. G. 2009. The origin of pterygotid eurypterids (Chelicerata: Eurypterida). *Palaeontology*, 52: 1141–1148.
- Tetlie, O. E. & Dunlop, J. A. 2008. *Geralinura carbonaria* (Arachnida; Uropygi) from Mazon Creek, Illinois, USA, and the origin of subchelate pedipalps in whip scorpions. *Journal of Paleontology*, 82: 299–312.
- Tetlie, O. E. & Van Roy, P. 2006. A reappraisal of *Eurypterus dumonti* Stainier, 1917 and its position within the Adelophthalmidae Tollerton, 1989. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre* 76: 79–90.
- Tetlie, O. E. & Poschmann, M. 2008. Phylogeny and palaeoecology of the Adelophthalmoidea (Arthropoda; Chelicerata; Eurypterida). *Journal of Systematic Palaeontology*, 6: 237–249.
- Tetlie, O. E., Selden, P. A. & Ren D. 2007. A new Silurian eurypterid (Arthropoda: Chelicerata) from China. *Palaeontology*, 50: 619–625.
- Tetlie O. E., Braddy, S. J., Butler, P.D. & Briggs, D.E.G. 2004. A new eurypterid (Chelicerata: Eurypterida) from the Upper Devonian Gogo Formation of Western Australia, with a review of the Rhenopteridae. *Palaeontology* 47: 801–809.
- Thevenin, A. 1901. Sur le découverte d'arachnides dans le Terrain Houiller de Commentry. *Bulletin de la Société Géologique de France, 4^e Série*, 1: 605–611.
- Thevenin, A. 1902. Sur une araignée du terrain houiller der Valenciennes. *Procès-Verbaux de la Société d'Histoire Naturelle de Autun*, 15: 195–203.
- Thompson, W. D'Arcy 1909. Pycnogonida. In Harmer, S. F. & Shipley, B. E. (eds). *The Cambridge Natural History*, pp. 501–542.
- Thor, S. 1900. Norske hydrachnider IV. – Archiv für Mathematik og Naturvidenskab 23: 1-56.
- Thor, S. 1905. Eine interessante neue Milbengattung aus der schweizerischen Sammlung des Herrn Dr. W. Volz. *Zoologischer Anzeiger*, 28: 505–509.
- Thor, S. 1911a. *Lebertia-Studien XXIV–XXV*. *Zoologischer Anzeiger*, 37: 385–394.
- Thor, S. 1911b. Eine neue Acarinengattung (Teneriffiidae) und zwei neue Gattungen, die eine von Teneriffa, die andre aus Paraguay. *Zoologischer Anzeiger*, 38: 171–179.
- Thor, S. 1927. Acarinologische Notizen. *Zoologischer Anzeiger*, 72: 155–159.
- Thor, S. 1933. Über die prostigmatische Familie: Eupodidae C.L.Koch 1842 und über die Teilung dieser Familie, mit Definitionen der neuen Familien. *Zoologischer Anzeiger*, 101: 271–277.
- Thor, S. 1934. Neue Beiträge zur Kenntnis der invertebraten Fauna von Svalbard. (Nach Sammlungen von Garteninspektor L. Lange, Dozent B. Lynge und dem Verfasser.). *Zoologischer Anzeiger*, 107: 114–139.
- Thor, S. 1935. Übersicht und Einteilung der Familie Trombidiidae W.E. Leach 1814 in Unterfamilien. *Zoologischer Anzeiger*, 109: 107–112.
- Thor, S. 1937. Übersicht der norwegischen *Cryptostigmata* mit einzelnen Nebenbemerkungen. *Saertrykk av Nytt Magasin for Naturvidenskapene*, 77: 275–307.

- Thorell, T. 1856. Recensio critica Aranearum Suecicarum quas descripserunt Clerckius, Linnaeus, de Geerus.
Nova Acta Societas Scientiae Uppsالensis, 2: 61–176.
- Thorell, T. 1869. On European spiders. Part I. Review of the European genera of spiders, preceded by some observations on zoological nomenclature. *Nova Acta Societas Scientiae Uppsالensis*, (3)7: 1–108.
- Thorell, T. 1870a. On European spiders. Part 2. *Nova Acta Societas Scientiae Uppsالensis*, (3)7: 109–242.
- Thorell, T. 1870b. *Remarks on synonyms of European spiders. Part I.* Uppsala, pp. 1–96.
- Thorell, T. 1873. *Remarks on synonyms of European spiders. Part IV.* Uppsala, pp. 375–645.
- Thorell, T. 1875. Diagnoses Aranearum Europaearum aliquot novarum. *Tijdschrift voor Entomologie*, 18: 81–108.
- Thorell, T. 1876a. Études Scorpiologiques. *Atti della Società Italiana di Scienze Naturali*, 19: 75–272.
- Thorell, T. 1876b. On the classification of scorpions. *Annals and Magazine of Natural History, series 4*, 17: 1–15.
- Thorell, T. 1876c. Sopra alcuni Opilioni (Phalangidea) d'Europa e dell'Asia occidentale, con un quadro dei generi europei di quest'Ordine. *Annali del Museo Civico di Storia Naturale (Genoa) series 1*, 8: 452–508.
- Thorell, T. 1881. Studi sui Ragni Malesi e Papuani. III. Ragni dell'Astro Malesia e del Capo York, conservati nel Museo civico di storia naturale di Genova. *Annali del Museo Civico di Storia Naturale di Genova*, 17: 1–727.
- Thorell, T. 1882. Descrizione di Alcuni Aracnidi Inferiori dell' Arcipelago Malese. *Annali del Museo Civico di Storia Naturale di Genova*, 18: 21–69.
- Thorell, T. 1887. Viaggio di L. Fea in Birmania e regioni vicine. II. Primo saggio sui ragni birmani. *Annali del Museo Civico di Storia Naturale di Genova*, 25: 5–417.
- Thorell, T. 1888. Pedipalpi e Scorpioni dell'Arcipelago Malese conservati nel Museo Civico di Storia Naturale di Genova. *Annali del Museo Civico di Storia Naturale di Genova*, 26: 327–428.
- Thorell, T. 1889. Viaggio di Leonardo Fea in Birmania e regioni vicine. XXI. Aracnidi Artrogastri Birmani raccolti da L. Fea nel 1885–1887. *Annali del Museo Civico di Storia Naturale di Genova*, 27: 521–729.
- Thorell, T. 1890. Studi sui ragni Malesi e Papuani. Part IV, 1. *Annali del Museo Civico di Storia Naturale di Genova*, 28: 1–419.
- Thorell, T. 1891. Spindlar från Nikobarerna och andra delar af södra Asien. *Bihang till Kongl. Svenska Vetenskaps-Akademiens Handlingar*, 24: 149 pp.
- Thorell, T. & Lindström, G. 1884. Discovery of a Silurian fossil scorpion. *The Glasgow Herald*, Dec. 19, 1884.
- Thorell, T. & Lindström, G. 1885. On a Silurian scorpion from Gotland. *Bihang till Kongl. Svenska Vetenskaps-Akademiens Handlingar*, 21(9): 1–33.
- Tollerton, V. P., Jr. 1989. Morphology, taxonomy, and classification of the order Eurypterida Burmeister, 1843. *Journal of Paleontology*, 63: 642–657.
- Trägårdh, I. 1902. Beiträge zur Kenntnis der schwedischen Acaridenfauna. *Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar*, 28: 1–26.

- Trägårdh, I. 1915. Bidrag till kännedomen om spinnkvalstren (*Tetranychus* Duf.). *Centralanstalten för försöksväsendet på jordbruksområdet. Entomologiska avdelningen*, 20: 1–60.
- Trägårdh, I. 1946. Outlines of a new classification of the Mesostigmata (Acarina) based on comparative morphological data. *Lunds Universitets Arsskrift, N.F.* 42: ?-?
- Trägårdh, I. 1950. Description of a new species of *Heterocheylus* Lombardini from Africa, with notes on the classification of the Pseudocheyletidae. *Entomologisk tidskrift*, 71: 104–110.
- Travé, J. 1959. Sur le genre *Niphocepheus* Balogh 1943. Les Niphocepheidae, famille nouvelle (Acariens, Oribates). *Acarologia*, 1: 475–498.
- Travé, J. 1967. *Phyllochthonius aoutii* nov. gen., nov. spec., un Enarthronota (Acarien, Oribate) nouveau de Côte d'Ivoire, avec la création d'une superfamille nouvelle, Phyllochthonoidea. *Zoologische Mededelingen*, 42: 83–105.
- Treat, A. E. 1955. An ectoparasite (Acarina: Mesostigmata) from moths of the genus *Zale*. *Journal of Parasitology* 41: 555–561.
- Türk, E. 1963. A new tyroglyphid deutonymph in amber from Chiapas, Mexico. *University of California Publications in Entomology* 31: 49–51.
- Ubick, D. & Dunlop, J. A. 2005. On the placement of the Baltic amber harvestman *Gonyleptes nemastomoides* Koch & Berendt, 1854, with notes on the phylogeny of Cladonychiidae (Opiliones, Laniatores, Travunioidea). *Mitteilungen aus dem Musuem für Naturkunde Berlin, Geowissenschaftliche Reihe* 8: 75–82.
- Vachon, M. & Heyler, D. 1985. Description d'une nouvelle espèce de Scorpion: *Buthiscorpius pescei* (Stéphanien de Montceau-les-Mines, France). Remarques sur la classification des Scorpions (Arachnida) du Carbonifère. *Bulletin de la Société d'Histoire Naturelle d'Autun* 113: 29–47.
- Vandenbergh, A. 1960. *Pringlia demaisteri* nov. sp., un xiphosure (Chélicératé) du Stéphanien de la Loire. – *Bulletin de la Société géologique de France* 7: 687–689.
- Vercammen-Grandjean, P. H. 1973. Study of the “Erythraeidae, R.O.M. No. 8” of Ewing, 1937. 329–335. In Daniel, M. and Rosicky, B. (eds). *Proceedings of the 3rd International Congress of Acarology*. Academia, Prague, 837 pp.
- Via Boada, L. & Villalta, J. F. de 1966. *Hetrolimulus gadeai*, nov. gen., nov. sp., représentant d'une nouvelle famille de Limulacés dans le Trias d'Espagne. *Comtes Rendues Sommaire Séances Société Géologique France*, 1966: 57–59.
- Viets, K. O. 1978. New water mites (Hydrachnellae: Acari) from Australia. *Australian Journal of Marine and Freshwater Research*, 29: 77–92.
- Villalta, J. F. 1957. Dos zoocecidas fósiles del Mioceno de Cerdaña (prov. de Lérida). *Cursillos y conferencias del Instituto Lucas Mallada*, 4: 63–64.

- Vitzthum, H. Graf 1931. Acari=Milben. In Kukenthal, W. (ed.) *Handbuch der Zoologie*, Vol. III 2. 3. Walter de Gruyter & Co., Berlin, pp. 1–160.
- Vitzthum, H. G. 1942. Acarina. In Brönn's *Klassen und Ordnungen des Tierreiches*, IV. Abt., 5. Buch, 5. Lieferung (1942), Leipzig, Akademische Verlagsgesellschaft Becker u. Erler: pp. 641–800.
- Wagner, W. A. 1887. Copulations organes des Männchens als Criterium für die Systematik der Spinnen. *Horae Societatis Entomologicae Rossicae*, 22: 3-132.
- Walcott, C. D. 1882. Description of a new genus of the order Eurypterida from the Utica Slate. *American Journal of Science*, 3rd Series, 23: 213–216.
- Walckenaer, C. A. 1802. *Faune parisienne. Insectes. Ou Histoire abrégée des Insectes des environs de Paris*. Paris, 2: 187–250.
- Walckenaer, C. A. 1805. *Tableau des Aranéides ou Caractères essentials des tribus, genres, familles et races que renferme le genre Aranea de Linné, avec la désignation des espèces comprises dans chacune de ces divisions*. Paris, 88 pp.
- Walckenaer, C. A. 1826. Aranéides. In *Faune française...*, Paris: 96 pp.
- Walckenaer, C. A. 1837. *Histoire naturelle des insects. Aptères*. Vol. 1. Librairie Encyclopédique de Roret, Paris, 682 pp.
- Walker, N. A. 1965. Euphthiracaroidea of California Sequoia litter : with a reclassification of the families and genera of the world (Acarina: Oribatei). *Fort Hays Studies, New Series, Science Series*, 3: 154 pp.
- Walossek, D., Li, C.S. & Brauckmann, C. 1990. A scorpion from the Upper Devonian of Hubei Province, China (Arachnida, Scorpiones). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1990(3): 169–180.
- Waloszek, D. & Dunlop, J. A. 2002. A larval sea spider (Arthropoda: Pycnogonida) from the Upper Cambrian 'Orsten' of Sweden and the phylogenetic position of pycnogonids. *Palaeontology*, 45: 421–446.
- Walter, D. E. 1997. Heatherellidae – a new family of Mesostigmata (Acari: Parasitiformes) based on two new species from rainforest litter in Australia. *International Journal of Acarology*, 23: 167–175.
- Walter, D. E. 2000. A jumping mesostigmatan mite, *Saltiseius hunteri* n. g., n. sp. (Acari: Mesostigmata: Trigynaspida: Saltiseiidae, n. fam.) from Australia. *International Journal of Acarology*, 26: 25–31.
- Walter, D. E. & Gerson, U. 1998. Dasythyreidae, new family, and *Xanthodasythyreus* n. g. (Acari: Prostigmata: Raphignathoidea) from Australia. *International Journal of Acarology*, 24: 189–197.
- Walter, D. E. & Krantz, G. W. 1999. New early derivative mesostigmatans from Australia: *Nothogynus* n. g., Nothogynidae n. fam. (Mesostigmata: Microgyniina). *International Journal of Acarology*, 25: 67–76.
- Waterston, C. D. 1962. *Pagea sturrocki* gen. et sp. nov., a new eurypterid from the Old Red Sandstone of Scotland. *Palaeontology*, 5: 137–148.
- Waterston, C. D. 1964. Observations on pterygotid eurypterids. *Transactions of the Royal Society of Edinburgh*, 66: 9–33.

- Waterston, C. D. 1968. Further observations on the Scottish Carboniferous eurypterids. *Transactions of the Royal Society of Edinburgh*, 68: 1–20.
- Waterston, C. D. 1979. Problems of functional morphology and classification in stylonurid eurypterids (Chelicerata, Merostomata), with observations on the Scottish Styloonoidea. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 70: 251–322.
- Waterston, C. D. 1985. Chelicerata from the Dinantian of Fouldon, Berwickshire, Scotland. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 76: 25–33.
- Waterston, C. D., Oelofsen, B. W. and Ooshuizen, R. D. F. 1985. *Cyrtocetus wittebergensis* sp. nov. (Chelicerata: Eurypterida), a large sweep-feeder from the Carboniferous of South Africa. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 76: 339–358.
- Watson, D. M. S. 1909. *Limulus woodwardi*, sp. nov., from the Lower Oolite of England. *Geological Magazine, New Series*, (5) 6: 14–15.
- Waterlot, G. 1934. *Étude de la Faune continentale du Terrain houiller Sarro-Lorrain – Études des gîtes minéraux de la France. Bassin houiller de la Sarre et de la Lorraine II. Faune fossile*. Lille, 317 pp.
- Weidner, H. 1964. Eine Zecke, *Ixodes succineus* sp. n. im Batischen Bernstein. *Veöffentlichungen aus dem Überseemuseum Bremen*, 3: 143–151.
- Weitschat, W. & Wichard, W. 2002. *Atlas of plants and animals in Baltic amber*. Dr. F. Pfeil, Munich, 256 pp.
- Westring, N. 1851. Förteckning öfver de till närvarande tid Kände, i Sverige förekommande Spindlarter, utgörande ett antal af 253, deraf 132 äro nya för svenska Faunan. *Göteborgs Kungliga Vetenskaps- och Vitterhets-Samhällens handlingar*, 2: 25–62.
- Westwood, J. O. 1835. Insectorum Arachnoidumque novorum Decades duo. *The Zoological Journal, London*, 5: 440–453.
- Westwood, J. O. 1874. *Thesaurus entomologicus oxoniensis*. Clarendon Press, Oxford, xx pp.
- Weyenbergh, H., Jr 1874. Notes sur quelques insectes du calcaire jurassique de la Bavière. *Archives Musée Teyler, Haarlem*, 3: 234–236.
- Weygoldt, P. 1996. Evolutionary morphology of whip spiders: towards a phylogenetic system (Chelicerata: Arachnida: Amblypygi). *Journal of Zoological Systematics and Evolutionary Research*, 34: 185–202.
- Weygoldt, P. & Paulus, H.F. 1979. Untersuchungen zur Morphologie, Taxonomie und Phylogenie der Chelicerata. *Zeitschrift für zoologische Systematik und Evolutionsforschung*, 17: 85–115, 177–200.
- White, D. 1908. Report on the fossil flora of the Coal Measures of Brazil. 377–607. In White, J. C. (ed.). *Final report on the coal measures and associated rocks of South Brazil*. Comissão de Estudos das Minas de Carvão de Pedra Do Brazil, Rio de Janeiro.
- Whiteaves, J. F. 1884. On some new, imperfectly characterized or previously unrecorded species of fossils from the Guelph Formations of Ontario. *Palaeozoic Fossils of Canada*, 3(1):1–43

- Whitfield, R. P. 1882. Descriptions of new species of fossils from Ohio, with remarks on some of the geological formations in which they occur. *Annals of the New York Academy of Science*, 2: 193–244.
- Whitfield, R. P. 1885a. An American Silurian scorpion. *Science*, 6: 87–88.
- Whitfield, R. P. 1885b. On a fossil scorpion from the Silurian rocks of America. *Bulletin of the American Museum of Natural History*, 1(9): 181–190.
- Wiles, P. R. 1996. A new family, genus and species of watermite (Acari: Hydrachnidia, Lebertioidea) from Brunei. *Quekett Journal of Microscopy*, 37: 692–695.
- Williams, H. 1915. An eurypterid horizon in the Niagara Formation of Ontario. *Geological Survey of Canada, Museum Bulletin*, 20: 1–9.
- Willmann, C. 1931. Oribatei (Acari), gesammelt von der Deutschen Limnologischen Sunda-Expedition. *Archiv für Hydrobiologie*, Supplement-Band IX: 240–305.
- Wills, L. J. 1910. On the fossiliferous Lower Keuper rocks of Worcestershire, with descriptions of some of the animals discovered therein. *Proceedings of the Geologists' Association*, 21: 249–331.
- Wills, L. J. 1947. *A monograph of the British Triassic scorpions*. The Palaeontographical Society, London, 100 & 101: 137 pp.
- Wills, L. J. 1959. The external anatomy of some Carboniferous “scorpions” Part 1. *Palaeontology*, 1: 261–282.
- Wills, L. J. 1960. The external anatomy of some Carboniferous “scorpions”. Part 2. *Palaeontology*, 3: 276–332.
- Wilson, E. B. 1878. Descriptions of two new genera of Pycnogonida. *American Journal of Science*, 15: 200–203.
- With, C. J. 1902. A new acaride *Opilioacarus segmentatus*. *Comptes Rendus du Congrès des Naturalistes et Médecins du Nord*, 20: 4–5.
- With, C. J. 1906. The Danish expedition to Siam 1899–1900. III. Chelonethi. An account of the Indian false-scorpions together with studies on the anatomy and classification of the order. *Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger*, 7(3): 1–214.
- Witaliński, W. 2000. *Aclerogamasus stenocornis* sp. n., a fossil mite from the Baltic amber (Acari: Gamasida: Parasitidae). *Genus*, 11: 619–626.
- Wolff, R.J. 1990. A new species of *Thiodina* (Araneae: Salticidae) from Dominican amber. *Acta Zoologica Fennica*, 190: 405–408.
- Womersley, H. 1956. On some new Acarina-Mesostigmata from Australia, New Zealand and New Guinea. *Zoological Journal of the Linnean Society of London*, 42: 505–599.
- Womersley, H. 1957. A fossil mite (*Acronothrus ramus* n.sp.) from Cainozoic resin at Allendale, Victoria. *Proceedings of the Royal Society of Victoria* 69: 21–23.
- Wood, T. G. 1969. The Homocaligidae, a new family of mites (Acari: Raphignathoidea), including a description of a new species from Malaya and the British Solomon Islands. *Acarologia*, 11: 711–729.
- Woodward, H. 1865. On a new genus of Eurypterida from the Lower Ludlow rock of Leintwardine, Shropshire. *Quarterly Journal of the Geological Society of London*, 21: 490–492.

- Woodward, H. 1868a. On a new limuloid crustacean (*Neolimulus falcatus*) from the Upper Silurian of Lesmahagow, Lanarkshire. *Geological Magazine*, 5: 1–3.
- Woodward, H. 1870. On *Necrogammarus salweyi* (H. Woodward), an amphipodus crustacean from the Lower Ludlow of Leintwardine. *Transactions of the Woolhope Naturalists Field Club*, 1870: 271–272.
- Woodward, H. 1871a. On the remains of a giant isopod *Praearcturus gigas*, (H. Woodward) from the Old Red Sandstone of Rowlestone Quarry, Herefordshire. *Transactions of the Woolhope Field Naturalist's Club*, 1871: 266–270.
- Woodward, H. 1871b. On the discovery of a new and very perfect Arachnid from the ironstone of the Dudley Coal-field. *Geological Magazine*, 8: 385–388.
- Woodward, H. 1872a. Notes on some British Palaeozoic Crustacea belonging to the order Merostomata. *Geological Magazine*, 9: 433–441.
- Woodward, H. 1872b. On a new Arachnid from the Coal-measures of Lancashire. *Geological Magazine*, 9: 385–387.
- Woodward, H. 1876. On the discovery of a fossil scorpion in the British Coal Measures. *Quarterly Journal of the Geological Society of London* 32: 57–59.
- Woodward, H. 1878b. Discovery of the remains of a fossil crab (Decapoda-Bracyura) in the Coal Measures of the Environs of Mons, Belgium. *Geological Magazine, new series, Decade 2*, 5: 433–436.
- Woodward, H. 1879. Contributions to the knowledge of fossil Crustacea. *Quarterly Journal of the Geological Society London*, 35: 549–555.
- Woodward, H. 1887. On a new species of *Eurypterus* from the Lower Carboniferous shales of Glencarholm, Eskdale, Scotland. *Geological Magazine, Decade 3*, 4: 481–484.
- Woodward, H. 1888. Note on *Eurypterus* from the Carboniferous. *Geological Magazine, Decade 3*, 5: 419–421.
- Woodward, H. 1907a. Two new species of *Eurypterus* from the Coal-Measures of Ilkeston, Derbyshire. *Geological Magazine*, 4: 277–282.
- Woodward, H. 1907b. Further notes on the Arthropoda of the British Coal Measures. *Geological Magazine*, 4: 539–549.
- Woodward, H. 1918. Fossil arthropods from the Carboniferous rocks of Cape Breton, Nova Scotia; and from the Upper Coal Measures, Sunderland, England. *Geological Magazine*, 5: 462–471.
- Woolley, T. A. 1969. Two new species of *Hydrozetes*, extant and fossil (Acari: Cryptostigmata, Hydrozetidae). *New York Entomological Society*, 77: 250–256.
- Woolley, T. A. 1971. Fossil oribatid mites in amber from Chiapas, Mexico (Acarina: Oribatei = Cryptostigmata). *University of California Publications in Entomology*, 63: 91–99.
- Woolley, T. A. & Higgins, H. G. 1968. Megeremaeidae: A New Family of Oribatid Mites (Acari: Cryptostigmata). *Great Basin Naturalist*, 28(4): 172–175.

- Wright, D. F. & Selden, P. A. 2011. A trigonotarbid arachnid from the Pennsylvanian of Kansas. *Journal of Paleontology*, 85: 871–876.
- Wunderlich, J. 1981. Fossile Zwergsechsaugenspinnen (Oonopidae) der Gattung *Orchestina* Simon, 1882 in Bernstein mit Anmerkungen zur Sexualbiologie (Arachnida: Araneae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 51: 83–113.
- Wunderlich, J. 1982. Die häufigsten Spinnen (Araneae) des Dominikanischen Bernsteins. *Neue Entomologische Nachrichten*, 1: 26–45.
- Wunderlich, J. 1985. Ein bisher unbekannte fossile Krabbenspinne aus dem Randecker Maar in Südwest-Deutschland (Arachnida: Araneae: Thomisidae). *Neue Entomologische Nachrichten*, 14: 4–13.
- Wunderlich, J. 1986. *Spinnenfauna Gestern und Heute. Fossile Spinnen in Bernstein und ihre heute lebenden Verwandten*. Erich Bauer Verlag bei Quelle und Meyer, Wiesbaden, 283 pp.
- Wunderlich, J. 1987. *Tama minor* n. sp., eine fossile Spinnenart der Familie Hersiliidae in Dominikanischem Bernstein (Arachnida: Araneae). *Entomologische Zeitschrift*, 97: 93–96.
- Wunderlich, J. 1988. Die fossilen Spinnen im dominikanischen Bernstein. *Beiträge zur Araneologie*, 2: 1–378.
- Wunderlich, J. 1991. Beschreibung der ersten fossilen Spinne der Familie Leptonetidae: *Eoleptona kutscheri* n. gen., n. sp. in Sächsischem Bernstein (Arachnida: Araneae). *Entomologische Zeitschrift*, 101: 21–26.
- Wunderlich, J. 1993a. Die ersten fossilen Speisspinnen (Fam. Scytodidae) im Baltischen Bernstein (Arachnida: Araneae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 75: 243–247.
- Wunderlich, J. 1993b. Die ersten fossilen Becherspinnen (Fam. Cyatholipidae) in Baltischem und Bitterfelder Bernstein (Arachnida: Araneae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 75: 231–241.
- Wunderlich, J. 1998. Beschreibung der ersten fossilen Spinnen der Unterfamilien Mysmeninae (Anapidae) und Erigoninae (Linyphiidae) im Dominikanischen Bernstein (Arachnida: Araneae). *Entomologische Zeitschrift*, 108: 363–367.
- Wunderlich, J. 2000. Zwei neue Arten der Familie Falltürspinnen (Araneae: Ctenizidae) aus dem Baltischen Bernstein. *Entomologische Zeitschrift*, 110: 345–348.
- Wunderlich, J. 2004a. Introduction, general findings and conclusions. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 5–329.
- Wunderlich, J. 2004b. The fossil mygalomorph spiders (Araneae) in Baltic and Dominican amber and about extant members of the family Micromygalidae. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 595–631.
- Wunderlich, J. 2004c. Fossil spiders (Araneae) of the superfamily Dysderoidea in Baltic and Dominican amber, with revised family diagnoses. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 633–746.

- Wunderlich, J. 2004d. Fossil and extant spiders (Araneae) of the superfamily Eresoidea s.l., with special reference to the Archaeidae and remarks on some higher taxa of the superfamily Araneoidea. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 747–808.
- Wunderlich, J. 2004e. On selected higher and lower taxa of fossil and extant spiders of the superfamily Oecobioidea, with a provisional Cladogram (Araneae: Hersiliidae and Oecobiidae). *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 809–848.
- Wunderlich, J. 2004f. Fossil spiders of the family Uloboridae (Araneae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 851–886.
- Wunderlich, J. 2004g. The fossil spiders of the family Deinopidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 887–897.
- Wunderlich, J. 2004h. The fossil spiders (Araneae) of the families Tetragnathidae and Zygelliidae n. stat. in Baltic and Dominican amber, with notes on higher extant and fossil taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 899–955.
- Wunderlich, J. 2004i. Fossil taxa of the family Araneidae (Araneae) inclusively Nephilinae in Baltic and Dominican amber, with the description of a new extinct subfamily and notes on selected extant taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 956–997.
- Wunderlich, J. 2004j. The fossil Theridiosomatidae (Araneae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 998–1019.
- Wunderlich, J. 2004k. The fossil spiders of the family Anapidae s. l. (Aeaneae [sic]) in Baltic, Dominican and Mexican amber and their extant relatives, with the description of a new subfamily Comarominae. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1020–1111.
- Wunderlich, J. 2004l. On the relationships of the families of the superfamily Araneoidea (Araneae) and their kin, with cladograms, remarks on the origin of the orb web and description of the new and extinct families Baltsuccinidae and Protheridiidae in Tertiary Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1112–1154.
- Wunderlich, J. 2004m. The fossil spiders (Araneae) of the family Cyatholipidae in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1155–1188.
- Wunderlich, J. 2004n. The fossil spiders (Araneae) of the family Synotaxidae in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1189–1239.
- Wunderlich, J. 2004o. Remarks on the fossil spiders (Araneae) of the family Nesticidae in amber, with the description of a new species in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1240–1244.
- Wunderlich, J. 2004p. Remarks on fossil spiders (Araneae) of the family Theridiidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1245–1248.

- Wunderlich, J. 2004q. Fossil pirate spiders (Araneae: Araneoidea: Mimetidae s. l.) in Baltic and Dominican amber, with notes on intrafamiliar higher taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1249–1278.
- Wunderlich, J. 2004r. Descriptions of the first fossil spiders (Araneae) of the family Pimoidae in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1279–1297.
- Wunderlich, J. 2004s. The fossil spiders of the family Linyphiidae in Baltic and Dominican amber (Araneae: Linyphiidae). *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1298–1373.
- Wunderlich, J. 2004t. No proof of fossil spiders (Araneae) of the family Psechridae in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1375–1376.
- Wunderlich, J. 2004u. Fossil spiders of the family Amaurobiidae (Arachnida: Araneae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1377–1379.
- Wunderlich, J. 2004v. Fossil spiders of the family Dictynidae s. l., including Cryphoecinae and Hahniinae in Baltic and Dominican amber and copal from Madagascar, and on selected extant Holarctic taxa, with new descriptions and diagnoses. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1380–1482.
- Wunderlich, J. 2004w. Fossil spiders (Araneae) of the family Agelenidae s. str. in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1483–1488.
- Wunderlich, J. 2004x. The fossil Zoropsidae in Baltic amber with revised diagnoses of the family Zoropsidae and its fossil and extant higher taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1489–1522.
- Wunderlich, J. 2004y. Spiders (Araneae) of the extinct family Insecutoridae Petrunkevitch 1942 in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1523–1531.
- Wunderlich, J. 2004z. Fossil spiders of the family Pisauridae (Araneae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1532–1541.
- Wunderlich, J. 2004aa. Members of the family Trechaleidae (Araneae) in Baltic and Dominican amber? *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1542–1553.
- Wunderlich, J. 2004ab. Fossil spiders (Araneae) of the family Oxyopidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1554–1556.
- Wunderlich, J. 2004ac. Proof of presence of the family Lycosidae (Araneae) in Baltic and Dominican amber? *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1557–1558.
- Wunderlich, J. 2004ad. Fossil spiders (Araneae) of the extinct family Ephalmatoridae Petrunkevitch 1950 in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1559–1577.
- Wunderlich, J. 2004ae. Fossil spiders (Araneae) of the family Zodariidae in Baltic amber, with remarks on their subfamilies including the Cryptothelinae and the Homalonychinae. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1578–1611.
- Wunderlich, J. 2004af. Fossil spiders (Araneae) of the families Clubionidae and Miturgidae (questionable) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1612–1622.

- Wunderlich, J. 2004ag. The fossil spiders of the family Liocranidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1623–1635.
- Wunderlich, J. 2004ah. Fossil spiders of the family Corinnidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1636–1680.
- Wunderlich, J. 2004ai. Fossil spiders (Araneae) of the family Gnaphosidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1681–1685.
- Wunderlich, J. 2004aj. Fossil spiders (Araneae) of the family Anyphaenidae in Baltic and Dominic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1686–1688.
- Wunderlich, J. 2004ak. Members of the family Philodromidae (Araneae) in Baltic amber? *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1689–1693.
- Wunderlich, J. 2004al. Fossil spiders (Araneae) of the family Sparassidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1694–1698.
- Wunderlich, J. 2004am. Fossil spiders of the family Trochanteriidae (Araneae) in Baltic, Dominican and Mexican amber, with a revision of the genus *Sosybius* Koch and Berendt 1854. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1699–1732.
- Wunderlich, J. 2004an. Fossil spiders of the family Selenopidae in Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1733–1736.
- Wunderlich, J. 2004ao. The new spider (Araneae) family Borboropactidae from the tropics and fossil in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1737–1746.
- Wunderlich, J. 2004ap. Fossil crab spiders (Araneae: Thomisidae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1747–1760.
- Wunderlich, J. 2004aq. Fossil jumping spiders (Araneae: Salticidae) in Baltic and Domican amber, with remarks on Salticidae subfamilies. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1761–1819.
- Wunderlich, J. 2004ar. Fossil spiders (Araneae) in Early Tertiary amber from the Ukraine. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1821–1829.
- Wunderlich, J. 2004as. Subrecent spiders (Araneae) in copal from Madagascar, with description of new species. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1830–1853.
- Wunderlich, J. 2004at. Two new fossil spider species in Copal from Colombia (Araneae: Oonopidae and Dictynidae). *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1854–1859.
- Wunderlich, J. 2004au. Description of two fossil taxa of spiders (Araneae: Oonopidae, Pholcidae) in Chinese amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1860–1863.
- Wunderlich, J. 2004av. Report on spider (Araneae) of the families Araneidae and Zygelliidae in Lebanese amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1864–1865.
- Wunderlich, J. 2006. *Spatiator martensi* n. sp., a second species of the extinct spider species Spatiatoriidae in Eocene Baltic amber. *Zootaxa*, 1325: 313–318.

- Wunderlich, J. 2008a. Descriptions of fossil spider (Araneae) taxa mainly in Baltic amber, as well as certain related extant taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 5: 44–139.
- Wunderlich, J. 2008b. On extant and fossil (Eocene) European comb-footed spiders (Araneae: Theridiidae), with notes on their subfamilies, and with descriptions of new taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 5: 140–469.
- Wunderlich, J. 2008c. On extant and fossil members of the RTA-clade in Eocene European ambers of the families Borboropactidae, Corinnidae, Selenopidae, Sparassidae, Trochanteriidae, Zoridae s. l., and of the superfamily Lycosoidea. *In* Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 5: 470–523.
- Wunderlich, J. 2008d. The dominance of ancient spider families of the Araneae: Haplogyne in the Cretaceous, and the late diversification of advanced ecribellate spiders of the Entelegynae after the Cretaceous–Tertiary boundary extinction events, with descriptions of new families. *In* Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 5: 524–675.
- Wunderlich, J. 2011a. On extant European spiders of the tribe Mangorini (Araneae: Araneidae) and two doubtful taxa in Baltic amber. *Beiträge zur Araneologie*, 6: 9–18.
- Wunderlich, J. 2011b. Taxonomy of extant and fossil (Eocene) European ground spiders of the family Gnaphosidae (Araneae), with a key to the genera, and descriptions of new taxa. *Beiträge zur Araneologie*, 6: 19–97.
- Wunderlich, J. 2011c. Spiders of the family Prodidomidae (Araneae) from Europe and Madagascar. *Beiträge zur Araneologie*, 6: 98–107.
- Wunderlich, J. 2011d. On extant and fossil (Eocene) Holarctic sac spiders (Araneae: Clubionidae), with descriptions of new taxa. *Beiträge zur Araneologie*, 6: 121–157.
- Wunderlich, J. 2011e. New extant taxa of the spider family Theridiosomatidae (Araneae) from Laos and on some fossil taxa. *Beiträge zur Araneologie*, 6: 427–444.
- Wunderlich, J. 2011f. Some subrecent spiders (Araneae) in copal from Madagascar. *Beiträge zur Araneologie*, 6: 445–460.
- Wunderlich, J. 2011g. Some fossil spiders in Dominican amber (Araneae: Hersiliidae, Theridiidae, Gnaphosidae). *Beiträge zur Araneologie*, 6: 461–471.
- Wunderlich, J. 2011h. Some fossil spiders (Araneae) in Eocene European ambers. *Beiträge zur Araneologie*, 6: 472–538.
- Wunderlich, J. 2011i. Some fossil spiders (Araneae) in Cretaceous ambers. *Beiträge zur Araneologie*, 6: 539–557.
- Wunderlich, J. 2012a. New subrecent species in copal from Madagascar, and on the relationships of the Copaldictyninae Wunderlich 2004 (Araneae: Linyphiidae, Theridiidae, Dictynidae, and Titanoecidae). *In* Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 75–88.

- Wunderlich, J. 2012b. New fossil spiders (Araneae) in Eocene amber from the Ukraine. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 89–93.
- Wunderlich, J. 2012c. New fossil spiders (Araneae) of eight families in Eocene Baltic amber, and revisions of selected taxa. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 94–149.
- Wunderlich, J. 2012d. On the fossil spider (Araneae) fauna in Cretaceous ambers, with descriptions of new taxa from Myanmar (Burma) and Jordan, and on the relationships of the superfamily Leptonetoidea. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 157–232.
- Wunderlich, J. 2012e. Description of the first fossil Ricinulei in amber from Burma (Myanmar), the first report of this arachnid order from the Mesozoic and from Asia, with notes on the related extinct order Trigonotarbida. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 233–244.
- Wunderlich, J. 2012f. Corrections and addenda to vol. 6 of the Beitr. Araneol. (2011). In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 245–246.
- Wunderlich, J. & Milki, R. 2004. Description of the extinct new subfamily Microsegstriinae (Araneae: Segestriidae) in Cretaceous Lebanese Amber. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1867–1873.
- Żabka, M. 1988. Fossil Eocene Salticidae (Araneae) from the collection of the Museum of Earth in Warsaw. *Annales Zoologici*, 41: 415–420.
- Zacharda, M. 1979. Strandtmanniidae – a new family of Eupodoidea (Acarina : Prostigmata). *Vestník Československé Společnosti Zoologické*, 43: 76–81.
- Zacharda, M. & Krivoluckij, D. A. 1985. Prostigmatic mites (Acarina: Prostigmata) from the Upper Cretaceous and Paleogene amber of the USSR. *Vestník Československé Společnosti Zoologické*, 49: 147–152.
- Zachvatkin, A. A. 1952. [The division of the Acarina into orders and their position in the system of the Chelicerata.] *Parazitologicheskii Sbornik Zoologicheskii Institut Akademii Nauk SSSR*, 14: 5–46. [in Russian]
- Zapfe, H. 1955. Filogenia y función en *Austrochilus manni* Gertsch y Zapfe (Araneae-Hypochilidae). *Trabajos del Laboratorio de Zoología de la Universidad de Chile*, 2: 1–53.
- Zhang, J., Sun, B. & Zhang, X. 1994. *Miocene insects and spiders from Shanwang, Shandong*. Science Press, Beijing, 298 pp. [in Chinese with English Summary].
- Zhang, Q.-y., Hu, S.-x., Zhou, C.-y., Lv, T. & Bai, J.-k. (2009): [New occurrence of Xiphosura in China.] *Progress in Nature Science*, 19: 1090–1093. [in Chinese]
- Zhang, Z.-Q. 1998: An unusual early-derivative larva of Parasitengona (Acari: Prostigmata) and proposal of a new superfamily. *Systematic & applied acarology*, 3: 159–170.
- Zhang, Z.-Q. & Fan, Q.-H. 2007. Allotanaupodidae, a new family of early derivative Parasitengona (Acari: Prostigmata). *Zootaxa*, 1517: 1–52.

- Zinken, C. 1862. *Limulus Decheni* aus dem Braunkohlensandstein bei Teuchern. *Zeitschrift für die Gesammten Naturwissenschaften*, 19: 329–331.
- Zittel, K. A. 1885. *Handbuch der Palaeontologie. I. Abtheilung, Palaeozoologie, 2 [Mollusca und Arthropoda]*. R, Oldenbourg, München, Leipzig, 893 pp.
- Zittel, K. A. & Eastman, C. R. 1913. *Textbook of Palaeontology (2nd Ed.) 1*. Macmillan, London, 839 pp.